

Edgar Dahl

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6434100/edgar-dahl-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

99
papers

4,750
citations

40
h-index

67
g-index

112
ext. papers

5,299
ext. citations

6.2
avg, IF

4.87
L-index

#	Paper	IF	Citations
99	Pax genes and organogenesis. <i>BioEssays</i> , 1997 , 19, 755-65	4.1	316
98	CD24 is expressed in ovarian cancer and is a new independent prognostic marker of patient survival. <i>American Journal of Pathology</i> , 2002 , 161, 1215-21	5.8	211
97	CD24 expression is a new prognostic marker in breast cancer. <i>Clinical Cancer Research</i> , 2003 , 9, 4906-13	12.9	191
96	Molecular cloning and functional expression of mouse connexin-30, a gap junction gene highly expressed in adult brain and skin. <i>Journal of Biological Chemistry</i> , 1996 , 271, 17903-10	5.4	153
95	Expression pattern of connexin gene products at the early developmental stages of the mouse cardiovascular system. <i>Circulation Research</i> , 1997 , 81, 423-37	15.7	146
94	Expression profiling of microdissected matched prostate cancer samples reveals CD166/MEMD and CD24 as new prognostic markers for patient survival. <i>Journal of Pathology</i> , 2005 , 205, 359-76	9.4	144
93	The death-domain fold of the ASC PYRIN domain, presenting a basis for PYRIN/PYRIN recognition. <i>Journal of Molecular Biology</i> , 2003 , 332, 1155-63	6.5	128
92	Deletions of chromosome 8p and loss of sFRP1 expression are progression markers of papillary bladder cancer. <i>Laboratory Investigation</i> , 2004 , 84, 465-78	5.9	125
91	Molecular profiling of laser-microdissected matched tumor and normal breast tissue identifies karyopherin alpha2 as a potential novel prognostic marker in breast cancer. <i>Clinical Cancer Research</i> , 2006 , 12, 3950-60	12.9	124
90	Frequent expression loss of Inter-alpha-trypsin inhibitor heavy chain (ITI-H) genes in multiple human solid tumors: a systematic expression analysis. <i>BMC Cancer</i> , 2008 , 8, 25	4.8	121
89	Frequent loss of SFRP1 expression in multiple human solid tumours: association with aberrant promoter methylation in renal cell carcinoma. <i>Oncogene</i> , 2007 , 26, 5680-91	9.2	118
88	Developmental regulation of connexin 40 gene expression in mouse heart correlates with the differentiation of the conduction system. <i>Developmental Dynamics</i> , 1995 , 204, 358-71	2.9	115
87	Tight correlation between expression of the Forkhead transcription factor FOXM1 and HER2 in human breast cancer. <i>BMC Cancer</i> , 2008 , 8, 42	4.8	114
86	Exhaustive mining of EST libraries for genes differentially expressed in normal and tumour tissues. <i>Nucleic Acids Research</i> , 1999 , 27, 4251-60	20.1	114
85	Targeting the Wnt pathway in cancer: the emerging role of Dickkopf-3. <i>Biochimica Et Biophysica Acta: Reviews on Cancer</i> , 2012 , 1825, 18-28	11.2	97
84	Dual role of macrophage migration inhibitory factor (MIF) in human breast cancer. <i>BMC Cancer</i> , 2009 , 9, 230	4.8	96
83	Expression of the glioma-associated oncogene homolog (GLI) 1 in human breast cancer is associated with unfavourable overall survival. <i>BMC Cancer</i> , 2009 , 9, 298	4.8	95

82	Promoter hypermethylation of the tumor-suppressor genes ITIH5, DKK3, and RASSF1A as novel biomarkers for blood-based breast cancer screening. <i>Breast Cancer Research</i> , 2013 , 15, R4	8.3	92
81	Prognostic relevance of AGR2 expression in breast cancer. <i>Clinical Cancer Research</i> , 2006 , 12, 1728-34	12.9	86
80	The DAPIN family: a novel domain links apoptotic and interferon response proteins. <i>Trends in Biochemical Sciences</i> , 2001 , 26, 83-5	10.3	86
79	The ubiquitin-like molecule interferon-stimulated gene 15 (ISG15) is a potential prognostic marker in human breast cancer. <i>Breast Cancer Research</i> , 2008 , 10, R58	8.3	80
78	Epigenetic inactivation of the secreted frizzled-related protein-5 (SFRP5) gene in human breast cancer is associated with unfavorable prognosis. <i>Carcinogenesis</i> , 2008 , 29, 991-8	4.6	80
77	Loss of SFRP1 is associated with breast cancer progression and poor prognosis in early stage tumors. <i>International Journal of Oncology</i> , 2004 , 25, 641-9	1	80
76	Decreased expression of angiogenesis antagonist EFEMP1 in sporadic breast cancer is caused by aberrant promoter methylation and points to an impact of EFEMP1 as molecular biomarker. <i>International Journal of Cancer</i> , 2009 , 124, 1727-35	7.5	77
75	Activation of Wnt signalling in stroma from pancreatic cancer identified by gene expression profiling. <i>Journal of Cellular and Molecular Medicine</i> , 2008 , 12, 2823-35	5.6	72
74	Wnt signalling in human breast cancer: expression of the putative Wnt inhibitor Dickkopf-3 (DKK3) is frequently suppressed by promoter hypermethylation in mammary tumours. <i>Breast Cancer Research</i> , 2008 , 10, R82	8.3	70
73	Prognostic relevance of Wnt-inhibitory factor-1 (WIF1) and Dickkopf-3 (DKK3) promoter methylation in human breast cancer. <i>BMC Cancer</i> , 2009 , 9, 217	4.8	69
72	Promoter methylation-associated loss of ID4 expression is a marker of tumour recurrence in human breast cancer. <i>BMC Cancer</i> , 2008 , 8, 154	4.8	67
71	Promoter hypermethylation of the SFRP2 gene is a high-frequent alteration and tumor-specific epigenetic marker in human breast cancer. <i>Molecular Cancer</i> , 2008 , 7, 83	42.1	63
70	Nuclear karyopherin alpha2 expression predicts poor survival in patients with advanced breast cancer irrespective of treatment intensity. <i>International Journal of Cancer</i> , 2008 , 123, 1433-8	7.5	63
69	Expression levels of the putative zinc transporter LIV-1 are associated with a better outcome of breast cancer patients. <i>International Journal of Cancer</i> , 2005 , 117, 961-73	7.5	62
68	The fractalkine receptor CX3CR1 is involved in liver fibrosis due to chronic hepatitis C infection. <i>Journal of Hepatology</i> , 2008 , 48, 208-15	13.4	58
67	Expression analysis of mammaglobin A (SCGB2A2) and lipophilin B (SCGB1D2) in more than 300 human tumors and matching normal tissues reveals their co-expression in gynecologic malignancies. <i>BMC Cancer</i> , 2006 , 6, 88	4.8	55
66	ITIH5, a novel member of the inter-alpha-trypsin inhibitor heavy chain family is downregulated in breast cancer. <i>Cancer Letters</i> , 2004 , 204, 69-77	9.9	54
65	A systematic comparison of quantitative high-resolution DNA methylation analysis and methylation-specific PCR. <i>Epigenetics</i> , 2012 , 7, 772-80	5.7	52

64	Liquid biopsy in colon cancer: comparison of different circulating DNA extraction systems following absolute quantification of mutations using Intplex allele-specific PCR. <i>Oncotarget</i> , 2017 , 8, 86253-86263 ^{3,3}	4.9	49
63	Nuclear detection of Y-box protein-1 (YB-1) closely associates with progesterone receptor negativity and is a strong adverse survival factor in human breast cancer. <i>BMC Cancer</i> , 2009 , 9, 410	4.8	46
62	Systematic identification and molecular characterization of genes differentially expressed in breast and ovarian cancer. <i>Journal of Pathology</i> , 2005 , 205, 21-8	9.4	45
61	KPNA2 protein expression in invasive breast carcinoma and matched peritumoral ductal carcinoma in situ. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2007 , 451, 877-81	5.1	42
60	Expression analysis and RNA localization of PAI-RBP1 (SERBP1) in epithelial ovarian cancer: association with tumor progression. <i>Gynecologic Oncology</i> , 2007 , 107, 266-73	4.9	41
59	The Bayesian group-Lasso for analyzing contingency tables 2009 ,		38
58	Frequent loss of endothelin-3 (EDN3) expression due to epigenetic inactivation in human breast cancer. <i>Breast Cancer Research</i> , 2009 , 11, R34	8.3	38
57	Endogenous myoglobin in breast cancer is hypoxia-inducible by alternative transcription and functions to impair mitochondrial activity: a role in tumor suppression?. <i>Journal of Biological Chemistry</i> , 2011 , 286, 43417-28	5.4	34
56	A new alternatively spliced transcript of the mouse connexin32 gene is expressed in embryonic stem cells, oocytes, and liver. <i>Experimental Cell Research</i> , 2001 , 266, 177-86	4.2	33
55	Low Input Whole-Exome Sequencing to Determine the Representation of the Tumor Exome in Circulating DNA of Non-Small Cell Lung Cancer Patients. <i>PLoS ONE</i> , 2016 , 11, e0161012	3.7	32
54	Epigenetic inactivation of ST6GAL1 in human bladder cancer. <i>BMC Cancer</i> , 2014 , 14, 901	4.8	31
53	Systematic characterisation of GABRP expression in sporadic breast cancer and normal breast tissue. <i>International Journal of Cancer</i> , 2006 , 118, 1453-9	7.5	31
52	AGR3 in breast cancer: prognostic impact and suitable serum-based biomarker for early cancer detection. <i>PLoS ONE</i> , 2015 , 10, e0122106	3.7	30
51	Epigenetic inactivation of ITIH5 promotes bladder cancer progression and predicts early relapse of pT1 high-grade urothelial tumours. <i>Carcinogenesis</i> , 2014 , 35, 727-36	4.6	28
50	Systematic isolation of genes differentially expressed in normal and cancerous tissue of the pancreas. <i>Pancreatology</i> , 2003 , 3, 169-78	3.8	27
49	Chromosomal assignments of mouse connexin genes, coding for gap junctional proteins, by somatic cell hybridization. <i>Somatic Cell and Molecular Genetics</i> , 1992 , 18, 351-9		25
48	Epigenetic inactivation of the novel candidate tumor suppressor gene ITIH5 in colon cancer predicts unfavorable overall survival in the CpG island methylator phenotype. <i>Epigenetics</i> , 2014 , 9, 1290-301	5.7	24
47	Low expression of ITIH5 in adenocarcinoma of the lung is associated with unfavorable patientsU outcome. <i>Epigenetics</i> , 2015 , 10, 903-12	5.7	23

46	Production and characterisation of monoclonal antibodies against RAI3 and its expression in human breast cancer. <i>BMC Cancer</i> , 2009 , 9, 200	4.8	23
45	Overexpression of SERBP1 (Plasminogen activator inhibitor 1 RNA binding protein) in human breast cancer is correlated with favourable prognosis. <i>BMC Cancer</i> , 2012 , 12, 597	4.8	22
44	Promoter methylation of DNA damage repair (DDR) genes in human tumor entities: / is almost exclusively methylated in bladder cancer. <i>Clinical Epigenetics</i> , 2018 , 10, 15	7.7	20
43	Altered expression pattern of topoisomerase IIalpha in ovarian tumor epithelial and stromal cells after platinum-based chemotherapy. <i>Neoplasia</i> , 2006 , 8, 38-45	6.4	20
42	ACE2 polymorphism and susceptibility for SARS-CoV-2 infection and severity of COVID-19. <i>Pharmacogenetics and Genomics</i> , 2021 , 31, 165-171	1.9	20
41	Loss of Dickkopf 3 Promotes the Tumorigenesis of Basal Breast Cancer. <i>PLoS ONE</i> , 2016 , 11, e0160077	3.7	19
40	Aberrant DNA hypermethylation of the ITIH5 tumor suppressor gene in acute myeloid leukemia. <i>Clinical Epigenetics</i> , 2011 , 2, 419-23	7.7	17
39	Different structural organization of the encephalopsin gene in man and mouse. <i>Gene</i> , 2002 , 295, 27-32	3.8	17
38	BDNF is associated with SFRP1 expression in luminal and basal-like breast cancer cell lines and primary breast cancer tissues: a novel role in tumor suppression?. <i>PLoS ONE</i> , 2014 , 9, e102558	3.7	17
37	Epigenetic loss of putative tumor suppressor SFRP3 correlates with poor prognosis of lung adenocarcinoma patients. <i>Epigenetics</i> , 2018 , 13, 214-227	5.7	16
36	OASIS/CREB3L1 is epigenetically silenced in human bladder cancer facilitating tumor cell spreading and migration in vitro. <i>Epigenetics</i> , 2014 , 9, 1626-40	5.7	16
35	Enhancer of the rudimentary gene homologue (ERH) expression pattern in sporadic human breast cancer and normal breast tissue. <i>BMC Cancer</i> , 2008 , 8, 145	4.8	16
34	ITIH5 mediates epigenetic reprogramming of breast cancer cells. <i>Molecular Cancer</i> , 2017 , 16, 44	42.1	15
33	Loss of anterior gradient-2 expression is an independent prognostic factor in colorectal carcinomas. <i>European Journal of Cancer</i> , 2014 , 50, 1722-1730	7.5	15
32	Inter- α -trypsin inhibitor heavy chain 5 (ITIH5) is overexpressed in inflammatory skin diseases and affects epidermal morphology in constitutive knockout mice and murine 3D skin models. <i>Experimental Dermatology</i> , 2015 , 24, 663-8	4	15
31	Multisystemic Cellular Tropism of SARS-CoV-2 in Autopsies of COVID-19 Patients. <i>Cells</i> , 2021 , 10,	7.9	12
30	ITIH5 induces a shift in TGF- β superfamily signaling involving Endoglin and reduces risk for breast cancer metastasis and tumor death. <i>Molecular Carcinogenesis</i> , 2018 , 57, 167-181	5	11
29	SNiPER: a novel hypermethylation biomarker panel for liquid biopsy based early breast cancer detection. <i>Oncotarget</i> , 2019 , 10, 6494-6508	3.3	11

28	Expression and localization of e-cadherin in epithelial ovarian cancer. <i>Anticancer Research</i> , 2010 , 30, 2525-30	11
27	Oncogenic features of neuromedin U in breast cancer are associated with NMUR2 expression involving crosstalk with members of the WNT signaling pathway. <i>Oncotarget</i> , 2017 , 8, 36246-36265	3-3 9
26	Distinct secreted Frizzled receptor protein 1 staining pattern in patients with hyperplastic polyposis coli syndrome. <i>Archives of Pathology and Laboratory Medicine</i> , 2004 , 128, 967-73	5 9
25	Gene expression analysis combined with functional genomics approach identifies ITIH5 as tumor suppressor gene in cervical carcinogenesis. <i>Molecular Carcinogenesis</i> , 2017 , 56, 1578-1589	5 8
24	The atypical cadherin Dachsous1 localizes to the base of the ciliary apparatus in airway epithelia. <i>Biochemical and Biophysical Research Communications</i> , 2016 , 473, 1177-1184	3-4 8
23	Paradox of sonic hedgehog (SHH) transcriptional regulation: Alternative transcription initiation overrides the effect of downstream promoter DNA methylation. <i>Epigenetics</i> , 2011 , 6, 465-77	5-7 8
22	Abundant NDRG2 Expression Is Associated with Aggressiveness and Unfavorable Patients Outcome in Basal-Like Breast Cancer. <i>PLoS ONE</i> , 2016 , 11, e0159073	3-7 8
21	and DNA Methylation Biomarker Test (EI-BLA) for Urine-Based Non-Invasive Detection of Bladder Cancer. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6-3 8
20	RNA expression analysis on formalin-fixed paraffin-embedded tissues in TMA format by RNA in situ hybridization. <i>Methods in Molecular Biology</i> , 2010 , 664, 135-50	1-4 8
19	Epigenetic Clocks Are Not Accelerated in COVID-19 Patients. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6-3 7
18	Mutation analysis of multiple pilomatricomas in a patient with myotonic dystrophy type 1 suggests a DM1-associated hypermutation phenotype. <i>PLoS ONE</i> , 2020 , 15, e0230003	3-7 6
17	Molecular Diagnostic Applications in Colorectal Cancer. <i>Microarrays (Basel, Switzerland)</i> , 2014 , 3, 168-79	6
16	Segment-specific expression of the gap junction gene connexin31 during hindbrain development. <i>Development Genes and Evolution</i> , 1997 , 207, 359-361	1-8 6
15	Deconvolution of cellular subsets in human tissue based on targeted DNA methylation analysis at individual CpG sites. <i>BMC Biology</i> , 2020 , 18, 178	7-3 6
14	Elevated serum SDMA and ADMA at hospital admission predict in-hospital mortality of COVID-19 patients. <i>Scientific Reports</i> , 2021 , 11, 9895	4-9 6
13	Infinite mixture-of-experts model for sparse survival regression with application to breast cancer. <i>BMC Bioinformatics</i> , 2010 , 11 Suppl 8, S8	3-6 5
12	Inter- α -Trypsin Inhibitor Heavy Chain 5 (ITIHS) Is a Natural Stabilizer of Hyaluronan That Modulates Biological Processes in the Skin. <i>Skin Pharmacology and Physiology</i> , 2020 , 33, 198-206	3 5
11	Epigenetic clocks are not accelerated in COVID-19 patients	4

10	SARS-CoV-2 RNA screening in routine pathology specimens. <i>Microbial Biotechnology</i> , 2021 , 14, 1627-1646.	6.3	4
9	Expression, Intracellular Localization, and Prognostic Value of Plasminogen Activator Inhibitor 1 and PAI-1 RNA-Binding Protein 1 in Primary and Recurrent Ovarian Cancer: A Study of the Tumor Bank Ovarian Cancer Network. <i>Gynecologic and Obstetric Investigation</i> , 2018 , 83, 508-514	2.5	4
8	The role of tumor-infiltrating lymphocytes in cholangiocarcinoma.. <i>Journal of Experimental and Clinical Cancer Research</i> , 2022 , 41, 127	12.8	4
7	Tissue pretreatment with formic acid might lower HercepTest scores in breast cancer. <i>Diagnostic Molecular Pathology</i> , 2006 , 15, 237-42		2
6	Towards sustainable data management in professional biobanking. <i>Studies in Health Technology and Informatics</i> , 2015 , 212, 94-102	0.5	2
5	Tumor- and stromal cell-specific expression of topoisomerase II α and HER-2/neu in primary and recurrent ovarian cancer: Results of a prospective study. <i>Molecular Medicine Reports</i> , 2009 , 2, 1011-6	2.9	1
4	Mutation analysis of multiple pilomatricomas in a patient with myotonic dystrophy type 1 suggests a DM1-associated hypermutation phenotype 2020 , 15, e0230003		
3	Mutation analysis of multiple pilomatricomas in a patient with myotonic dystrophy type 1 suggests a DM1-associated hypermutation phenotype 2020 , 15, e0230003		
2	Mutation analysis of multiple pilomatricomas in a patient with myotonic dystrophy type 1 suggests a DM1-associated hypermutation phenotype 2020 , 15, e0230003		
1	Mutation analysis of multiple pilomatricomas in a patient with myotonic dystrophy type 1 suggests a DM1-associated hypermutation phenotype 2020 , 15, e0230003		