
## Thomas J Dichristina

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6443658/publications.pdf Version: 2024-02-01



| #  | Article  | IF   | CITATIONS |
|----|--|------|-----------|
| 1  | lodate Reduction by Shewanella oneidensis Requires Genes Encoding an Extracellular<br>Dimethylsulfoxide Reductase. Frontiers in Microbiology, 2022, 13, 852942.  | 3.5  | 7         |
| 2  | Resistance of Perfluorooctanoic Acid to Degradation by the Microbially-Driven Fenton Reaction. FEMS Microbiology Letters, 2021, , .  | 1.8  | 0         |
| 3  | Fe(III) Oxide Reduction by Anaerobic Biofilm Formation-DeficientS-Ribosylhomocysteine Lyase (LuxS)<br>Mutant ofShewanella oneidensis. Geomicrobiology Journal, 2019, 36, 639-650.  | 2.0  | 0         |
| 4  | Metal Reduction and Protein Secretion Genes Required for Iodate Reduction by Shewanella oneidensis. Applied and Environmental Microbiology, 2019, 85, .  | 3.1  | 18        |
| 5  | lodate Reduction by <i>Shewanella oneidensis</i> Does Not Involve Nitrate Reductase.<br>Geomicrobiology Journal, 2018, 35, 570-579.  | 2.0  | 17        |
| 6  | Whole-genome sequencing reveals that Shewanella haliotis Kim et al. 2007 can be considered a later<br>heterotypic synonym of Shewanella algae Simidu et al. 1990. International Journal of Systematic and<br>Evolutionary Microbiology, 2018, 68, 1356-1360. | 1.7  | 20        |
| 7  | Microbial manganese(III) reduction fuelled by anaerobic acetate oxidation. Environmental<br>Microbiology, 2017, 19, 3475-3486.   | 3.8  | 17        |
| 8  | Degradation of the recalcitrant oil spill components anthracene and pyrene by a microbially driven<br>Fenton reaction. FEMS Microbiology Letters, 2017, 364, .   | 1.8  | 16        |
| 9  | Activation of an Otherwise Silent Xylose Metabolic Pathway in Shewanella oneidensis. Applied and<br>Environmental Microbiology, 2016, 82, 3996-4005.   | 3.1  | 16        |
| 10 | Detection of Metalâ€reducing Enzyme Complexes by Scanning Electrochemical Microscopy.<br>Electroanalysis, 2016, 28, 2459-2465.   | 2.9  | 7         |
| 11 | Simultaneous Transformation of Commingled Trichloroethylene, Tetrachloroethylene, and 1,4-Dioxane by a Microbially Driven Fenton Reaction in Batch Liquid Cultures. Applied and Environmental Microbiology, 2016, 82, 6335-6343.                             | 3.1  | 25        |
| 12 | Direct conversion of cellulose and hemicellulose to fermentable sugars by a microbially-driven Fenton reaction. Bioresource Technology, 2016, 218, 1133-1139.  | 9.6  | 20        |
| 13 | Identification of a molecular signature unique to metal-reducing <i>Gammaproteobacteria</i> . FEMS<br>Microbiology Letters, 2014, 350, 90-99.  | 1.8  | 22        |
| 14 | Electron transport and protein secretion pathways involved in <scp>Mn</scp> ( <scp>III</scp> )<br>reduction by <scp><i>S</i></scp> <i>hewanella oneidensis</i> . Environmental Microbiology Reports,<br>2014, 6, 490-500.                                    | 2.4  | 27        |
| 15 | Sulfur-mediated electron shuttling during bacterial iron reduction. Science, 2014, 344, 1039-1042.   | 12.6 | 175       |
| 16 | Microbially Driven Fenton Reaction for Degradation of the Widespread Environmental Contaminant<br>1,4-Dioxane. Environmental Science & Technology, 2014, 48, 12858-12867.  | 10.0 | 77        |
| 17 | Microbial Mn(IV) reduction requires an initial one-electron reductive solubilization step. Geochimica<br>Et Cosmochimica Acta, 2012, 99, 179-192.  | 3.9  | 57        |
| 18 | <i>Shewanella oneidensis</i> MRâ€1 mutants selected for their inability to produce soluble<br>organicâ€Fe(III) complexes are unable to respire Fe(III) as anaerobic electron acceptor. Environmental<br>Microbiology, 2010, 12, 938-950.                     | 3.8  | 38        |

THOMAS J DICHRISTINA

| #  | Article   | IF   | CITATIONS |
|----|---|------|-----------|
| 19 | Siderophores Are Not Involved in Fe(III) Solubilization during Anaerobic Fe(III) Respiration by<br><i>Shewanella oneidensis</i> MR-1. Applied and Environmental Microbiology, 2010, 76, 2425-2432.  | 3.1  | 39        |
| 20 | Outer Membrane-Associated Serine Protease Involved in Adhesion of <i>Shewanella oneidensis</i> to<br>Fe(III) Oxides. Environmental Science & Technology, 2010, 44, 68-73.   | 10.0 | 41        |
| 21 | Anaerobic Respiration of Elemental Sulfur and Thiosulfate by <i>Shewanella oneidensis</i> MR-1<br>Requires <i>psrA</i> , a Homolog of the <i>phsA</i> Gene of <i>Salmonella enterica</i> Serovar<br>Typhimurium LT2. Applied and Environmental Microbiology, 2009, 75, 5209-5217. | 3.1  | 117       |
| 22 | Dissolution Morphology of Iron (Oxy)(Hydr)Oxides Exposed to the Dissimilatory Iron-Reducing<br>Bacterium <i>Shewanella oneidensis</i> MR-1. Geomicrobiology Journal, 2009, 26, 83-92.   | 2.0  | 9         |
| 23 | A Conserved Histidine in Cytochrome c Maturation Permease CcmB of Shewanella putrefaciens Is<br>Required for Anaerobic Growth below a Threshold Standard Redox Potential. Journal of<br>Bacteriology, 2007, 189, 1036-1043.   | 2.2  | 12        |
| 24 | Shewanella putrefaciens produces an Fe(III)-solubilizing organic ligand during anaerobic respiration on insoluble Fe(III) oxides. Journal of Inorganic Biochemistry, 2007, 101, 1760-1767.  | 3.5  | 102       |
| 25 | A rapid mutant screening technique for detection of technetium [Tc(VII)] reduction-deficient mutants ofShewanella oneidensisMR-1. FEMS Microbiology Letters, 2006, 259, 282-287.  | 1.8  | 30        |
| 26 | Dissimilatory Fe(III) and Mn(IV) Reduction by Shewanella putrefaciens Requires ferE, a Homolog of the pulE (gspE) Type II Protein Secretion Gene. Journal of Bacteriology, 2002, 184, 142-151.  | 2.2  | 149       |
| 27 | Isolation of U(VI) reduction-deficient mutants ofShewanella putrefaciens. FEMS Microbiology Letters, 2000, 184, 143-148.  | 1.8  | 85        |
| 28 | Design and application of a rapid screening technique for isolation of selenite reduction-deficient mutants of Shewanella putrefaciens. Microbiological Research, 2000, 155, 79-85.   | 5.3  | 30        |
| 29 | Isolation of U(VI) reduction-deficient mutants of Shewanella putrefaciens. FEMS Microbiology<br>Letters, 2000, 184, 143-148.  | 1.8  | 4         |
| 30 | Microbially Driven Fenton Reaction for Transformation of Pentachlorophenol. Environmental<br>Science & Technology, 1999, 33, 1886-1891.   | 10.0 | 37        |
| 31 | Design and Application of Two Rapid Screening Techniques for Isolation of Mn(IV) Reduction-Deficient<br>Mutants of Shewanella putrefaciens. Applied and Environmental Microbiology, 1998, 64, 2716-2720.  | 3.1  | 28        |
| 32 | Bioextraction (Reductive Dissolution) of Iron from Low-Grade Iron Ores: Fundamental and Applied<br>Studiesa. Annals of the New York Academy of Sciences, 1994, 721, 440-449.  | 3.8  | 3         |
| 33 | Genetic Approaches in Bacteria with No Natural Genetic Systems. , 0, , 581-602.   |      | 1         |
| 34 | Breathing Iron: Molecular Mechanism of Microbial Iron Reduction by <i>Shewanella oneidensis</i> . ,<br>0, , 5.2.1-1-5.2.1-13.   |      | 4         |