## Kristin Taraldsen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6446950/publications.pdf

Version: 2024-02-01

44 papers

1,947 citations

394421 19 h-index 254184 43 g-index

47 all docs

47
docs citations

47 times ranked

2668 citing authors

#	Article	IF	CITATIONS
1	Adherence to mHealth and Paper-Based Versions of Lifestyle-Integrated Functional Exercise: A Secondary Analysis of Data From the PreventIT Feasibility Randomized Controlled Trial. Journal of Aging and Physical Activity, 2022, , 1-8.	1.0	O
2	The use of technology in creating individualized, meaningful activities for people living with dementia: A systematic review. Dementia, 2021, 20, 1442-1469.	2.0	34
3	Supporting identity and relationships amongst people with dementia through the use of technology: a qualitative interview study. International Journal of Qualitative Studies on Health and Well-being, 2021, 16, 1920349.	1.6	10
4	Sensitivity to Change and Responsiveness of the Original and the Shortened Version of the Community Balance & Mobility Scale for Young Seniors. Archives of Physical Medicine and Rehabilitation, 2021, 102, 2102-2108.	0.9	2
5	Change of physical activity parameters of hip and pelvic fracture patients during inpatient rehabilitation and after discharge: analysis of global and in-depth parameters. European Review of Aging and Physical Activity, 2021, 18, 9.	2.9	4
6	Towards personalized dementia care through meaningful activities supported by technology: A multisite qualitative study with care professionals. BMC Geriatrics, 2021, 21, 468.	2.7	4
7	Walking on common ground: a cross-disciplinary scoping review on the clinical utility of digital mobility outcomes. Npj Digital Medicine, 2021, 4, 149.	10.9	54
8	Gait, physical function, and physical activity in three groups of home-dwelling older adults with different severity of cognitive impairment $\hat{a}\in$ a cross-sectional study. BMC Geriatrics, 2021, 21, 670.	2.7	10
9	Creating and Validating a Shortened Version of the Community Balance and Mobility Scale for Application in People Who Are 61 to 70 Years of Age. Physical Therapy, 2020, 100, 180-191.	2.4	11
10	The association of basic and challenging motor capacity with mobility performance and falls in young seniors. Archives of Gerontology and Geriatrics, 2020, 90, 104134.	3.0	5
11	Digital Technology to Deliver a Lifestyle-Integrated Exercise Intervention in Young Seniors—The PreventIT Feasibility Randomized Controlled Trial. Frontiers in Digital Health, 2020, 2, 10.	2.8	12
12	Effects of an intervention to reduce fear of falling and increase physical activity during hip and pelvic fracture rehabilitation. Age and Ageing, 2020, 49, 771-778.	1.6	19
13	Predicting Advanced Balance Ability and Mobility with an Instrumented Timed Up and Go Test. Sensors, 2020, 20, 4987.	3.8	15
14	Client, caregiver, volunteer, and therapist views on a voluntary supported group exercise programme for older adults with dementia. BMC Geriatrics, 2020, 20, 235.	2.7	5
15	App-based Self-administrable Clinical Tests of Physical Function: Development and Usability Study. JMIR MHealth and UHealth, 2020, 8, e16507.	3.7	33
16	Attitudes Towards Adapted Lifestyle-Integrated Functional Exercise Developed for 60–70-Year-Olds: Perceptions of Participants and Trainers. Gerontology, 2019, 65, 599-609.	2.8	7
17	Delirium motor subtypes and prognosis in hospitalized geriatric patients – A prospective observational study. Journal of Psychosomatic Research, 2019, 122, 24-28.	2.6	13
18	The Adapted Lifestyle-Integrated Functional Exercise Program for Preventing Functional Decline in Young Seniors: Development and Initial Evaluation. Gerontology, 2019, 65, 362-374.	2.8	32

#	Article	IF	Citations
19	Protocol for the PreventIT feasibility randomised controlled trial of a lifestyle-integrated exercise intervention in young older adults. BMJ Open, 2019, 9, e023526.	1.9	34
20	My husband is not ill; he has memory loss - caregivers $\hat{A}$ perspectives on health care services for persons with dementia. BMC Geriatrics, 2019, 19, 75.	2.7	14
21	Motor activity across delirium motor subtypes in geriatric patients assessed using body-worn sensors: a Norwegian cross-sectional study. BMJ Open, 2019, 9, e026401.	1.9	10
22	Short and long-term clinical effectiveness and cost-effectiveness of a late-phase community-based balance and gait exercise program following hip fracture. The EVA-Hip Randomised Controlled Trial. PLoS ONE, 2019, 14, e0224971.	2.5	25
23	Performance-based clinical tests of balance and muscle strength used in young seniors: a systematic literature review. BMC Geriatrics, 2019, 19, 9.	2.7	47
24	The Use of Virtual and Immersive Technology in Creating Personalized Multisensory Spaces for People Living With Dementia (SENSE-GARDEN): Protocol for a Multisite Before-After Trial. JMIR Research Protocols, 2019, 8, e14096.	1.0	10
25	Environmental factors and risk of delirium in geriatric patients: an observational study. BMC Geriatrics, 2018, 18, 282.	2.7	14
26	Time spent lying, sitting, and upright during hospitalization after stroke: a prospective observation study. BMC Neurology, 2018, 18, 138.	1.8	25
27	Complexity of Daily Physical Activity Is More Sensitive Than Conventional Metrics to Assess Functional Change in Younger Older Adults. Sensors, 2018, 18, 2032.	3.8	18
28	Concurrent validity and reliability of the Community Balance and Mobility scale in young-older adults. BMC Geriatrics, 2018, 18, 156.	2.7	30
29	Daily Physical Activity in Total Hip Arthroplasty Patients Undergoing Different Surgical Approaches. American Journal of Physical Medicine and Rehabilitation, 2017, 96, 473-478.	1.4	19
30	Physical activity among hospitalized older adults – an observational study. BMC Geriatrics, 2017, 17, 110.	2.7	36
31	Mobile Health Applications to Promote Active and Healthy Ageing. Sensors, 2017, 17, 622.	3.8	151
32	Who benefits from orthogeriatric treatment? Results from the Trondheim hip-fracture trial. BMC Geriatrics, 2016, 16, 49.	2.7	38
33	The long-term effect of comprehensive geriatric care on gait after hip fracture: the Trondheim Hip Fracture Trial—a randomised controlled trial. Osteoporosis International, 2016, 27, 933-942.	3.1	55
34	Identification of gait domains and key gait variables following hip fracture. BMC Geriatrics, 2015, 15, 150.	2.7	45
35	The long-term effect of being treated in a geriatric ward compared to an orthopaedic ward on six measures of free-living physical behavior 4 and 12 months after a hip fracture - a randomised controlled trial. BMC Geriatrics, 2015, 15, 160.	2.7	28
36	Quantification of Outdoor Mobility by Use of Accelerometer-Measured Physical Behaviour. BioMed Research International, 2015, 2015, 1-7.	1.9	6

#	Article	IF	CITATION
37	Comprehensive geriatric care for patients with hip fractures: a prospective, randomised, controlled trial. Lancet, The, 2015, 385, 1623-1633.	13.7	449
38	Effectiveness of Task Specific Gait and Balance Exercise 4 Months After Hip Fracture: Protocol of a Randomized Controlled Trial — The Evaâ€Hip Study. Physiotherapy Research International, 2015, 20, 87-99.	1.5	9
39	Physical Behavior and Function Early After Hip Fracture Surgery in Patients Receiving Comprehensive Geriatric Care or Orthopedic Care—A Randomized Controlled Trial. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2014, 69A, 338-345.	3.6	84
40	Multiple Days of Monitoring Are Needed to Obtain a Reliable Estimate of Physical Activity in Hip-Fracture Patients. Journal of Aging and Physical Activity, 2014, 22, 173-177.	1.0	18
41	Physical activity monitoring by use of accelerometer-based body-worn sensors in older adults: A systematic literature review of current knowledge and applications. Maturitas, 2012, 71, 13-19.	2.4	164
42	Effect of in-hospital comprehensive geriatric assessment (CGA) in older people with hip fracture. The protocol of the Trondheim Hip Fracture Trial. BMC Geriatrics, 2011, 11, 18.	2.7	47
43	Evaluation of a Body-Worn Sensor System to Measure Physical Activity in Older People With Impaired Function. Physical Therapy, 2011, 91, 277-285.	2.4	225
44	Validation of the Falls Efficacy Scale-International in fall-prone older persons. Age and Ageing, 2010, 39, 259-259.	1.6	75