

# Minna Ilmakunnas

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6448315/publications.pdf>

Version: 2024-02-01

12  
papers

293  
citations

1163117

8  
h-index

1281871

11  
g-index

12  
all docs

12  
docs citations

12  
times ranked

510  
citing authors

#	ARTICLE	IF	CITATIONS
1	High mobility group box 1 protein as a marker of hepatocellular injury in human liver transplantation. Liver Transplantation, 2008, 14, 1517-1525.	2.4	76
2	Prothrombin complex concentrate in the reduction of blood loss during orthotopic liver transplantation: PROTON-trial. BMC Surgery, 2013, 13, 22.	1.3	65
3	Activation of neutrophils and monocytes by a leukocyte-depleting filter used throughout cardiopulmonary bypass. Journal of Thoracic and Cardiovascular Surgery, 2005, 129, 851-859.	0.8	44
4	Hepatic IL-8 release during graft procurement is associated with impaired graft function after human liver transplantation. Clinical Transplantation, 2010, 24, 29-35.	1.6	26
5	Factor XIII deficiency enhances thrombin generation due to impaired fibrin polymerization – An effect corrected by Factor XIII replacement. Thrombosis Research, 2017, 149, 56-61.	1.7	17
6	Activation of protein C during reperfusion in clinical liver transplantation. Transplantation, 2003, 75, 467-472.	1.0	14
7	Graft glycocalyx degradation in human liver transplantation. PLoS ONE, 2019, 14, e0221010.	2.5	14
8	Heparin Binding Protein in Adult Heart Surgery. Annals of Thoracic Surgery, 2019, 107, 1154-1159.	1.3	13
9	Graft protein C entrapment is associated with reduced phagocyte activation during reperfusion in human liver transplantation. Critical Care Medicine, 2006, 34, 426-432.	0.9	8
10	Relative Quantification of Several Plasma Proteins during Liver Transplantation Surgery. Journal of Biomedicine and Biotechnology, 2011, 2011, 1-12.	3.0	8
11	Endogenous protease inhibitor uptake within the graft during reperfusion in human liver transplantation. Journal of Hepato-Biliary-Pancreatic Sciences, 2010, 17, 158-165.	2.6	7
12	Neutrophil gelatinase-associated lipocalin does not originate from the kidney during reperfusion in clinical renal transplantation. Intensive Care Medicine Experimental, 2021, 9, 56.	1.9	1