Byung Sun Min

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/6463800/byung-sun-min-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

2,616 164 38 27 h-index g-index citations papers 168 5.06 3,098 4.1 avg, IF L-index ext. citations ext. papers

| # | Paper | IF | Citations |
|-----|---|-----|-----------|
| 164 | PTP1B and Eglucosidase inhibitory activities of the chemical constituents from Hedera rhombea fruits: Kinetic analysis and molecular docking simulation <i>Phytochemistry</i> , 2022 , 197, 113100 | 4 | 2 |
| 163 | Anti-osteoclastogenic Effects of Indole Alkaloids Isolated from Barley (Var.) Grass. <i>Journal of Agricultural and Food Chemistry</i> , 2021 , 69, 12994-13005 | 5.7 | 0 |
| 162 | PTP1B Inhibitory and Anti-inflammatory Properties of Constituents from Eclipta prostrata L. <i>Biological and Pharmaceutical Bulletin</i> , 2021 , 44, 298-304 | 2.3 | 4 |
| 161 | SARS-CoV-2 main protease inhibition by compounds isolated from Luffa cylindrica using molecular docking. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2021 , 40, 127972 | 2.9 | 4 |
| 160 | Diterpenoids isolated from the root of and their anti-inflammatory activity. <i>Natural Product Research</i> , 2021 , 35, 726-732 | 2.3 | 9 |
| 159 | Triterpenoids from Celastrus orbiculatus Thunb. inhibit RANKL-induced osteoclast formation and bone resorption via c-Fos signaling. <i>Journal of Natural Medicines</i> , 2021 , 75, 56-65 | 3.3 | 1 |
| 158 | Flavonoids from the peels of Citrus unshiu Markov. and their inhibitory effects on RANKL-induced osteoclastogenesis through the downregulation of c-Fos signaling in vitro. <i>Bioorganic Chemistry</i> , 2021 , 107, 104613 | 5.1 | 3 |
| 157 | Polyacetylenes and Flavonoids Isolated from Flowers of Carthamus tinctorius. <i>Chemistry of Natural Compounds</i> , 2021 , 57, 635-640 | 0.7 | O |
| 156 | Structural characterization of prenylated compounds from Broussonetia kazinoki and their antiosteoclastogenic activity. <i>Phytochemistry</i> , 2021 , 188, 112791 | 4 | O |
| 155 | PTP1B inhibition studies of biological active phloroglucinols from the rhizomes of Dryopteris crassirhizoma: Kinetic properties and molecular docking simulation. <i>International Journal of Biological Macromolecules</i> , 2021 , 188, 719-728 | 7.9 | 6 |
| 154 | Antioxidant and Antidiabetic Activities of Flavonoid Derivatives from the Outer Skins of L. <i>Journal of Agricultural and Food Chemistry</i> , 2020 , 68, 8797-8811 | 5.7 | 19 |
| 153 | C5, A Cassaine Diterpenoid Amine, Induces Apoptosis via the Extrinsic Pathways in Human Lung Cancer Cells and Human Lymphoma Cells. <i>International Journal of Molecular Sciences</i> , 2020 , 21, | 6.3 | 4 |
| 152 | Tetra-aryl cyclobutane and stilbenes from the rhizomes of Rheum undulatum and their Eglucosidase inhibitory activity: Biological evaluation, kinetic analysis, and molecular docking simulation. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2020 , 30, 127049 | 2.9 | 4 |
| 151 | Stilbenes with Potent Protein Tyrosine Phosphatase-1B Inhibitory Activity from the Roots of. <i>Journal of Natural Products</i> , 2020 , 83, 323-332 | 4.9 | 10 |
| 150 | Chemical constituents from the roots of Kadsura coccinea with their protein tyrosine phosphatase 1B and acetylcholinesterase inhibitory activities. <i>Archives of Pharmacal Research</i> , 2020 , 43, 204-213 | 6.1 | 6 |
| 149 | Identification of Anti-Inflammatory Constituents from Vietnamese Piper hymenophyllum. <i>Revista Brasileira De Farmacognosia</i> , 2020 , 30, 312-316 | 2 | |
| 148 | Inhibition of PTP1B by farnesylated 2-arylbenzofurans isolated from Morus alba root bark: unraveling the mechanism of inhibition based on in vitro and in silico studies. <i>Archives of Pharmacal Research</i> , 2020 , 43, 961-975 | 6.1 | 7 |

(2019-2020)

| 147 | Antioxidant and anti-browning property of 2-arylbenzofuran derivatives from Morus alba Linn root bark. <i>Food Chemistry</i> , 2020 , 309, 125739 | 8.5 | 12 |
|-----|---|-----|----|
| 146 | Glycerols and fatty acids isolated from Micractinium sp. KSF0031. <i>Biochemical Systematics and Ecology</i> , 2020 , 89, 104000 | 1.4 | O |
| 145 | Subchronic Toxicity Assessment of Phytolacca americana L. (Phytolaccaceae) in F344 Rats. <i>Natural Product Communications</i> , 2020 , 15, 1934578X2094165 | 0.9 | 1 |
| 144 | Sappanone A Prevents Left Ventricular Dysfunction in a Rat Myocardial Ischemia Reperfusion Injury Model. <i>International Journal of Molecular Sciences</i> , 2020 , 21, | 6.3 | 2 |
| 143 | Characterization of hydrocoptisonine metabolites in human liver microsomes using a high-resolution quadrupole-orbitrap mass spectrometer. <i>Xenobiotica</i> , 2020 , 50, 1423-1433 | 2 | 1 |
| 142 | Phytochemical and pharmacological properties of Myristica fragrans Houtt.: an updated review. <i>Archives of Pharmacal Research</i> , 2020 , 43, 1067-1092 | 6.1 | 13 |
| 141 | 3-Hydroxyolean-12-en-27-oic Acids Inhibit RANKL-Induced Osteoclastogenesis in Vitro and Inflammation-Induced Bone Loss in Vivo. <i>International Journal of Molecular Sciences</i> , 2020 , 21, | 6.3 | 3 |
| 140 | Albanol B from Mulberries Exerts Anti-Cancer Effect through Mitochondria ROS Production in Lung Cancer Cells and Suppresses In Vivo Tumor Growth. <i>International Journal of Molecular Sciences</i> , 2020 , 21, | 6.3 | 3 |
| 139 | 6,7,4RTrihydroxyflavone inhibits osteoclast formation and bone resorption in vitro and in vivo. <i>Phytotherapy Research</i> , 2019 , 33, 2948-2959 | 6.7 | 9 |
| 138 | Quantitation and Radical Scavenging Activity Evaluation of Iridoids and Phenylethanoids from the Roots of Phlomis umbrosa (Turcz.) using DPPH Free Radical and DPPH-HPLC Methods, and their Cytotoxicity. <i>Natural Product Sciences</i> , 2019 , 25, 122 | 1.1 | 6 |
| 137 | Water Extract of Pleurotus eryngii var. ferulae Prevents High-Fat Diet-Induced Obesity by Inhibiting Pancreatic Lipase. <i>Journal of Medicinal Food</i> , 2019 , 22, 178-185 | 2.8 | 3 |
| 136 | Identification of anti-osteoclastogenic compounds from Cleistocalyx operculatus flower buds and their effects on RANKL-induced osteoclastogenesis. <i>Journal of Functional Foods</i> , 2019 , 60, 103388 | 5.1 | 5 |
| 135 | Five new diterpenoids from the barks of Cinnamomum cassia (L.) J. Presl. <i>Phytochemistry Letters</i> , 2019 , 32, 23-28 | 1.9 | 8 |
| 134 | Protective effects of extract of Cleistocalyx operculatus flower buds and its isolated major constituent against LPS-induced endotoxic shock by activating the Nrf2/HO-1 pathway. <i>Food and Chemical Toxicology</i> , 2019 , 129, 125-137 | 4.7 | 7 |
| 133 | Arylbenzofurans from the Root Bark of as Triple Inhibitors of Cholinesterase, Esite Amyloid Precursor Protein Cleaving Enzyme 1, and Glycogen Synthase Kinase-3ERelevance to Alzheimer Disease. ACS Omega, 2019, 4, 6283-6294 | 3.9 | 15 |
| 132 | Insight into the PTP1B Inhibitory Activity of Arylbenzofurans: An In Vitro and In Silico Study. <i>Molecules</i> , 2019 , 24, | 4.8 | 10 |
| 131 | Anti-inflammatory and cytotoxic activities of constituents isolated from the fruits of Ziziphus jujuba var. inermis Rehder. <i>Floterap</i> [2019 , 137, 104261 | 3.2 | 7 |
| 130 | Cytotoxic Lactones from the Pericarps of Litsea japonica. <i>Natural Product Sciences</i> , 2019 , 25, 23 | 1.1 | 1 |

| 129 | Development of Analytical Method and Validation using HPLC/PDA for Discrimination between Artemisiae Argyi Folium and Artemisiae Iwayomogii Herba. <i>Natural Product Sciences</i> , 2019 , 25, 275 | 1.1 | | |
|-----|---|-----|----|--|
| 128 | Lignans from Saururus chinensis exhibit anti-inflammatory activity by influencing the Nrf2/HO-1 activation pathway. <i>Archives of Pharmacal Research</i> , 2019 , 42, 332-343 | 6.1 | 8 | |
| 127 | Investigation of chemical compounds from Chlamydomonas sp. KSF108 (Chlamydomonadaceae). <i>Biochemical Systematics and Ecology</i> , 2019 , 83, 4-6 | 1.4 | 1 | |
| 126 | Trichosanhemiketal A and B: Two 13,14-seco-13,14-epoxyporiferastanes from the root of Trichosanthes kirilowii Maxim. <i>Bioorganic Chemistry</i> , 2019 , 83, 105-110 | 5.1 | 7 | |
| 125 | Assessing the safety of an Ephedrae Herba aqueous extract in rats: A repeat dose toxicity study. <i>Regulatory Toxicology and Pharmacology</i> , 2018 , 94, 144-151 | 3.4 | 12 | |
| 124 | Cholinesterase inhibitory alkaloids from the rhizomes of Coptis chinensis. <i>Bioorganic Chemistry</i> , 2018 , 77, 625-632 | 5.1 | 7 | |
| 123 | Lactones from the pericarps of Litsea japonica and their anti-inflammatory activities. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2018 , 28, 2109-2115 | 2.9 | 8 | |
| 122 | A comparative study of Mentha arvensis L. and Mentha haplocalyx Briq. by HPLC. <i>Natural Product Research</i> , 2018 , 32, 239-242 | 2.3 | 7 | |
| 121 | Hepatoprotective effect of Cassia obtusifolia seed extract and constituents against oxidative damage induced by tert-butyl hydroperoxide in human hepatic HepG2 cells. <i>Journal of Food Biochemistry</i> , 2018 , 42, e12439 | 3.3 | 9 | |
| 120 | Anti-inflammatory activity of compounds from the rhizome of Cnidium officinale. <i>Archives of Pharmacal Research</i> , 2018 , 41, 977-985 | 6.1 | 9 | |
| 119 | Chalcone derivatives from the root bark of Morus alba L. act as inhibitors of PTP1B and Eglucosidase. <i>Phytochemistry</i> , 2018 , 155, 114-125 | 4 | 44 | |
| 118 | Two new naphthalenic lactone glycosides from Cassia obtusifolia L. seeds. <i>Archives of Pharmacal Research</i> , 2018 , 41, 737-742 | 6.1 | 6 | |
| 117 | Triterpenoids from Ziziphus jujuba induce apoptotic cell death in human cancer cells through mitochondrial reactive oxygen species production. <i>Food and Function</i> , 2018 , 9, 3895-3905 | 6.1 | 20 | |
| 116 | Anti-inflammatory activity of caffeic acid derivatives isolated from the roots of Salvia miltiorrhiza Bunge. <i>Archives of Pharmacal Research</i> , 2018 , 41, 64-70 | 6.1 | 35 | |
| 115 | Protein tyrosine phosphatase 1B inhibitors from natural sources. <i>Archives of Pharmacal Research</i> , 2018 , 41, 130-161 | 6.1 | 44 | |
| 114 | Desoxyrhapontigenin inhibits RANKL-induced osteoclast formation and prevents inflammation-mediated bone loss. <i>International Journal of Molecular Medicine</i> , 2018 , 42, 569-578 | 4.4 | 8 | |
| 113 | PTP1B inhibitory activity and molecular docking analysis of stilbene derivatives from the rhizomes of Rheum undulatum L. <i>Floterap</i> [2018 , 131, 119-126 | 3.2 | 14 | |
| 112 | Antioxidant Compounds Isolated from the Roots of Phlomis umbrosa Turcz <i>Natural Product Sciences</i> , 2018 , 24, 119 | 1.1 | 6 | |

| 111 | Moracin derivatives from Morus Radix as dual BACE1 and cholinesterase inhibitors with antioxidant and anti-glycation capacities. <i>Life Sciences</i> , 2018 , 210, 20-28 | 6.8 | 22 |
|-----|---|------------------|------------------|
| 110 | Alkaloids from Piper nigrum Exhibit Antiinflammatory Activity via Activating the Nrf2/HO-1 Pathway. <i>Phytotherapy Research</i> , 2017 , 31, 663-670 | 6.7 | 21 |
| 109 | Flavanonol glucosides from the aerial parts of Agrimonia pilosa Ledeb. and their acetylcholinesterase inhibitory effects. <i>Carbohydrate Research</i> , 2017 , 445, 75-79 | 2.9 | 13 |
| 108 | Structure-related protein tyrosine phosphatase 1B inhibition by naringenin derivatives. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2017 , 27, 2274-2280 | 2.9 | 22 |
| 107 | Cytotoxic and apoptosis-inducing activities against human lung cancer cell lines of cassaine diterpenoids from the bark of Erythrophleum fordii. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2017 , 27, 2946-2952 | 2.9 | 8 |
| 106 | PTP1B inhibitors from Selaginella tamariscina (Beauv.) Spring and their kinetic properties and molecular docking simulation. <i>Bioorganic Chemistry</i> , 2017 , 72, 273-281 | 5.1 | 16 |
| 105 | Ellagitannin and flavonoid constituents from Agrimonia pilosa Ledeb. with their protein tyrosine phosphatase and acetylcholinesterase inhibitory activities. <i>Bioorganic Chemistry</i> , 2017 , 72, 293-300 | 5.1 | 23 |
| 104 | Anti-inflammatory activities of compounds from twigs of Morus alba. Floterap 2017, 120, 17-24 | 3.2 | 17 |
| 103 | Chemical constituents from the fruits of Ligustrum japonicum and their inhibitory effects on T cell activation. <i>Phytochemistry</i> , 2017 , 141, 147-155 | 4 | 17 |
| 102 | Lignan derivatives from Selaginella tamariscina and their nitric oxide inhibitory effects in LPS-stimulated RAW 264.7 cells. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2017 , 27, 524-529 | 2.9 | 20 |
| 101 | Antifungal activity of sterols and dipsacus saponins isolated from Dipsacus asper roots against phytopathogenic fungi. <i>Pesticide Biochemistry and Physiology</i> , 2017 , 141, 103-108 | 4.9 | 21 |
| 100 | Sappanone A inhibits RANKL-induced osteoclastogenesis in BMMs and prevents inflammation-mediated bone loss. <i>International Immunopharmacology</i> , 2017 , 52, 230-237 | 5.8 | 21 |
| 99 | Anti-allergic and Cytotoxic Effects of Sesquiterpenoids and Phenylpropanoids Isolated from Magnolia biondii. <i>Natural Product Communications</i> , 2017 , 12, 1934578X1701201 | 0.9 | 1 |
| 98 | Cytotoxic Activity of Compounds from Styrax obassia. <i>Natural Product Communications</i> , 2017 , 12, 193 | 457&) (1 | 70 <u>/</u> 1200 |
| 97 | PTP1B inhibitory and cytotoxic activities of triterpenoids from the aerial parts of Agrimonia pilosa. <i>Medicinal Chemistry Research</i> , 2017 , 26, 2870-2878 | 2.2 | 11 |
| 96 | Triterpenoids and sterols from the grains of Echinochloa utilis Ohwi & Yabuno and their cytotoxic activity. <i>Biomedicine and Pharmacotherapy</i> , 2017 , 93, 202-207 | 7.5 | 4 |
| 95 | EMethyl artoflavanocoumarin from Juniperus chinensis exerts anti-diabetic effects by inhibiting PTP1B and activating the PI3K/Akt signaling pathway in insulin-resistant HepG2 cells. <i>Archives of Pharmacal Research</i> , 2017 , 40, 1403-1413 | 6.1 | 7 |
| 94 | Inhibitory effects of serratene-type triterpenoids from Lycopodium complanatum on cholinesterases and Becretase 1. <i>Chemico-Biological Interactions</i> , 2017 , 274, 150-157 | 5 | 11 |

| 93 | Prunin is a highly potent flavonoid from Prunus davidiana stems that inhibits protein tyrosine phosphatase 1B and stimulates glucose uptake in insulin-resistant HepG2 cells. <i>Archives of Pharmacal Research</i> , 2017 , 40, 37-48 | 6.1 | 26 |
|----|---|------------------|----|
| 92 | Inhibition of C1-Ten PTPase activity reduces insulin resistance through IRS-1 and AMPK pathways. <i>Scientific Reports</i> , 2017 , 7, 17777 | 4.9 | 6 |
| 91 | Four New Lignans and IL-2 Inhibitors from Magnoliae Flos. <i>Chemical and Pharmaceutical Bulletin</i> , 2017 , 65, 840-847 | 1.9 | 13 |
| 90 | Effect of Sipjeondaebo-Tang on the Pharmacokinetics of S-1, an Anticancer Agent, in Rats Evaluated by Population Pharmacokinetic Modeling. <i>Molecules</i> , 2017 , 22, | 4.8 | 1 |
| 89 | Sesquiterpenoids from the heartwood of Juniperu s chinensis. <i>Natural Product Sciences</i> , 2017 , 23, 208 | 1.1 | 6 |
| 88 | A cassaine diterpene alkaloid, 3日acetyl-nor-erythrophlamide, suppresses VEGF-induced angiogenesis and tumor growth via inhibiting eNOS activation. <i>Oncotarget</i> , 2017 , 8, 92346-92358 | 3.3 | 9 |
| 87 | Protein tyrosine phosphatase 1B (PTP1B) inhibitory activity and glucosidase inhibitory activity of compounds isolated from Agrimonia pilosa. <i>Pharmaceutical Biology</i> , 2016 , 54, 474-80 | 3.8 | 32 |
| 86 | Isolation of a New Homomonoterpene from Madhuca Pasquieri and Effect of Isolated Compounds on NO Production. <i>Natural Product Communications</i> , 2016 , 11, 1934578X1601100 | 0.9 | 1 |
| 85 | A subchronic toxicity study of Radix Dipsaci water extract by oral administration in F344 rats. <i>Regulatory Toxicology and Pharmacology</i> , 2016 , 81, 136-145 | 3.4 | 11 |
| 84 | Fucosterol activates the insulin signaling pathway in insulin resistant HepG2 cells via inhibiting PTP1B. <i>Archives of Pharmacal Research</i> , 2016 , 39, 1454-1464 | 6.1 | 11 |
| 83 | BACE1 molecular docking and anti-Alzheimerß disease activities of ginsenosides. <i>Journal of Ethnopharmacology</i> , 2016 , 190, 219-30 | 5 | 34 |
| 82 | Anti-Inflammatory Activity of a Novel Acetylene Isolated from the Roots of Angelica tenuissima Nakai. <i>Helvetica Chimica Acta</i> , 2016 , 99, 447-451 | 2 | 4 |
| 81 | Anti-inflammatory terpenylated coumarins from the leaves of Zanthoxylum schinifolium with Eglucosidase inhibitory activity. <i>Journal of Natural Medicines</i> , 2016 , 70, 276-81 | 3.3 | 18 |
| 80 | Inhibitory evaluation of oligonol on Eglucosidase, protein tyrosine phosphatase 1B, cholinesterase, and Esecretase 1 related to diabetes and Alzheimer disease. <i>Archives of Pharmacal Research</i> , 2016 , 39, 409-20 | 6.1 | 12 |
| 79 | Two-Week Repeated Dose Toxicity of Atractylodis Rhizoma Alba in F344 Rats. <i>Natural Product Sciences</i> , 2016 , 22, 180 | 1.1 | 1 |
| 78 | Antioxidant and Anti-Inflammatory Effects of Rhei Rhizoma and Coptidis Rhizoma Mixture on Reflux Esophagitis in Rats. <i>Evidence-based Complementary and Alternative Medicine</i> , 2016 , 2016, 205218 | 3g ^{.3} | 23 |
| 77 | PTP1B, Eglucosidase, and DPP-IV inhibitory effects for chromene derivatives from the leaves of Smilax china L. <i>Chemico-Biological Interactions</i> , 2016 , 253, 27-37 | 5 | 34 |
| 76 | Kinetics and molecular docking studies of loganin, morroniside and 7-O-galloyl-D-sedoheptulose derived from Corni fructus as cholinesterase and Execretase 1 inhibitors. <i>Archives of Pharmacal Research</i> , 2016 , 39, 794-805 | 6.1 | 23 |

(2015-2016)

| 75 | Potential pancreatic lipase inhibitory activity of phenolic constituents from the root bark of Morus alba L. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2016 , 26, 2788-2794 | 2.9 | 28 |
|----------------|---|-----|----|
| 74 | Anti-cholinesterases and memory improving effects of Vietnamese Xylia xylocarpa. <i>Chemistry Central Journal</i> , 2016 , 10, 48 | | 8 |
| 73 | Cytotoxic and anti-angiogenic effects of lanostane triterpenoids from Ganoderma lucidum. <i>Phytochemistry Letters</i> , 2015 , 12, 69-74 | 1.9 | 32 |
| 7 ² | Sappanone A exhibits anti-inflammatory effects via modulation of Nrf2 and NF- B . <i>International Immunopharmacology</i> , 2015 , 28, 328-36 | 5.8 | 38 |
| 71 | Caffeoylglycolic acid methyl ester, a major constituent of sorghum, exhibits anti-inflammatory activity via the Nrf2/heme oxygenase-1 pathway. <i>RSC Advances</i> , 2015 , 5, 17786-17796 | 3.7 | 18 |
| 70 | Anti-adipogenic effect of epiberberine is mediated by regulation of the Raf/MEK1/2/ERK1/2 and AMPK[Akt pathways. <i>Archives of Pharmacal Research</i> , 2015 , 38, 2153-62 | 6.1 | 15 |
| 69 | Selaginellin and biflavonoids as protein tyrosine phosphatase 1B inhibitors from Selaginella tamariscina and their glucose uptake stimulatory effects. <i>Bioorganic and Medicinal Chemistry</i> , 2015 , 23, 3730-7 | 3.4 | 45 |
| 68 | An improved HPLC-DAD method for quantitative comparisons of triterpenes in Ganoderma lucidum and its five related species originating from Vietnam. <i>Molecules</i> , 2015 , 20, 1059-77 | 4.8 | 10 |
| 67 | Isolation of cholinesterase and Esecretase 1 inhibiting compounds from Lycopodiella cernua. <i>Bioorganic and Medicinal Chemistry</i> , 2015 , 23, 3126-34 | 3.4 | 27 |
| 66 | Protein tyrosine phosphatase 1B (PTP1B) inhibitory constituents from the aerial parts of Tradescantia spathacea Sw. <i>Floterap</i> [12015, 103, 113-21 | 3.2 | 17 |
| 65 | Inhibitory effects of compounds from Styrax obassia on NO production. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2015 , 25, 5087-91 | 2.9 | 13 |
| 64 | Simultaneous quantitation and validation of method for the quality evaluation of Eucommiae cortex by HPLC/UV. <i>Archives of Pharmacal Research</i> , 2015 , 38, 2183-92 | 6.1 | 5 |
| 63 | A Novel Arginase Inhibitor Derived from Scutellavia indica Restored Endothelial Function in ApoE-Null Mice Fed a High-Cholesterol Diet. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2015 , 355, 57-65 | 4.7 | 14 |
| 62 | Kinetics and molecular docking studies of cholinesterase inhibitors derived from water layer of Lycopodiella cernua (L.) Pic. Serm. (II). <i>Chemico-Biological Interactions</i> , 2015 , 240, 74-82 | 5 | 13 |
| 61 | Inhibition of advanced glycation endproducts formation by Korean thistle, Cirsium maackii. <i>Asian Pacific Journal of Tropical Medicine</i> , 2015 , 8, 1-5 | 2.1 | 17 |
| 60 | Isolation of benzoic and cinnamic acid derivatives from the grains of Sorghum bicolor and their inhibition of lipopolysaccharide-induced nitric oxide production in RAW 264.7 cells. <i>Food Chemistry</i> , 2015 , 168, 512-9 | 8.5 | 43 |
| 59 | Chelidonine suppresses migration and invasion of MDA-MB-231 cells by inhibiting formation of the integrin-linked kinase/PINCH/Eparvin complex. <i>Molecular Medicine Reports</i> , 2015 , 12, 2161-8 | 2.9 | 13 |
| 58 | Chemical Constituents of Euonymus alatus (Thunb.) Sieb. and Their PTP1B and EGlucosidase Inhibitory Activities. <i>Phytotherapy Research</i> , 2015 , 29, 1540-8 | 6.7 | 19 |

| 57 | Quantitative and Pattern Recognition Analyses of Five Marker Compounds in Raphani Semen using High-Performance Liquid Chromatography. <i>Bulletin of the Korean Chemical Society</i> , 2015 , 36, 2307-2319 |) ^{1.2} | 1 |
|----|---|------------------|----|
| 56 | Anti-inflammatory Flavonoids Isolated from Passiflora foetida. <i>Natural Product Communications</i> , 2015 , 10, 1934578X1501000 | 0.9 | 6 |
| 55 | Anti-inflammatory Compounds from Ampelopsis cantoniensis. <i>Natural Product Communications</i> , 2015 , 10, 1934578X1501000 | 0.9 | 1 |
| 54 | Compounds from the aerial parts of Piper bavinum and their anti-cholinesterase activity. <i>Archives of Pharmacal Research</i> , 2015 , 38, 677-82 | 6.1 | 9 |
| 53 | Anti-inflammatory activity of phenolic compounds from the whole plant of Scutellaria indica. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2015 , 25, 1129-34 | 2.9 | 23 |
| 52 | Anti-inflammatory and heme oxygenase-1 inducing activities of lanostane triterpenes isolated from mushroom Ganoderma lucidum in RAW264.7 cells. <i>Toxicology and Applied Pharmacology</i> , 2014 , 280, 434 | 1-42 | 30 |
| 51 | In vitro apoptotic effect of cassaine-type diterpene amides from Erythrophleum fordii on PC-3 prostate cancer cells. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2014 , 24, 4989-94 | 2.9 | 9 |
| 50 | Coptis chinensis alkaloids exert anti-adipogenic activity on 3T3-L1 adipocytes by downregulating C/EBP-land PPAR-[] <i>Flioterap</i> [] 2014 , 98, 199-208 | 3.2 | 68 |
| 49 | The anti-inflammatory effect of 3-deoxysappanchalcone is mediated by inducing heme oxygenase-1 via activating the AKT/mTOR pathway in murine macrophages. <i>International Immunopharmacology</i> , 2014 , 22, 420-6 | 5.8 | 21 |
| 48 | Cholinesterase inhibitors from the roots of Harpagophytum procumbens. <i>Archives of Pharmacal Research</i> , 2014 , 37, 1124-9 | 6.1 | 16 |
| 47 | Cassaine diterpene alkaloids from Erythrophleum fordii and their anti-angiogenic effect. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2014 , 24, 168-72 | 2.9 | 14 |
| 46 | Endothelial nitric oxide synthase activation through obacunone-dependent arginase inhibition restored impaired endothelial function in ApoE-null mice. <i>Vascular Pharmacology</i> , 2014 , 60, 102-9 | 5.9 | 14 |
| 45 | Pharmacokinetic alteration of baclofen by multiple oral administration of herbal medicines in rats. Evidence-based Complementary and Alternative Medicine, 2014, 2014, 402126 | 2.3 | 6 |
| 44 | Trogopterins A-C: Three new neolignans from feces of Trogopterus xanthipes. <i>Beilstein Journal of Organic Chemistry</i> , 2014 , 10, 2955-2962 | 2.5 | 9 |
| 43 | Potent Acetylcholinesterase Inhibitory Compounds from Myristica fragrans. <i>Natural Product Communications</i> , 2014 , 9, 1934578X1400900 | 0.9 | 8 |
| 42 | Arginase II inhibitory activity of flavonoid compounds from Scutellaria indica. <i>Archives of Pharmacal Research</i> , 2013 , 36, 922-6 | 6.1 | 27 |
| 41 | Inhibitory effect on NO production of triterpenes from the fruiting bodies of Ganoderma lucidum. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2013 , 23, 1428-32 | 2.9 | 40 |
| 40 | Kinetics and molecular docking studies of pimarane-type diterpenes as protein tyrosine phosphatase (PTP1B) inhibitors from Aralia continentalis roots. <i>Archives of Pharmacal Research</i> , 2013 36 957-65 | 6.1 | 17 |

(2009-2013)

| 39 | Apoptosis-inducing and antitumor activity of neolignans isolated from Magnolia officinalis in HeLa cancer cells. <i>Phytotherapy Research</i> , 2013 , 27, 1419-22 | 6.7 | 10 | |
|----|---|-----|----|--|
| 38 | Anti-platelet activity of erythro-(7S,8R)-7-acetoxy-3,4,3R5Rtetramethoxy-8-O-4Rneolignan from Myristica fragrans. <i>Phytotherapy Research</i> , 2013 , 27, 1694-9 | 6.7 | 12 | |
| 37 | Compounds from the heartwood of Caesalpinia sappan and their anti-inflammatory activity. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2012 , 22, 7436-9 | 2.9 | 44 | |
| 36 | Anticomplementary activity of oleanane-type triterpenes from the roots of Aceriphyllum rossii. <i>Archives of Pharmacal Research</i> , 2012 , 35, 1003-8 | 6.1 | 5 | |
| 35 | Arginase II Inhibitory Activity of Phenolic Compounds from Saururus chinensis. <i>Bulletin of the Korean Chemical Society</i> , 2012 , 33, 3079-3082 | 1.2 | 12 | |
| 34 | Inhibitory effect on NO production of phenolic compounds from Myristica fragrans. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2011 , 21, 6884-7 | 2.9 | 24 | |
| 33 | Simultaneous determination of bioactive flavonoids in some selected Korean thistles by high-performance liquid chromatography. <i>Archives of Pharmacal Research</i> , 2011 , 34, 455-61 | 6.1 | 27 | |
| 32 | Ergosta-7,22-diene-2即卧riol from the fruit bodies of Ganoderma lucidum induces apoptosis in human myelocytic HL-60 cells. <i>Phytotherapy Research</i> , 2011 , 25, 1579-85 | 6.7 | 11 | |
| 31 | Inhibitory Effect of Lignans from Myristica fragrans on LPS-induced NO Production in RAW264.7 Cells. <i>Bulletin of the Korean Chemical Society</i> , 2011 , 32, 4059-4062 | 1.2 | 8 | |
| 30 | 3-Oxoolean-12-en-27-oic acid inhibits the proliferation of non-small cell lung carcinoma cells by inducing cell-cycle arrest at G0/G1 phase. <i>Anticancer Research</i> , 2011 , 31, 2179-85 | 2.3 | 3 | |
| 29 | Protective action of 9-hydroxypinoresinol against oxidative damage in brain of mice challenged with kainic acid. <i>Journal of Pharmacy and Pharmacology</i> , 2010 , 59, 1585-1585 | 4.8 | | |
| 28 | Addition: Anticholinesterase and Antioxidant Constituents from Gloiopeltis furcata [Chem. Pharm. Bull. 58(9): 1236-1239 (2010)]. <i>Chemical and Pharmaceutical Bulletin</i> , 2010 , 58, 1554 | 1.9 | | |
| 27 | Cholinesterase inhibitors from Cleistocalyx operculatus buds. <i>Archives of Pharmacal Research</i> , 2010 , 33, 1665-70 | 6.1 | 33 | |
| 26 | Dihydrobenzofuran Norlignans from the Leaves of Cedrela sinensis A. Juss. <i>Helvetica Chimica Acta</i> , 2010 , 93, 272-276 | 2 | 19 | |
| 25 | Study on the Constituents of Roots of Aceriphyllum rossii. <i>Helvetica Chimica Acta</i> , 2010 , 93, 1803-1807 | 2 | 3 | |
| 24 | Homoisoflavonoid derivatives from the roots of Ophiopogon japonicus and their in vitro anti-inflammation activity. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2010 , 20, 2412-6 | 2.9 | 47 | |
| 23 | Antioxidant activities of coumarins from Korean medicinal plants and their structure-activity relationships. <i>Phytotherapy Research</i> , 2010 , 24, 101-6 | 6.7 | 88 | |
| 22 | Two New Diterpenes from the Twigs of Cinnamomum cassia. <i>Helvetica Chimica Acta</i> , 2009 , 92, 2058-206 | 62 | 17 | |

| 21 | Steroids and triterpenes from the fruit bodies of Ganoderma lucidum and their anti-complement activity. <i>Archives of Pharmacal Research</i> , 2009 , 32, 1573-9 | 6.1 | 55 |
|----|---|-----|-----|
| 20 | Phenolic glycosides from Alangium salviifolium leaves with inhibitory activity on LPS-induced NO, PGE(2), and TNF-alpha production. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009 , 19, 4389-93 | 2.9 | 25 |
| 19 | Tyrosinase-inhibitory constituents from the twigs of Cinnamomum cassia. <i>Journal of Natural Products</i> , 2009 , 72, 1205-8 | 4.9 | 37 |
| 18 | Protein tyrosine phosphatase 1B inhibitory by dammaranes from Vietnamese Giao-Co-Lam tea. <i>Journal of Ethnopharmacology</i> , 2009 , 124, 240-5 | 5 | 28 |
| 17 | Anticomplementary activity of triterpenoids from the whole plant of Aceriphyllum rossii against the classical pathway. <i>Planta Medica</i> , 2008 , 74, 726-9 | 3.1 | 18 |
| 16 | Antioxidative flavonoids from Cleistocalyx operculatus buds. <i>Chemical and Pharmaceutical Bulletin</i> , 2008 , 56, 1725-8 | 1.9 | 18 |
| 15 | Lipoxygenase inhibitory constituents from rhubarb. <i>Archives of Pharmacal Research</i> , 2008 , 31, 598-605 | 6.1 | 60 |
| 14 | Triterpenoids and a sterol from the stem-bark of Styrax japonica and their protein tyrosine phosphatase 1B inhibitory activities. <i>Phytotherapy Research</i> , 2008 , 22, 1303-6 | 6.7 | 27 |
| 13 | Isolation of coumarins and ferulate from the roots of Angelica purpuraefolia and the antitumor activity of khellactone. <i>Phytotherapy Research</i> , 2007 , 21, 406-9 | 6.7 | 12 |
| 12 | A phenylpropanoid glycoside with antioxidant activity from Picria tel-ferae. <i>Archives of Pharmacal Research</i> , 2007 , 30, 1062-6 | 6.1 | 13 |
| 11 | Anticomplement activity of cycloartane glycosides from the rhizome of Cimicifuga foetida. <i>Phytotherapy Research</i> , 2006 , 20, 945-8 | 6.7 | 30 |
| 10 | Two novel furan derivatives from Phellinus linteus with anti-complement activity. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2006 , 16, 3255-7 | 2.9 | 20 |
| 9 | A new furofuran lignan with antioxidant and antiseizure activities from the leaves of Petasites japonicus. <i>Archives of Pharmacal Research</i> , 2005 , 28, 1023-6 | 6.1 | 19 |
| 8 | Inhibitory effects of Korean plants on HIV-1 activities. <i>Phytotherapy Research</i> , 2001 , 15, 481-6 | 6.7 | 36 |
| 7 | Cytotoxic triterpenes from Crataegus pinnatifida. Archives of Pharmacal Research, 2000 , 23, 155-8 | 6.1 | 39 |
| 6 | Antifungal activity of magnolol and honokiol. <i>Archives of Pharmacal Research</i> , 2000 , 23, 46-9 | 6.1 | 102 |
| 5 | Inhibitory constituents against HIV-1 protease from Agastache rugosa. <i>Archives of Pharmacal Research</i> , 1999 , 22, 75-7 | 6.1 | 22 |
| 4 | Screening of Korean plants against human immunodeficiency virus type 1 protease. <i>Phytotherapy Research</i> , 1999 , 13, 680-2 | 6.7 | 22 |

LIST OF PUBLICATIONS

| 3 | A cytotoxic constituent fromSophora flavescens. <i>Archives of Pharmacal Research</i> , 1997 , 20, 342-5 | 6.1 | 13 |
|---|---|-----|----|
| 2 | Antitumor activity of 2(S)-5,2R5Rtrihydroxy-7,8-dimethoxyflavanone and its analogues. <i>Archives of Pharmacal Research</i> , 1997 , 20, 368-71 | 6.1 | 5 |
| 1 | Synthesis and structure-activity relationship of cytotoxic 5,2?,5?-trihydroxy-7,8-dimethoxyflavanone analogues. <i>Archives of Pharmacal Research</i> , 1996 , 19, 543-550 | 6.1 | 6 |