

Lin Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/647765/publications.pdf>

Version: 2024-02-01

15
papers

746
citations

687363

13
h-index

996975

15
g-index

15
all docs

15
docs citations

15
times ranked

1046
citing authors

#	ARTICLE	IF	CITATIONS
1	Receptor-Like Cytoplasmic Kinases Directly Link Diverse Pattern Recognition Receptors to the Activation of Mitogen-Activated Protein Kinase Cascades in Arabidopsis. <i>Plant Cell</i> , 2018, 30, 1543-1561.	6.6	219
2	Two novel NAC transcription factors regulate gene expression and flowering time by associating with the histone demethylase JM14. <i>Nucleic Acids Research</i> , 2015, 43, 1469-1484.	14.5	94
3	Plant Mitogen-Activated Protein Kinase Cascades in Environmental Stresses. <i>International Journal of Molecular Sciences</i> , 2021, 22, 1543.	4.1	61
4	A plant-specific SWR1 chromatin remodeling complex couples histone H2A.Z deposition with nucleosome sliding. <i>EMBO Journal</i> , 2020, 39, e102008.	7.8	57
5	IDN2 and Its Paralogs Form a Complex Required for RNA-Directed DNA Methylation. <i>PLoS Genetics</i> , 2012, 8, e1002693.	3.5	52
6	A methylated-DNA-binding complex required for plant development mediates transcriptional activation of promoter methylated genes. <i>Journal of Integrative Plant Biology</i> , 2019, 61, 120-139.	8.5	45
7	The PEAT protein complexes are required for histone deacetylation and heterochromatin silencing. <i>EMBO Journal</i> , 2018, 37, .	7.8	42
8	Dual Recognition of H3K4me3 and DNA by the ISWI Component ARID5 Regulates the Floral Transition in Arabidopsis. <i>Plant Cell</i> , 2020, 32, 2178-2195.	6.6	34
9	The CBP/p300 histone acetyltransferases function as plant-specific MEDIATOR subunits in <i>Arabidopsis</i> . <i>Journal of Integrative Plant Biology</i> , 2021, 63, 755-771.	8.5	29
10	AtINO80 represses photomorphogenesis by modulating nucleosome density and H2A.Z incorporation in light-related genes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 33679-33688.	7.1	22
11	Three functionally redundant plant-specific paralogs are core subunits of the SAGA histone acetyltransferase complex in Arabidopsis. <i>Molecular Plant</i> , 2021, 14, 1071-1087.	8.3	20
12	A histone H3K27me3 reader cooperates with a family of PHD finger-containing proteins to regulate flowering time in <i>Arabidopsis</i> . <i>Journal of Integrative Plant Biology</i> , 2021, 63, 787-802.	8.5	19
13	Arabidopsis RPD3-like histone deacetylases form multiple complexes involved in stress response. <i>Journal of Genetics and Genomics</i> , 2021, 48, 369-383.	3.9	18
14	COMPASS functions as a module of the INO80 chromatin remodeling complex to mediate histone H3K4 methylation in Arabidopsis. <i>Plant Cell</i> , 2021, 33, 3250-3271.	6.6	17
15	The <i>Arabidopsis</i> NuA4 histone acetyltransferase complex is required for chlorophyll biosynthesis and photosynthesis. <i>Journal of Integrative Plant Biology</i> , 2022, 64, 901-914.	8.5	17