

Lin Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/647765/publications.pdf>

Version: 2024-02-01

15
papers

746
citations

687363

13
h-index

996975

15
g-index

15
all docs

15
docs citations

15
times ranked

1046
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | The <i>Arabidopsis</i> NuA4 histone acetyltransferase complex is required for chlorophyll biosynthesis and photosynthesis. <i>Journal of Integrative Plant Biology</i> , 2022, 64, 901-914. | 8.5 | 17 |
| 2 | The CBP/p300 histone acetyltransferases function as plant-specific MEDIATOR subunits in <i>Arabidopsis</i> . <i>Journal of Integrative Plant Biology</i> , 2021, 63, 755-771. | 8.5 | 29 |
| 3 | Plant Mitogen-Activated Protein Kinase Cascades in Environmental Stresses. <i>International Journal of Molecular Sciences</i> , 2021, 22, 1543. | 4.1 | 61 |
| 4 | A histone H3K27me3 reader cooperates with a family of PHD finger-containing proteins to regulate flowering time in <i>Arabidopsis</i> . <i>Journal of Integrative Plant Biology</i> , 2021, 63, 787-802. | 8.5 | 19 |
| 5 | <i>Arabidopsis</i> RPD3-like histone deacetylases form multiple complexes involved in stress response. <i>Journal of Genetics and Genomics</i> , 2021, 48, 369-383. | 3.9 | 18 |
| 6 | Three functionally redundant plant-specific paralogs are core subunits of the SAGA histone acetyltransferase complex in <i>Arabidopsis</i> . <i>Molecular Plant</i> , 2021, 14, 1071-1087. | 8.3 | 20 |
| 7 | COMPASS functions as a module of the INO80 chromatin remodeling complex to mediate histone H3K4 methylation in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2021, 33, 3250-3271. | 6.6 | 17 |
| 8 | Dual Recognition of H3K4me3 and DNA by the ISWI Component ARID5 Regulates the Floral Transition in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2020, 32, 2178-2195. | 6.6 | 34 |
| 9 | A plant-specific SWR1 chromatin remodeling complex couples histone H2A.Z deposition with nucleosome sliding. <i>EMBO Journal</i> , 2020, 39, e102008. | 7.8 | 57 |
| 10 | AtINO80 represses photomorphogenesis by modulating nucleosome density and H2A.Z incorporation in light-related genes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 33679-33688. | 7.1 | 22 |
| 11 | A methylated-DNA-binding complex required for plant development mediates transcriptional activation of promoter methylated genes. <i>Journal of Integrative Plant Biology</i> , 2019, 61, 120-139. | 8.5 | 45 |
| 12 | The PEAT protein complexes are required for histone deacetylation and heterochromatin silencing. <i>EMBO Journal</i> , 2018, 37, . | 7.8 | 42 |
| 13 | Receptor-Like Cytoplasmic Kinases Directly Link Diverse Pattern Recognition Receptors to the Activation of Mitogen-Activated Protein Kinase Cascades in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2018, 30, 1543-1561. | 6.6 | 219 |
| 14 | Two novel NAC transcription factors regulate gene expression and flowering time by associating with the histone demethylase JMJ14. <i>Nucleic Acids Research</i> , 2015, 43, 1469-1484. | 14.5 | 94 |
| 15 | IDN2 and Its Paralogs Form a Complex Required for RNA-Directed DNA Methylation. <i>PLoS Genetics</i> , 2012, 8, e1002693. | 3.5 | 52 |