

Mary H Grace

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6480339/mary-h-grace-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

92
papers

3,071
citations

33
h-index

53
g-index

99
ext. papers

3,522
ext. citations

4.4
avg, IF

5.27
L-index

#	Paper	IF	Citations
92	Spray-dried and freeze-dried protein-spinach particles; effect of drying technique and protein type on the bioaccessibility of carotenoids, chlorophylls, and phenolics.. <i>Food Chemistry</i> , 2022 , 388, 133017	8.5	0
91	Alaskan Bog Blueberry (<i>Vaccinium uliginosum</i>) Extract as an Innovative Topical Approach to Prevent UV-Induced Skin Damage. <i>Cosmetics</i> , 2021 , 8, 112	2.7	1
90	Novel Spray Dried Algae-Rosemary Particles Attenuate Pollution-Induced Skin Damage. <i>Molecules</i> , 2021 , 26,	4.8	3
89	The same anthocyanins served four different ways: Insights into anthocyanin structure-function relationships from the wintergreen orchid, <i>Tipularia discolor</i> . <i>Plant Science</i> , 2021 , 303, 110793	5.3	2
88	Whey and soy proteins as wall materials for spray drying rosemary: Effects on polyphenol composition, antioxidant activity, bioaccessibility after in vitro gastrointestinal digestion and stability during storage. <i>LWT - Food Science and Technology</i> , 2021 , 149, 111901	5.4	3
87	Neuroprotective mechanisms of red clover and soy isoflavones in Parkinson's disease models. <i>Food and Function</i> , 2021 , 12, 11987-12007	6.1	0
86	Phytoecdysteroids Do Not Have Anabolic Effects in Skeletal Muscle in Sedentary Aging Mice. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	2
85	Hypoglycaemic, insulin releasing, and hepatoprotective effect of the aqueous extract of Aloe perryi Baker resin (Socotran Aloe) in streptozotocin-induced diabetic rats. <i>Journal of Taibah University for Science</i> , 2020 , 14, 1671-1685	3	2
84	Phenolic content, anti-inflammatory properties, and dermal wound repair properties of industrially processed and non-processed acai from the Brazilian Amazon. <i>Food and Function</i> , 2020 , 11, 4903-4914	6.1	10
83	Enhanced stability of berry pomace polyphenols delivered in protein-polyphenol aggregate particles to an in vitro gastrointestinal digestion model. <i>Food Chemistry</i> , 2020 , 331, 127279	8.5	23
82	LC-MS characterization of bioactive metabolites from two Yemeni Aloe spp. with antioxidant and antidiabetic properties. <i>Arabian Journal of Chemistry</i> , 2020 , 13, 5040-5049	5.9	15
81	Diversity in Metabolites and Fruit Quality Traits in Blueberry Enables Ploidy and Species Differentiation and Establishes a Strategy for Future Genetic Studies. <i>Frontiers in Plant Science</i> , 2020 , 11, 370	6.2	9
80	In Vitro Evaluation of a Novel Synthetic Bilirubin Analog as an Antioxidant and Cytoprotective Agent for Pancreatic Islet Transplantation. <i>Cell Transplantation</i> , 2020 , 29, 963689720906417	4	3
79	Blueberry Extracts as a Novel Approach to Prevent Ozone-Induced Cutaneous Inflammation Activation. <i>Oxidative Medicine and Cellular Longevity</i> , 2020 , 2020, 9571490	6.7	5
78	Changes in the bioactive properties of strawberries caused by the storage in oxygen- and carbon dioxide-enriched atmospheres. <i>Food Science and Nutrition</i> , 2019 , 7, 2527-2536	3.2	3
77	Alaskan Berry Extracts Promote Dermal Wound Repair Through Modulation of Bioenergetics and Integrin Signaling. <i>Frontiers in Pharmacology</i> , 2019 , 10, 1058	5.6	17
76	Photosynthetic Profiles of Green, Purple, and Spotted-Leaf Morphotypes of <i>Tipularia discolor</i> (Orchidaceae). <i>Southeastern Naturalist</i> , 2019 , 18, 641	0.4	1

75	Changes due to high oxygen and high carbon dioxide atmospheres on the general quality and the polyphenolic profile of strawberries. <i>Postharvest Biology and Technology</i> , 2019 , 148, 49-57	6.2	14
74	Simultaneous LC-MS quantification of anthocyanins and non-anthocyanin phenolics from blueberries with widely divergent profiles and biological activities. <i>Food Chemistry</i> , 2019 , 277, 336-346	8.5	45
73	Anti-inflammatory and wound healing properties of polyphenolic extracts from strawberry and blackberry fruits. <i>Food Research International</i> , 2019 , 121, 453-462	7	47
72	In vitro mineral nutrition of <i>Curcuma longa</i> L. affects production of volatile compounds in rhizomes after transfer to the greenhouse. <i>BMC Plant Biology</i> , 2018 , 18, 122	5.3	2
71	Increased Plasma Levels of Gut-Derived Phenolics Linked to Walking and Running Following Two Weeks of Flavonoid Supplementation. <i>Nutrients</i> , 2018 , 10,	6.7	24
70	One-step isolation of carnosic acid and carnosol from rosemary by centrifugal partition chromatography. <i>Journal of Separation Science</i> , 2017 , 40, 1057-1062	3.4	10
69	Inter- and intra-seasonal changes in anthocyanin accumulation and global metabolite profiling of six blueberry genotypes. <i>Journal of Food Composition and Analysis</i> , 2017 , 59, 105-110	4.1	11
68	Wild blueberry polyphenol-protein food ingredients produced by three drying methods: Comparative physico-chemical properties, phytochemical content, and stability during storage. <i>Food Chemistry</i> , 2017 , 235, 76-85	8.5	52
67	αAmylase and βGlucosidase Inhibitory Activities of Phenolic Extracts from <i>Eucalyptus grandis</i> L. urophylla Bark. <i>Journal of Chemistry</i> , 2017 , 2017, 1-7	2.3	9
66	Pendulaosides A and B. Two acylated triterpenoid saponins from <i>Harpullia pendula</i> seed extract. <i>Phytochemistry Letters</i> , 2017 , 21, 278-282	1.9	2
65	Influence of Ingesting a Flavonoid-Rich Supplement on the Metabolome and Concentration of Urine Phenolics in Overweight/Obese Women. <i>Journal of Proteome Research</i> , 2017 , 16, 2924-2935	5.6	18
64	Polyphenols isolated from <i>Acacia mearnsii</i> bark with anti-inflammatory and carbolytic enzyme inhibitory activities. <i>Chinese Journal of Natural Medicines</i> , 2017 , 15, 816-824	2.8	14
63	Metabolic Effects of Berries with Structurally Diverse Anthocyanins. <i>International Journal of Molecular Sciences</i> , 2017 , 18,	6.3	76
62	Quantitative comparison of phytochemical profile, antioxidant, and anti-inflammatory properties of blackberry fruits adapted to Argentina. <i>Journal of Food Composition and Analysis</i> , 2016 , 47, 82-91	4.1	39
61	Unraveling Anthocyanin Bioavailability for Human Health. <i>Annual Review of Food Science and Technology</i> , 2016 , 7, 375-93	14.7	130
60	Impact of a new postharvest disinfection method based on peracetic acid fogging on the phenolic profile of strawberries. <i>Postharvest Biology and Technology</i> , 2016 , 117, 197-205	6.2	15
59	A polyphenol-rich fraction obtained from table grapes decreases adiposity, insulin resistance and markers of inflammation and impacts gut microbiota in high-fat-fed mice. <i>Journal of Nutritional Biochemistry</i> , 2016 , 31, 150-65	6.3	72
58	Phytochemical Characterization and Anti-inflammatory Properties of <i>Acacia mearnsii</i> Leaves. <i>Natural Product Communications</i> , 2016 , 11, 1934578X1601100	0.9	8

57	Chemical composition, antioxidant and anti-inflammatory properties of pistachio hull extracts. <i>Food Chemistry</i> , 2016 , 210, 85-95	8.5	58
56	In vitro lipolytic, antioxidant and anti-inflammatory activities of roasted pistachio kernel and skin constituents. <i>Food and Function</i> , 2016 , 7, 4285-4298	6.1	22
55	Novel strategies for capturing health-protective mango phytochemicals in shelf stable food matrices. <i>International Journal of Food Sciences and Nutrition</i> , 2015 , 66, 175-85	3.7	5
54	Black Currant Anthocyanins Attenuate Weight Gain and Improve Glucose Metabolism in Diet-Induced Obese Mice with Intact, but Not Disrupted, Gut Microbiome. <i>Journal of Agricultural and Food Chemistry</i> , 2015 , 63, 6172-80	5.7	101
53	Alaskan seaweeds lower inflammation in RAW 264.7 macrophages and decrease lipid accumulation in 3T3-L1 adipocytes. <i>Journal of Functional Foods</i> , 2015 , 15, 396-407	5.1	31
52	Characterization of Phenolic Compounds and Antioxidant and Anti-inflammatory Activities from Mamuyo (<i>Styrax ramirezii</i> Greenm.) Fruit. <i>Journal of Agricultural and Food Chemistry</i> , 2015 , 63, 10459-65	5.7	24
51	Novel value-added uses for sweet potato juice and flour in polyphenol- and protein-enriched functional food ingredients. <i>Food Science and Nutrition</i> , 2015 , 3, 415-24	3.2	13
50	Neuroprotective effects of anthocyanin- and proanthocyanidin-rich extracts in cellular models of Parkinson's disease. <i>Brain Research</i> , 2014 , 1555, 60-77	3.7	125
49	Effects of a high fat meal matrix and protein complexation on the bioaccessibility of blueberry anthocyanins using the TNO gastrointestinal model (TIM-1). <i>Food Chemistry</i> , 2014 , 142, 349-57	8.5	115
48	Inhibitory effects of wild blueberry anthocyanins and other flavonoids on biomarkers of acute and chronic inflammation in vitro. <i>Journal of Agricultural and Food Chemistry</i> , 2014 , 62, 7022-8	5.7	114
47	Anthocyanin profiling of wild maqui berries (<i>Aristotelia chilensis</i> [Mol.] Stuntz) from different geographical regions in Chile. <i>Journal of the Science of Food and Agriculture</i> , 2014 , 94, 2639-48	4.3	67
46	Efficient preparative isolation and identification of walnut bioactive components using high-speed counter-current chromatography and LC-ESI-IT-TOF-MS. <i>Food Chemistry</i> , 2014 , 158, 229-38	8.5	61
45	Comparative analysis of phenolic content and profile, antioxidant capacity, and anti-inflammatory bioactivity in wild Alaskan and commercial <i>Vaccinium</i> berries. <i>Journal of Agricultural and Food Chemistry</i> , 2014 , 62, 4007-17	5.7	92
44	Leishmanicidal activity of a daucane sesquiterpene isolated from <i>Eryngium foetidum</i> . <i>Pharmaceutical Biology</i> , 2014 , 52, 398-401	3.8	14
43	Novel strategy to create hypoallergenic peanut protein-polyphenol edible matrices for oral immunotherapy. <i>Journal of Agricultural and Food Chemistry</i> , 2014 , 62, 7010-21	5.7	43
42	Quinoa seeds leach phytoecdysteroids and other compounds with anti-diabetic properties. <i>Food Chemistry</i> , 2014 , 163, 178-85	8.5	63
41	Phlorotannins from Alaskan seaweed inhibit carbolytic enzyme activity. <i>Marine Drugs</i> , 2014 , 12, 5277-946		59
40	Concentrating immunoprotective phytoactive compounds from fruits and vegetables into shelf-stable protein-rich ingredients. <i>Plant Foods for Human Nutrition</i> , 2014 , 69, 317-24	3.9	11

39	Bioactive capacity, sensory properties, and nutritional analysis of a shelf stable protein-rich functional ingredient with concentrated fruit and vegetable phytoactives. <i>Plant Foods for Human Nutrition</i> , 2014 , 69, 372-8	3.9	13
38	Cytotoxic effects of ellagitannins isolated from walnuts in human cancer cells. <i>Nutrition and Cancer</i> , 2014 , 66, 1304-14	2.8	23
37	Phytochemical changes in phenolics, anthocyanins, ascorbic acid, and carotenoids associated with sweetpotato storage and impacts on bioactive properties. <i>Food Chemistry</i> , 2014 , 145, 717-24	8.5	101
36	Antiplasmodial activity of cucurbitacin glycosides from <i>Datisca glomerata</i> (C. Presl) Baill. <i>Phytochemistry</i> , 2013 , 87, 78-85	4	9
35	Stable binding of alternative protein-enriched food matrices with concentrated cranberry bioflavonoids for functional food applications. <i>Journal of Agricultural and Food Chemistry</i> , 2013 , 61, 6856-64	5.7	45
34	Efficient quantification of the health-relevant anthocyanin and phenolic acid profiles in commercial cultivars and breeding selections of blueberries (<i>Vaccinium</i> spp.). <i>Journal of Agricultural and Food Chemistry</i> , 2013 , 61, 4806-15	5.7	71
33	Flavonoid glycosides and pharmacological activity of <i>Amphilophium paniculatum</i> . <i>Pharmacognosy Research (discontinued)</i> , 2013 , 5, 17-21	0.7	10
32	Bioactive polyphenols from muscadine grape and blackcurrant stably concentrated onto protein-rich matrices for topical applications. <i>International Journal of Cosmetic Science</i> , 2013 , 35, 394-407	2.7	26
31	Acute phytoecdysteroid treatment increases PI3k-Akt signaling in aged mouse skeletal muscle. <i>FASEB Journal</i> , 2013 , 27, 713.6	0.9	
30	Comparison of health-relevant flavonoids in commonly consumed cranberry products. <i>Journal of Food Science</i> , 2012 , 77, H176-83	3.4	56
29	In vitro antiplasmodial activity of indole alkaloids from the stem bark of <i>Geissospermum vellosii</i> . <i>Journal of Ethnopharmacology</i> , 2012 , 139, 471-7	5	36
28	Antiparasitic compounds from <i>Cornus florida</i> L. with activities against <i>Plasmodium falciparum</i> and <i>Leishmania tarentolae</i> . <i>Journal of Ethnopharmacology</i> , 2012 , 142, 456-61	5	28
27	Isolation and structural elucidation of indole alkaloids from <i>Geissospermum vellosii</i> by mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012 , 885-886, 83-9	3.2	8
26	Complementary approaches to gauge the bioavailability and distribution of ingested berry polyphenolics. <i>Journal of Agricultural and Food Chemistry</i> , 2012 , 60, 5763-71	5.7	41
25	Antiplasmodial Activity of the Ethnobotanical Plant <i>Cassia fistula</i> . <i>Natural Product Communications</i> , 2012 , 7, 1934578X1200701	0.9	8
24	Isolation and characterization of flavonols from blackcurrant by high-performance counter-current chromatography and electrospray ionization tandem mass spectrometry. <i>Journal of Separation Science</i> , 2012 , 35, 1682-9	3.4	13
23	and Anti-Diabetic Effects of Anthocyanins from Maqui Berry (). <i>Food Chemistry</i> , 2012 , 131, 387-396	8.5	146
22	Efficient sorption of polyphenols to soybean flour enables natural fortification of foods. <i>Food Chemistry</i> , 2012 , 131, 1193-1200	8.5	55

21	Antiplasmodial activity of the ethnobotanical plant <i>Cassia fistula</i> . <i>Natural Product Communications</i> , 2012 , 7, 1263-6	0.9	11
20	Antiplasmodial activity of aporphine alkaloids and sesquiterpene lactones from <i>Liriodendron tulipifera</i> L. <i>Journal of Ethnopharmacology</i> , 2011 , 133, 26-30	5	49
19	Isolation and identification of antiplasmodial N-alkylamides from <i>Spilanthes acmella</i> flowers using centrifugal partition chromatography and ESI-IT-TOF-MS. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2011 , 879, 1886-92	3.2	32
18	Antiplasmodial and Cytotoxic Activities of Drimane Sesquiterpenes from <i>Canella winterana</i> . <i>Natural Product Communications</i> , 2010 , 5, 1934578X1000501	0.9	2
17	Tracking deposition of a ¹⁴ C-radiolabeled kudzu hairy root-derived isoflavone-rich fraction into bone. <i>Experimental Biology and Medicine</i> , 2010 , 235, 1224-35	3.7	6
16	Antiplasmodial and cytotoxic activities of drimane sesquiterpenes from <i>Canella winterana</i> . <i>Natural Product Communications</i> , 2010 , 5, 1869-72	0.9	7
15	Hypoglycemic activity of a novel anthocyanin-rich formulation from lowbush blueberry, <i>Vaccinium angustifolium</i> Aiton. <i>Phytomedicine</i> , 2009 , 16, 406-15	6.5	161
14	Phytochemical Characterization of an Adaptogenic Preparation from <i>Rhodiola heterodonta</i> . <i>Natural Product Communications</i> , 2009 , 4, 1934578X0900400	0.9	2
13	Characteristic Flavonoids from <i>Acacia burkittii</i> and <i>A. Acuminata</i> Heartwoods and their Differential Cytotoxicity to Normal and Leukemia Cells. <i>Natural Product Communications</i> , 2009 , 4, 1934578X0900400	0.9	
12	Phytochemical characterization of an adaptogenic preparation from <i>Rhodiola heterodonta</i> . <i>Natural Product Communications</i> , 2009 , 4, 1053-8	0.9	13
11	Characteristic flavonoids from <i>Acacia burkittii</i> and <i>A. acuminata</i> heartwoods and their differential cytotoxicity to normal and leukemia cells. <i>Natural Product Communications</i> , 2009 , 4, 69-76	0.9	7
10	In vitro production of metabolism-enhancing phytoecdysteroids from <i>Ajuga turkestanica</i> . <i>Plant Cell, Tissue and Organ Culture</i> , 2008 , 93, 73-83	2.7	39
9	Neo-Clerodane Diterpenes from <i>Ajuga turkestanica</i> . <i>Phytochemistry Letters</i> , 2008 , 1, 81-84	1.9	15
8	ent-Beyerane diterpenoids from the heartwood of <i>Excoecaria parvifolia</i> . <i>Phytochemistry</i> , 2007 , 68, 546-54	4	7
7	Comparative phytochemical characterization of three <i>Rhodiola</i> species. <i>Phytochemistry</i> , 2006 , 67, 2380-91	4	83
6	Chemopreventive Potential of Flavonoid Extracts from Plantation-Bred and Wild <i>Aronia melanocarpa</i> (Black Chokeberry) Fruits. <i>Journal of Food Science</i> , 2006 , 71, C480-C488	3.4	45
5	Structures, biogenetic relationships, and cytotoxicity of pimarane-derived diterpenes from <i>Petalostigma pubescens</i> . <i>Phytochemistry</i> , 2006 , 67, 1708-15	4	25
4	Polyphenolics in <i>Rhizophora mangle</i> L. leaves and their changes during leaf development and senescence. <i>Trees - Structure and Function</i> , 2004 , 18, 518	2.6	98

3	Chemical composition and biological activity of the volatiles of <i>Anthemis melampodina</i> and <i>Pluchea dioscoridis</i> . <i>Phytotherapy Research</i> , 2002 , 16, 183-5	6.7	51
2	Polyphenols from <i>Cornulaca monacantha</i> . <i>Phytochemistry</i> , 2001 , 58, 611-3	4	16
1	A new eudesmanolide from <i>Crataegus flava</i> fruits. <i>Phytotherapy Research</i> , 2001 , 72, 756-9	3.2	6