

Edward Wargent

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6484563/publications.pdf>

Version: 2024-02-01

12
papers

456
citations

758635

12
h-index

1199166

12
g-index

12
all docs

12
docs citations

12
times ranked

914
citing authors

#	ARTICLE	IF	CITATIONS
1	Male mice that lack the G-protein-coupled receptor GPR41 have low energy expenditure and increased body fat content. <i>British Journal of Nutrition</i> , 2013, 109, 1755-1764.	1.2	106
2	Chronic Activation of $\hat{1}^3$ AMPK Induces Obesity and Reduces $\hat{1}^2$ Cell Function. <i>Cell Metabolism</i> , 2016, 23, 821-836.	7.2	87
3	Modulation of susceptibility to weight gain and insulin resistance in low birthweight rats by treatment of their mothers with leptin during pregnancy and lactation. <i>International Journal of Obesity</i> , 2004, 28, 129-136.	1.6	79
4	Preventive effects of <i>Salvia officinalis</i> leaf extract on insulin resistance and inflammation in a model of high fat diet-induced obesity in mice that responds to rosiglitazone. <i>PeerJ</i> , 2018, 6, e4166.	0.9	26
5	Contrasts between the effects of zinc- $\hat{1}^2$ -glycoprotein, a putative $\hat{1}^3/2$ -adrenoceptor agonist and the $\hat{1}^3/2$ -adrenoceptor agonist BRL35135 in C57Bl/6 (ob/ob) mice. <i>Journal of Endocrinology</i> , 2013, 216, 157-168.	1.2	25
6	5-HT _{2A} and 5-HT _{2C} receptors as hypothalamic targets of developmental programming in male rats. <i>DMM Disease Models and Mechanisms</i> , 2016, 9, 401-12.	1.2	25
7	Evidence from studies in rodents and in isolated adipocytes that agonists of the chemerin receptor CMKLR1 may be beneficial in the treatment of type 2 diabetes. <i>PeerJ</i> , 2015, 3, e753.	0.9	24
8	Identification of a novel agonist of peroxisome proliferator-activated receptors $\hat{1}^{\pm}$ and $\hat{1}^3$ that may contribute to the anti-diabetic activity of guggulipid in Lepob/Lepob mice. <i>Journal of Nutritional Biochemistry</i> , 2009, 20, 806-815.	1.9	21
9	Improvement of glucose tolerance in Zucker diabetic fatty rats by long-term treatment with the dipeptidyl peptidase inhibitor P32/98: comparison with and combination with rosiglitazone. <i>Diabetes, Obesity and Metabolism</i> , 2005, 7, 170-181.	2.2	17
10	A new, highly selective murine peroxisome proliferator-activated receptor $\hat{1}^{\pm}$ agonist increases responsiveness to thermogenic stimuli and glucose uptake in skeletal muscle in obese mice. <i>Diabetes, Obesity and Metabolism</i> , 2011, 13, 455-464.	2.2	17
11	Circulating levels of the cytokines IL10, IFN $\hat{1}^3$ and resistin in an obese mouse model of developmental programming. <i>Journal of Developmental Origins of Health and Disease</i> , 2013, 4, 491-498.	0.7	15
12	Leanness in postnatally nutritionally programmed rats is associated with increased sensitivity to leptin and a melanocortin receptor agonist and decreased sensitivity to neuropeptide Y. <i>International Journal of Obesity</i> , 2012, 36, 1040-1046.	1.6	14