

Ryoichi Banno

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6485017/publications.pdf>

Version: 2024-02-01

16
papers

425
citations

1163117

8
h-index

996975

15
g-index

16
all docs

16
docs citations

16
times ranked

541
citing authors

#	ARTICLE	IF	CITATIONS
1	Patients With Antithyroid Antibodies Are Prone To Develop Destructive Thyroiditis by Nivolumab: A Prospective Study. <i>Journal of the Endocrine Society</i> , 2018, 2, 241-251.	0.2	146
2	Pituitary dysfunction induced by immune checkpoint inhibitors is associated with better overall survival in both malignant melanoma and non-small cell lung carcinoma: a prospective study. , 2020, 8, e000779.		75
3	Anti-thyroid antibodies and thyroid echo pattern at baseline as risk factors for thyroid dysfunction induced by anti-programmed cell death-1 antibodies: a prospective study. <i>British Journal of Cancer</i> , 2020, 122, 771-777.	6.4	48
4	CD4 ⁺ T cells are essential for the development of destructive thyroiditis induced by anti- α PD-1 antibody in thyroglobulin-immunized mice. <i>Science Translational Medicine</i> , 2021, 13, .	12.4	47
5	Anti-pituitary antibodies against corticotrophs in IgG4-related hypophysitis. <i>Pituitary</i> , 2017, 20, 301-310.	2.9	21
6	Critical role of rabphilin α 3A in the pathophysiology of experimental lymphocytic neurohypophysitis. <i>Journal of Pathology</i> , 2018, 244, 469-478.	4.5	20
7	Hypothalamic glial cells isolated by MACS reveal that microglia and astrocytes induce hypothalamic inflammation via different processes under high-fat diet conditions. <i>Neurochemistry International</i> , 2020, 136, 104733.	3.8	15
8	Increased Risk of Thyroid Dysfunction by PD-1 and CTLA-4 Blockade in Patients Without Thyroid Autoantibodies at Baseline. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, e1620-e1630.	3.6	15
9	GABAB Receptor Signaling in the Mesolimbic System Suppresses Binge-like Consumption of a High-Fat Diet. <i>IScience</i> , 2019, 20, 337-347.	4.1	10
10	High-fat Feeding Causes Inflammation and Insulin Resistance in the Ventral Tegmental Area in Mice. <i>Neuroscience</i> , 2021, 461, 72-79.	2.3	8
11	d-Allulose Ameliorates Skeletal Muscle Insulin Resistance in High-Fat Diet-Fed Rats. <i>Molecules</i> , 2021, 26, 6310.	3.8	7
12	Protein Tyrosine Phosphatase 1B Deficiency Improves Glucose Homeostasis in Type 1 Diabetes Treated With Leptin. <i>Diabetes</i> , 2022, 71, 1902-1914.	0.6	5
13	Basigin deficiency prevents anaplerosis and ameliorates insulin resistance and hepatosteatosis. <i>JCI Insight</i> , 2021, 6, .	5.0	3
14	Arginine vasopressin-Venus reporter mice as a tool for studying magnocellular arginine vasopressin neurons. <i>Peptides</i> , 2021, 139, 170517.	2.4	2
15	Peripheral combination treatment of leptin and an SGLT2 inhibitor improved glucose metabolism in insulin-dependent diabetes mellitus mice. <i>Journal of Pharmacological Sciences</i> , 2021, 147, 340-347.	2.5	2
16	d-Allulose Improves Endurance and Recovery from Exhaustion in Male C57BL/6J Mice. <i>Nutrients</i> , 2022, 14, 404.	4.1	1