

Charles W Kaspar

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6495883/charles-w-kaspar-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

21
papers

601
citations

12
h-index

21
g-index

21
ext. papers

692
ext. citations

3.6
avg, IF

3.1
L-index

#	Paper	IF	Citations
21	A Single Nucleotide Polymorphism in Increases Tolerance to Bile Salts, Acid, and Staining of Calcofluor-Binding Polysaccharides in Serovar Typhimurium E40. <i>Frontiers in Microbiology</i> , 2021 , 12, 671453	5.7	53
20	Genome Sequences of 14 Escherichia coli O157:H7 Strains Isolated before and during the Time Frame of the 2018 Multistate Outbreak Associated with Romaine Lettuce. <i>Microbiology Resource Announcements</i> , 2020 , 9,	1.3	1
19	Chronological set of E. coli O157:H7 bovine strains establishes a role for repeat sequences and mobile genetic elements in genome diversification. <i>BMC Genomics</i> , 2020 , 21, 562	4.5	2
18	Evaluating the efficacy of beef slaughter line interventions by quantifying the six major non-O157 Shiga toxin producing Escherichia coli serogroups using real-time multiplex PCR. <i>Food Microbiology</i> , 2017 , 63, 228-238	6	9
17	Gene markers of generic Escherichia coli associated with colonization and persistence of Escherichia coli O157 in cattle. <i>Preventive Veterinary Medicine</i> , 2014 , 117, 140-8	3.1	2
16	Phylogenetic characterization of Escherichia coli O157 : H7 based on IS629 distribution and Shiga toxin genotype. <i>Microbiology (United Kingdom)</i> , 2014 , 160, 502-513	2.9	9
15	Response of Medicago truncatula seedlings to colonization by Salmonella enterica and Escherichia coli O157:H7. <i>PLoS ONE</i> , 2014 , 9, e87970	3.7	12
14	Prevalent and persistent Escherichia coli O157:H7 strains on farms are selected by bovine passage. <i>Veterinary Microbiology</i> , 2013 , 162, 912-920	3.3	11
13	Evolution of the Stx2-encoding prophage in persistent bovine Escherichia coli O157:H7 strains. <i>Applied and Environmental Microbiology</i> , 2013 , 79, 1563-72	4.8	22
12	Osmotic and desiccation tolerance in Escherichia coli O157:H7 requires rpoS ((B8)). <i>Current Microbiology</i> , 2012 , 65, 660-5	2.4	16
11	Differences in colonization and shedding patterns after oral challenge of cattle with three Escherichia coli O157:H7 strains. <i>Applied and Environmental Microbiology</i> , 2012 , 78, 8045-55	4.8	17
10	Reduction of Escherichia coli O157:H7 shedding in cattle by addition of chitosan microparticles to feed. <i>Applied and Environmental Microbiology</i> , 2011 , 77, 2611-6	4.8	28
9	Acid stress damage of DNA is prevented by Dps binding in Escherichia coli O157:H7. <i>BMC Microbiology</i> , 2008 , 8, 181	4.5	67
8	Molybdate treatment and sulfate starvation decrease ATP and DNA levels in Ferroplasma acidarmanu. <i>Archaea</i> , 2008 , 2, 205-9	2	6
7	Production of methanethiol and volatile sulfur compounds by the archaeon "Ferroplasma acidarmanus". <i>Extremophiles</i> , 2007 , 11, 841-51	3	15
6	Sulfate requirement for heterotrophic growth of "Ferroplasma acidarmanus" strain fer1. <i>Research in Microbiology</i> , 2005 , 156, 492-8	4	13
5	Temperature-dependent genome degradation in the coccoid form of Campylobacter jejuni. <i>Current Microbiology</i> , 2005 , 50, 110-3	2.4	7

4	Tetraether-linked membrane monolayers in <i>Ferroplasma</i> spp: a key to survival in acid. <i>Extremophiles</i> , 2004 , 8, 411-9	3	124
3	Contribution of <i>dps</i> to acid stress tolerance and oxidative stress tolerance in <i>Escherichia coli</i> O157:H7. <i>Applied and Environmental Microbiology</i> , 2000 , 66, 3911-6	4.8	122
2	Role of <i>rpoS</i> in acid resistance and fecal shedding of <i>Escherichia coli</i> O157:H7. <i>Applied and Environmental Microbiology</i> , 2000 , 66, 632-7	4.8	90
1	Genomic analysis using pulsed-field gel electrophoresis of <i>Escherichia coli</i> O157: H7 isolated from dairy calves during the United State National Dairy Heifer Evaluation Project (1992-1992). <i>Veterinary Microbiology</i> , 1996 , 48, 223-30	3.3	28