Catherine M Sackley

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6501687/publications.pdf

Version: 2024-02-01

140 papers 4,629 citations

32 h-index 61 g-index

145 all docs

145 docs citations

145 times ranked 5558 citing authors

#	Article	IF	CITATIONS
1	Physiotherapy intervention in Parkinson's disease: systematic review and meta-analysis. BMJ, The, 2012, 345, e5004-e5004.	3.0	250
2	Physiotherapy versus placebo or no intervention in Parkinson's disease. The Cochrane Library, 2013, , CD002817.	1.5	235
3	Effectiveness of physiotherapy exercise after knee arthroplasty for osteoarthritis: systematic review and meta-analysis of randomised controlled trials. BMJ: British Medical Journal, 2007, 335, 812.	2.4	225
4	The Prevalence of Joint Contractures, Pressure Sores, Painful Shoulder, Other Pain, Falls, and Depression in the Year After a Severely Disabling Stroke. Stroke, 2008, 39, 3329-3334.	1.0	195
5	Effectiveness of physiotherapy exercise following total knee replacement: systematic review and meta-analysis. BMC Musculoskeletal Disorders, 2015, 16, 15.	0.8	178
6	A Population-Based Study of the Prevalence of Fatigue After Transient Ischemic Attack and Minor Stroke. Stroke, 2009, 40, 757-761.	1.0	138
7	Single blind randomized controlled trial of visual feedback after stroke: Effects on stance symmetry and function. Disability and Rehabilitation, 1997, 19, 536-546.	0.9	134
8	Falls, sway, and symmetry of weight-bearing after stroke. International Disability Studies, 1991, 13, 1-4.	0.4	131
9	Fatigue, psychological and cognitive impairment following transient ischaemic attack and minor stroke: a systematic review. European Journal of Neurology, 2014, 21, 1258-1267.	1.7	129
10	Occupational therapy for patients with problems in personal activities of daily living after stroke: systematic review of randomised trials. BMJ: British Medical Journal, 2007, 335, 922.	2.4	127
11	Priority setting partnership to identify the top 10 research priorities for the management of Parkinson's disease. BMJ Open, 2014, 4, e006434.	0.8	120
12	Effectiveness of physiotherapy exercise following hip arthroplasty for osteoarthritis: a systematic review of clinical trials. BMC Musculoskeletal Disorders, 2009, 10, 98.	0.8	112
13	Physiotherapy for Parkinson's disease: a comparison of techniques. The Cochrane Library, 2014, , CD002815.	1.5	100
14	Use of information feedback and attentional focus of feedback in treating the person with a hemiplegic arm. Physiotherapy Research International, 2009, 14, 77-90.	0.7	91
15	Prognostic factors of functional outcome after hip fracture surgery: a systematic review. Age and Ageing, 2018, 47, 661-670.	0.7	90
16	Physiotherapy treatment for stroke patients: A survey of current practice. Physiotherapy Theory and Practice, 1996, 12, 87-96.	0.6	82
17	The relationships between weight-bearing asymmetry after stroke, motor function and activities of daily living. Physiotherapy Theory and Practice, 1990, 6, 179-185.	0.6	74
18	Physiotherapy and Occupational Therapy vs No Therapy in Mild to Moderate Parkinson Disease. JAMA Neurology, 2016, 73, 291.	4.5	73

#	Article	IF	Citations
19	Physiotherapy versus placebo or no intervention in Parkinson's disease., 2012, , CD002817.		72
20	Clinical effectiveness and cost-effectiveness of physiotherapy and occupational therapy versus no therapy in mild to moderate Parkinson's disease: a large pragmatic randomised controlled trial (PD) Tj ETQ	q0 01 0 3rgB1	Г/О ве rlock 10
21	Cluster Randomized Pilot Controlled Trial of an Occupational Therapy Intervention for Residents With Stroke in UK Care Homes. Stroke, 2006, 37, 2336-2341.	1.0	59
22	Home-based therapy programmes for upper limb functional recovery following stroke. The Cochrane Library, 2012, , CD006755.	1.5	52
23	Towards a programme theory for fidelity in the evaluation of complex interventions. Journal of Evaluation in Clinical Practice, 2014, 20, 445-452.	0.9	52
24	A study of perceived facilitators to physical activity in neurological conditions. International Journal of Therapy and Rehabilitation, 2009, 16, 17-24.	0.1	51
25	Effects of a physiotherapy and occupational therapy intervention on mobility and activity in care home residents: a cluster randomised controlled trial. BMJ: British Medical Journal, 2009, 339, b3123-b3123.	2.4	51
26	Health-related quality of life and supportive care in patients with rare long-term neurological conditions. Quality of Life Research, 2013, 22, 1231-1238.	1.5	50
27	An occupational therapy intervention for residents with stroke related disabilities in UK care homes (OTCH): cluster randomised controlled trial. BMJ, The, 2015, 350, h468-h468.	3.0	47
28	Treatment of chronic back pain by sensory discrimination training. A Phase I RCT of a novel device (FairMed) vs. TENS. BMC Musculoskeletal Disorders, 2008, 9, 97.	0.8	44
29	Rehabilitation interventions for foot drop in neuromuscular disease. , 2009, , CD003908.		44
30	Physiotherapy versus placebo or no intervention in Parkinson's disease., 2012, , CD002817.		40
31	Safety and efficacy of co-careldopa as an add-on therapy to occupational and physical therapy in patients after stroke (DARS): a randomised, double-blind, placebo-controlled trial. Lancet Neurology, The, 2019, 18, 530-538.	4.9	39
32	Pedometer step counts in individuals with neurological conditions. Clinical Rehabilitation, 2009, 23, 171-175.	1.0	38
33	Effectiveness of land-based physiotherapy exercise following hospital discharge following hip arthroplasty for osteoarthritis: an updated systematic review. Physiotherapy, 2015, 101, 252-265.	0.2	38
34	Physiotherapy versus placebo or no intervention in Parkinson's disease., 2001, , CD002817.		37
35	Attentional focus of feedback for improving performance of reach-to-grasp after stroke: a randomised crossover study. Physiotherapy, 2014, 100, 108-115.	0.2	37
36	Comparison of speech and language therapy techniques for speech problems in Parkinson's disease. The Cochrane Library, 2012, , CD002814.	1.5	36

#	Article	IF	Citations
37	A pilot randomized controlled trial comparing effectiveness of prism glasses, visual search training and standard care in hemianopia. Acta Neurologica Scandinavica, 2017, 136, 310-321.	1.0	36
38	Feasibility and Preliminary Efficacy of Visual Cue Training to Improve Adaptability of Walking after Stroke: Multi-Centre, Single-Blind Randomised Control Pilot Trial. PLoS ONE, 2015, 10, e0139261.	1.1	36
39	Inequity in rehabilitation interventions after hip fracture: a systematic review. Age and Ageing, 2019, 48, 489-497.	0.7	34
40	Speech and language therapy versus placebo or no intervention for speech problems in Parkinson's disease. The Cochrane Library, 2012, 2012, CD002812.	1.5	32
41	Community occupational therapy for people with dementia and family carers (COTiD-UK) versus treatment as usual (Valuing Active Life in Dementia [VALID] programme): study protocol for a randomised controlled trial. Trials, 2016, 17, 65.	0.7	31
42	A phase II exploratory cluster randomized controlled trial of a group mobility training and staff education intervention to promote urinary continence in UK care homes. Clinical Rehabilitation, 2008, 22, 714-721.	1.0	30
43	Discharge after hip fracture surgery by mobilisation timing: secondary analysis of the UK National Hip Fracture Database. Age and Ageing, 2021, 50, 415-422.	0.7	29
44	An exploratory cluster randomized controlled trial of group exercise on mobility and depression in care home residents. Clinical Rehabilitation, 2009, 23, 146-154.	1.0	28
45	Lee Silverman Voice Treatment versus standard speech and language therapy versus control in Parkinson's disease: a pilot randomised controlled trial (PD COMM pilot). Pilot and Feasibility Studies, 2018, 4, 30.	0.5	27
46	The effect of rehabilitation interventions on physical function and immobility-related complications in severe stroke: a systematic review. BMJ Open, 2020, 10, e033642.	0.8	27
47	Recovery of function following hip resurfacing arthroplasty: a randomized controlled trial comparing an accelerated versus standard physiotherapy rehabilitation programme. Clinical Rehabilitation, 2013, 27, 771-784.	1.0	26
48	Fracture in the Elderly Multidisciplinary Rehabilitation (FEMuR): a phase II randomised feasibility study of a multidisciplinary rehabilitation package following hip fracture. BMJ Open, 2016, 6, e012422.	0.8	26
49	UK survey of occupational therapist's and physiotherapist's experiences and attitudes towards hip replacement precautions and equipment. BMC Musculoskeletal Disorders, 2016, 17, 228.	0.8	25
50	Is there a difference in physical activity levels in patients before and up to one year after unilateral total hip replacement? A systematic review and meta-analysis. Clinical Rehabilitation, 2017, 31, 639-650.	1.0	25
51	Metronome Cueing of Walking Reduces Gait Variability after a Cerebellar Stroke. Frontiers in Neurology, 2016, 7, 84.	1.1	24
52	Assistive devices, hip precautions, environmental modifications and training to prevent dislocation and improve function after hip arthroplasty. The Cochrane Library, 2016, 2016, CD010815.	1.5	24
53	Occupational Therapy in Nursing and Residential Care Settings: A Description of a Randomised Controlled Trial Intervention. British Journal of Occupational Therapy, 2004, 67, 104-110.	0.5	23
54	The effect of rehabilitation interventions on physical function and immobility-related complications in severe strokeâ€"protocol for a systematic review. Systematic Reviews, 2018, 7, 197.	2.5	23

#	Article	IF	CITATIONS
55	Practical research-based guidance for motor imagery practice in neurorehabilitation. Disability and Rehabilitation, 2012, 34, 2192-2200.	0.9	22
56	A systematic review of the efficiency of recruitment to stroke rehabilitation randomised controlled trials. Trials, 2020, 21, 68.	0.7	22
57	Comparison of postdischarge physiotherapy versus usual care following primary total knee arthroplasty for osteoarthritis: an exploratory pilot randomized clinical trial. Clinical Rehabilitation, 2012, 26, 629-641.	1.0	21
58	A randomized controlled trial to compare two methods of constraint-induced movement therapy to improve functional ability in the affected upper limb in pre-school children with hemiplegic cerebral palsy: CATCH TRIAL. Clinical Rehabilitation, 2018, 32, 909-918.	1.0	21
59	Long-Term Trends in Stroke Survivors Discharged to Care Homes. Stroke, 2020, 51, 179-185.	1.0	21
60	Effect of a Home-Based, Walking Exercise Behavior Change Intervention vs Usual Care on Walking in Adults With Peripheral Artery Disease. JAMA - Journal of the American Medical Association, 2022, 327, 1344.	3.8	21
61	Metronome-Cued Stepping in Place after Hemiparetic Stroke: Comparison of a One- and Two-Tone Beat. ISRN Rehabilitation, 2013, 2013, 1-5.	0.6	20
62	Lee Silverman voice treatment versus standard NHS speech and language therapy versus control in Parkinson's disease (PD COMM pilot): study protocol for a randomized controlled trial. Trials, 2014, 15, 213.	0.7	20
63	The DARS (Dopamine Augmented Rehabilitation in Stroke) trial: protocol for a randomised controlled trial of Co-careldopa treatment in addition to routine NHS occupational and physical therapy after stroke. Trials, 2014, 15, 316.	0.7	19
64	Huntington's Disease: Current Epidemiology and Pharmacological Management in UK Primary Care. Neuroepidemiology, 2011, 37, 216-221.	1.1	18
65	Developing a multidisciplinary rehabilitation package following hip fracture and testing in a randomised feasibility study: Fracture in the Elderly Multidisciplinary Rehabilitation (FEMuR). Health Technology Assessment, 2017, 21, 1-528.	1.3	18
66	Improving patients' experience and outcome of total joint replacement: the RESTORE programme. Programme Grants for Applied Research, 2016, 4, 1-508.	0.4	18
67	Comparison of speech and language therapy techniques for speech problems in Parkinson's disease. , 2001, , CD002814.		17
68	Rehabilitation interventions for foot drop in neuromuscular disease., 2007,, CD003908.		17
69	Does physical activity change following hip and knee replacement? Matched case-control study evaluating Physical Activity Scale for the Elderly data from the Osteoarthritis Initiative. Physiotherapy, 2018, 104, 80-90.	0.2	17
70	Development of an evidence-based complex intervention for community rehabilitation of patients with hip fracture using realist review, survey and focus groups. BMJ Open, 2017, 7, e014362.	0.8	17
71	Stepping to the Beat: Feasibility and Potential Efficacy of a Home-Based Auditory-Cued Step Training Program in Chronic Stroke. Frontiers in Neurology, 2017, 8, 412.	1.1	16
72	Community Occupational Therapy for people with dementia and family carers (COTiD-UK) versus treatment as usual (Valuing Active Life in Dementia [VALID]) study: A single-blind, randomised controlled trial. PLoS Medicine, 2021, 18, e1003433.	3.9	16

#	Article	IF	Citations
73	Subjective Experiences of Speech and Language Therapy in Patients with Parkinson's Disease: A Pilot Study. Rehabilitation Research and Practice, 2015, 2015, 1-11.	0.5	15
74	Home-based Reach-to-Grasp training for people after stroke is feasible: a pilot randomised controlled trial. Clinical Rehabilitation, 2017, 31, 891-903.	1.0	15
75	Short-term effects on mobility, activities of daily living and health-related quality of life of a Conductive Education programme for adults with multiple sclerosis, Parkinson's disease and stroke. Clinical Rehabilitation, 2008, 22, 329-337.	1.0	13
76	Occupational therapy for care home residents with stroke. The Cochrane Library, 2013, , CD010116.	1.5	13
77	Delayed Recovery of Leg Fatigue Symptoms Following a Maximal Exercise Session in People With Multiple Sclerosis. Neurorehabilitation and Neural Repair, 2014, 28, 139-148.	1.4	13
78	Stroke patients entering nursing home care: a content analysis of discharge letters. Clinical Rehabilitation, 2002, 16, 736-740.	1.0	12
79	Visual cue training to improve walking and turning after stroke: a study protocol for a multi-centre, single blind randomised pilot trial. Trials, 2013, 14, 276.	0.7	12
80	A feasibility randomised controlled trial of pre-operative occupational therapy to optimise recovery for patients undergoing primary total hip replacement for osteoarthritis (PROOF-THR). Clinical Rehabilitation, 2016, 30, 156-166.	1.0	12
81	Conceptual Framework for an Episode of Rehabilitative Care After Surgical Repair of Hip Fracture. Physical Therapy, 2019, 99, 276-285.	1.1	12
82	Rating the intelligibility of dysarthic speech amongst people with Parkinson's Disease: a comparison of trained and untrained listeners. Clinical Linguistics and Phonetics, 2019, 33, 1063-1070.	0.5	12
83	Functional and emotional outcomes after transient ischemic attack: A 12-month prospective controlled cohort study. International Journal of Stroke, 2019, 14, 522-529.	2.9	12
84	An investigation into the utility of the Stroke Aphasic Depression Questionnaire (SADQ) in care home settings. Clinical Rehabilitation, 2006, 20, 598-602.	1.0	11
85	Occupational Therapy to Optimise Independence in Parkinson's Disease: The Designing and Recording of a Randomised Controlled Trial Intervention. British Journal of Occupational Therapy, 2010, 73, 178-185.	0.5	11
86	Reaching the †hard to reach': strategies to recruit black and minority ethnic service users with rare long-term neurological conditions. International Journal of Social Research Methodology: Theory and Practice, 2012, 15, 485-495.	2.3	11
87	Subjective experiences of transient ischaemic attack: a repertory grid approach. Disability and Rehabilitation, 2013, 35, 2205-2212.	0.9	11
88	A randomised controlled trial to compare the clinical and cost-effectiveness of prism glasses, visual search training and standard care in patients with hemianopia following stroke: a protocol. BMJ Open, 2014, 4, e005885-e005885.	0.8	11
89	An Occupational Therapy intervention for residents with stroke-related disabilities in UK Care Homes (OTCH): cluster randomised controlled trial with economic evaluation. Health Technology Assessment, 2016, 20, 1-138.	1.3	11
90	A group memory rehabilitation programme for people with traumatic brain injuries: the ReMemBrIn RCT. Health Technology Assessment, 2019, 23, 1-194.	1.3	11

#	Article	IF	CITATIONS
91	Hydration and independence in activities of daily living in people with multiple sclerosis: a pilot investigation. Disability and Rehabilitation, 2011, 33, 1822-1825.	0.9	10
92	Fracture in the Elderly Multidisciplinary Rehabilitation (FEMuR): study protocol for a phase II randomised feasibility study of a multidisciplinary rehabilitation package following hip fracture [ISRCTN22464643]. Pilot and Feasibility Studies, 2015, 1, 13.	0.5	10
93	Orthopaedic physiotherapists' perceptions of mechanisms for observed variation in the implementation of physiotherapy practices in the early postoperative phase after hip fracture: a UK qualitative study. Age and Ageing, 2021, 50, 1961-1970.	0.7	10
94	The effectiveness of rehabilitation interventions including outdoor mobility on older adults' physical activity, endurance, outdoor mobility and falls-related self-efficacy: systematic review and meta-analysis. Age and Ageing, 2022, 51, .	0.7	10
95	A systematic review investigating fatigue, psychological and cognitive impairment following TIA and minor stroke: protocol paper. Systematic Reviews, 2013, 2, 72.	2.5	9
96	Rehabilitation of Memory following Brain Injury (ReMemBrIn): study protocol for a randomised controlled trial. Trials, 2015, 16, 6.	0.7	9
97	Changes in physical activity following total hip or knee arthroplasty: a matched case-control study from the EPIC-Norfolk cohort. Clinical Rehabilitation, 2017, 31, 1548-1557.	1.0	9
98	Clinical and cost effectiveness of memory rehabilitation following traumatic brain injury: a pragmatic cluster randomized controlled trial. Clinical Rehabilitation, 2019, 33, 1171-1184.	1.0	9
99	Visual Function Questionnaire as an outcome measure for homonymous hemianopia: subscales and supplementary questions, analysis from the VISION trial. Eye, 2019, 33, 1485-1493.	1.1	9
100	Technology-enhanced learning in physiotherapy education: Student satisfaction and knowledge acquisition of entry-level students in the United Kingdom. Research in Learning Technology, 2019, 27, .	2.3	9
101	Nominal group technique to establish the core components of home-based rehabilitation for survivors of stroke with severe disability. BMJ Open, 2021, 11, e052593.	0.8	9
102	VISTA-Rehab: A Resource for Stroke Rehabilitation Trials. International Journal of Stroke, 2010, 5, 447-452.	2.9	8
103	A Randomised Controlled Trial of Treatment for Post-Stroke Homonymous Hemianopia: Screening and Recruitment. Neuro-Ophthalmology, 2016, 40, 1-7.	0.4	8
104	Survival and outcomes for stroke survivors living in care homes: a prospective cohort study. Age and Ageing, 2021, 50, 2079-2087.	0.7	7
105	Discharge after hip fracture surgery in relation to mobilisation timing by patient characteristics: linked secondary analysis of the UK National Hip Fracture Database. BMC Geriatrics, 2021, 21, 694.	1.1	7
106	The Hospital Anxiety and Depression Scale: a pilot study to examine its latent structure and the link between psychological state and symptom severity in transient ischaemic attack patients. Psychology, Health and Medicine, 2016, 21, 632-638.	1.3	6
107	The PD COMM trial: a protocol for the process evaluation of a randomised trial assessing the effectiveness of two types of SLT for people with Parkinson's disease. Trials, 2017, 18, 397.	0.7	6
108	Finding the right outcome measures for care home research. Age and Ageing, 2010, 39, 119-122.	0.7	5

#	Article	lF	CITATIONS
109	Motivating Structured walking Activity in people with Intermittent Claudication (MOSAIC): protocol for a randomised controlled trial of a physiotherapist-led, behavioural change intervention versus usual care in adults with intermittent claudication. BMJ Open, 2019, 9, e030002.	0.8	5
110	The rehabilitation of physical function after severely disabling stroke: a survey of UK therapist practice. International Journal of Therapy and Rehabilitation, 2021, 28, 1-25.	0.1	5
111	Using the Barthel Index and modified Rankin Scale as Outcome Measures for Stroke Rehabilitation Trials; A Comparison of Minimum Sample Size Requirements. Journal of Stroke and Cerebrovascular Diseases, 2022, 31, 106229.	0.7	5
112	Occupational Therapy for Care Home Residents with Stroke: What is Routine National Practice?. British Journal of Occupational Therapy, 2014, 77, 265-273.	0.5	4
113	Conductive Education as a Method of Stroke Rehabilitation: A Single Blinded Randomised Controlled Feasibility Study. Stroke Research and Treatment, 2016, 2016, 1-6.	0.5	4
114	Do changes in outcomes following primary and revision hip replacement differ and relate to markers of socioeconomic status? A 1-year population-based cohort study. Monthly Notices of the Royal Astronomical Society: Letters, 2022, 93, 397-404.	1.2	4
115	178â€fDoes Physical Activity Change Following Hip and Knee Replacement? An Analysis of Data from the Osteoarthritis Initiative. Rheumatology, 2016, 55, i134-i134.	0.9	3
116	Physiotherapy and Occupational Therapy and Mild to Moderate Parkinson Diseaseâ€"Reply. JAMA Neurology, 2016, 73, 894.	4.5	3
117	Recruitment challenges in stroke rehabilitation randomized controlled trials: a qualitative exploration of trialists' perspectives using Framework analysis. Clinical Rehabilitation, 2020, 34, 1122-1133.	1.0	3
118	Effectiveness of community-based rehabilitation interventions incorporating outdoor mobility on ambulatory ability and falls-related self-efficacy after hip fracture: a systematic review and meta-analysis. Archives of Osteoporosis, 2021, 16, 99.	1.0	3
119	Letter to the Editor. Clinical Orthopaedics and Related Research, 2013, 471, 3706-3707.	0.7	2
120	An occupational therapy intervention for residents with stroke living in care homes in the United Kingdom: A content analysis of occupational therapy records from the OTCH trial. British Journal of Occupational Therapy, 2015, 78, 422-430.	0.5	1
121	Rehabilitation interventions for foot drop in neuromuscular disease. The Cochrane Library, 2015, , CD003908.	1.5	1
122	Stepping to an Auditory Metronome Improves Weight-Bearing Symmetry in Poststroke Hemiparesis. Journal of Applied Biomechanics, 2018, 34, 469-473.	0.3	1
123	A multicentre, randomised controlled trial to compare the clinical and cost-effectiveness of Lee Silverman Voice Treatment versus standard NHS Speech and Language Therapy versus control in Parkinson's disease: a study protocol for a randomised controlled trial. Trials, 2020, 21, 436.	0.7	1
124	Dopamine Augmented Rehabilitation in Stroke (DARS): a multicentre double-blind, randomised controlled trial of co-careldopa compared with placebo, in addition to routine NHS occupational and physical therapy, delivered early after stroke on functional recovery. Efficacy and Mechanism Evaluation, 2019, 6, 1-138.	0.9	1
125	Exploring the experiences of stroke survivors, informal caregivers and healthcare providers in Sierra Leone: a qualitative study protocol. BMJ Open, 2021, 11, e051276.	0.8	1
126	Gait rehabilitation for foot and ankle impairments in early rheumatoid arthritis: a feasibility study of a new gait rehabilitation programme (GREAT Strides). Pilot and Feasibility Studies, 2022, 8, .	0.5	1

#	Article	IF	CITATIONS
127	PD REHAB RCT OF PHYSIOTHERAPY AND OT IN PARKINSON'S DISEASE. Journal of Neurology, Neurosurgery and Psychiatry, 2014, 85, e4.124-e4.	0.9	O
128	Conductive Education as a Method of Stroke Rehabilitation: A Single Blind Randomised Controlled Feasibility Study. Archives of Physical Medicine and Rehabilitation, 2017, 98, e48.	0.5	0
129	Choice of outcome measures for the VISION pilot trial of interventions for hemianopia. Acta Neurologica Scandinavica, 2017, 136, 551-553.	1.0	0
130	P105â€fTherapists acceptability of delivering a psychologically informed gait rehabilitation intervention in early rheumatoid arthritis (GREAT): a qualitative interview study. Rheumatology, 2020, 59, .	0.9	0
131	P125 A mixed methods feasibility study of a gait rehabilitation programme for people with early rheumatoid arthritis and foot pain. Rheumatology, 2020, 59, .	0.9	0
132	P221 $\hat{a} \in f$ Treatment fidelity in the Gait Rehabilitation in Early Rheumatoid Arthritis Trial (GREAT) feasibility study. Rheumatology, 2020, 59, .	0.9	0
133	Protocol for a definitive randomised controlled trial and economic evaluation of a community-based rehabilitation programme following hip fracture: fracture in the elderly multidisciplinary rehabilitation-phase III (FEMuR III). BMJ Open, 2020, 10, e039791.	0.8	0
134	Title is missing!. , 2021, 18, e1003433.		0
135	Title is missing!. , 2021, 18, e1003433.		0
136	Title is missing!. , 2021, 18, e1003433.		0
137	Title is missing!. , 2021, 18, e1003433.		0
138	Title is missing!. , 2021, 18, e1003433.		0
139	Title is missing!. , 2021, 18, e1003433.		0
140	Speech and language therapy interventions for speech problems in Parkinson's disease. The Cochrane Library, 2022, 2022, .	1.5	0