## Catherine M Sackley

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6501687/publications.pdf

Version: 2024-02-01

140 papers 4,629 citations

32 h-index 61 g-index

145 all docs

145 docs citations

145 times ranked 5558 citing authors

#	Article	IF	CITATIONS
1	Using the Barthel Index and modified Rankin Scale as Outcome Measures for Stroke Rehabilitation Trials; A Comparison of Minimum Sample Size Requirements. Journal of Stroke and Cerebrovascular Diseases, 2022, 31, 106229.	0.7	5
2	Do changes in outcomes following primary and revision hip replacement differ and relate to markers of socioeconomic status? A 1-year population-based cohort study. Monthly Notices of the Royal Astronomical Society: Letters, 2022, 93, 397-404.	1.2	4
3	Effect of a Home-Based, Walking Exercise Behavior Change Intervention vs Usual Care on Walking in Adults With Peripheral Artery Disease. JAMA - Journal of the American Medical Association, 2022, 327, 1344.	3.8	21
4	Gait rehabilitation for foot and ankle impairments in early rheumatoid arthritis: a feasibility study of a new gait rehabilitation programme (GREAT Strides). Pilot and Feasibility Studies, 2022, 8, .	0.5	1
5	The effectiveness of rehabilitation interventions including outdoor mobility on older adults' physical activity, endurance, outdoor mobility and falls-related self-efficacy: systematic review and meta-analysis. Age and Ageing, 2022, 51, .	0.7	10
6	Speech and language therapy interventions for speech problems in Parkinson's disease. The Cochrane Library, 2022, 2022, .	1.5	0
7	Discharge after hip fracture surgery by mobilisation timing: secondary analysis of the UK National Hip Fracture Database. Age and Ageing, 2021, 50, 415-422.	0.7	29
8	Community Occupational Therapy for people with dementia and family carers (COTiD-UK) versus treatment as usual (Valuing Active Life in Dementia [VALID]) study: A single-blind, randomised controlled trial. PLoS Medicine, 2021, 18, e1003433.	3.9	16
9	Orthopaedic physiotherapists' perceptions of mechanisms for observed variation in the implementation of physiotherapy practices in the early postoperative phase after hip fracture: a UK qualitative study. Age and Ageing, 2021, 50, 1961-1970.	0.7	10
10	Effectiveness of community-based rehabilitation interventions incorporating outdoor mobility on ambulatory ability and falls-related self-efficacy after hip fracture: a systematic review and meta-analysis. Archives of Osteoporosis, 2021, 16, 99.	1.0	3
11	Survival and outcomes for stroke survivors living in care homes: a prospective cohort study. Age and Ageing, 2021, 50, 2079-2087.	0.7	7
12	The rehabilitation of physical function after severely disabling stroke: a survey of UK therapist practice. International Journal of Therapy and Rehabilitation, 2021, 28, 1-25.	0.1	5
13	Nominal group technique to establish the core components of home-based rehabilitation for survivors of stroke with severe disability. BMJ Open, 2021, 11, e052593.	0.8	9
14	Discharge after hip fracture surgery in relation to mobilisation timing by patient characteristics: linked secondary analysis of the UK National Hip Fracture Database. BMC Geriatrics, 2021, 21, 694.	1.1	7
15	Exploring the experiences of stroke survivors, informal caregivers and healthcare providers in Sierra Leone: a qualitative study protocol. BMJ Open, 2021, 11, e051276.	0.8	1
16	Title is missing!. , 2021, 18, e1003433.		0
17	Title is missing!. , 2021, 18, e1003433.		O
18	Title is missing!. , 2021, 18, e1003433.		0

#	Article	IF	Citations
19	Title is missing!. , 2021, 18, e1003433.		O
20	Title is missing!. , 2021, 18, e1003433.		0
21	Title is missing!. , 2021, 18, e1003433.		0
22	Long-Term Trends in Stroke Survivors Discharged to Care Homes. Stroke, 2020, 51, 179-185.	1.0	21
23	P105â $\in$ fTherapists acceptability of delivering a psychologically informed gait rehabilitation intervention in early rheumatoid arthritis (GREAT): a qualitative interview study. Rheumatology, 2020, 59, .	0.9	0
24	P125â€fA mixed methods feasibility study of a gait rehabilitation programme for people with early rheumatoid arthritis and foot pain. Rheumatology, 2020, 59, .	0.9	0
25	P221 $\hat{a} \in \mathcal{F}$ Treatment fidelity in the Gait Rehabilitation in Early Rheumatoid Arthritis Trial (GREAT) feasibility study. Rheumatology, 2020, 59, .	0.9	0
26	A multicentre, randomised controlled trial to compare the clinical and cost-effectiveness of Lee Silverman Voice Treatment versus standard NHS Speech and Language Therapy versus control in Parkinson's disease: a study protocol for a randomised controlled trial. Trials, 2020, 21, 436.	0.7	1
27	Recruitment challenges in stroke rehabilitation randomized controlled trials: a qualitative exploration of trialists' perspectives using Framework analysis. Clinical Rehabilitation, 2020, 34, 1122-1133.	1.0	3
28	The effect of rehabilitation interventions on physical function and immobility-related complications in severe stroke: a systematic review. BMJ Open, 2020, 10, e033642.	0.8	27
29	A systematic review of the efficiency of recruitment to stroke rehabilitation randomised controlled trials. Trials, 2020, 21, 68.	0.7	22
30	Protocol for a definitive randomised controlled trial and economic evaluation of a community-based rehabilitation programme following hip fracture: fracture in the elderly multidisciplinary rehabilitation-phase III (FEMuR III). BMJ Open, 2020, 10, e039791.	0.8	0
31	Conceptual Framework for an Episode of Rehabilitative Care After Surgical Repair of Hip Fracture. Physical Therapy, 2019, 99, 276-285.	1.1	12
32	Safety and efficacy of co-careldopa as an add-on therapy to occupational and physical therapy in patients after stroke (DARS): a randomised, double-blind, placebo-controlled trial. Lancet Neurology, The, 2019, 18, 530-538.	4.9	39
33	Clinical and cost effectiveness of memory rehabilitation following traumatic brain injury: a pragmatic cluster randomized controlled trial. Clinical Rehabilitation, 2019, 33, 1171-1184.	1.0	9
34	Rating the intelligibility of dysarthic speech amongst people with Parkinson's Disease: a comparison of trained and untrained listeners. Clinical Linguistics and Phonetics, 2019, 33, 1063-1070.	0.5	12
35	Visual Function Questionnaire as an outcome measure for homonymous hemianopia: subscales and supplementary questions, analysis from the VISION trial. Eye, 2019, 33, 1485-1493.	1.1	9
36	Inequity in rehabilitation interventions after hip fracture: a systematic review. Age and Ageing, 2019, 48, 489-497.	0.7	34

#	Article	IF	Citations
37	Motivating Structured walking Activity in people with Intermittent Claudication (MOSAIC): protocol for a randomised controlled trial of a physiotherapist-led, behavioural change intervention versus usual care in adults with intermittent claudication. BMJ Open, 2019, 9, e030002.	0.8	5
38	Functional and emotional outcomes after transient ischemic attack: A 12-month prospective controlled cohort study. International Journal of Stroke, 2019, 14, 522-529.	2.9	12
39	Technology-enhanced learning in physiotherapy education: Student satisfaction and knowledge acquisition of entry-level students in the United Kingdom. Research in Learning Technology, 2019, 27, .	2.3	9
40	A group memory rehabilitation programme for people with traumatic brain injuries: the ReMemBrIn RCT. Health Technology Assessment, 2019, 23, 1-194.	1.3	11
41	Dopamine Augmented Rehabilitation in Stroke (DARS): a multicentre double-blind, randomised controlled trial of co-careldopa compared with placebo, in addition to routine NHS occupational and physical therapy, delivered early after stroke on functional recovery. Efficacy and Mechanism Evaluation, 2019, 6, 1-138.	0.9	1
42	Prognostic factors of functional outcome after hip fracture surgery: a systematic review. Age and Ageing, 2018, 47, 661-670.	0.7	90
43	A randomized controlled trial to compare two methods of constraint-induced movement therapy to improve functional ability in the affected upper limb in pre-school children with hemiplegic cerebral palsy: CATCH TRIAL. Clinical Rehabilitation, 2018, 32, 909-918.	1.0	21
44	Does physical activity change following hip and knee replacement? Matched case-control study evaluating Physical Activity Scale for the Elderly data from the Osteoarthritis Initiative. Physiotherapy, 2018, 104, 80-90.	0.2	17
45	The effect of rehabilitation interventions on physical function and immobility-related complications in severe strokeâ€"protocol for a systematic review. Systematic Reviews, 2018, 7, 197.	2.5	23
46	Stepping to an Auditory Metronome Improves Weight-Bearing Symmetry in Poststroke Hemiparesis. Journal of Applied Biomechanics, 2018, 34, 469-473.	0.3	1
47	Lee Silverman Voice Treatment versus standard speech and language therapy versus control in Parkinson's disease: a pilot randomised controlled trial (PD COMM pilot). Pilot and Feasibility Studies, 2018, 4, 30.	0.5	27
48	Changes in physical activity following total hip or knee arthroplasty: a matched case-control study from the EPIC-Norfolk cohort. Clinical Rehabilitation, 2017, 31, 1548-1557.	1.0	9
49	A pilot randomized controlled trial comparing effectiveness of prism glasses, visual search training and standard care in hemianopia. Acta Neurologica Scandinavica, 2017, 136, 310-321.	1.0	36
50	Conductive Education as a Method of Stroke Rehabilitation: A Single Blind Randomised Controlled Feasibility Study. Archives of Physical Medicine and Rehabilitation, 2017, 98, e48.	0.5	0
51	Choice of outcome measures for the VISION pilot trial of interventions for hemianopia. Acta Neurologica Scandinavica, 2017, 136, 551-553.	1.0	0
52	Is there a difference in physical activity levels in patients before and up to one year after unilateral total hip replacement? A systematic review and meta-analysis. Clinical Rehabilitation, 2017, 31, 639-650.	1.0	25
53	Home-based Reach-to-Grasp training for people after stroke is feasible: a pilot randomised controlled trial. Clinical Rehabilitation, 2017, 31, 891-903.	1.0	15
54	Stepping to the Beat: Feasibility and Potential Efficacy of a Home-Based Auditory-Cued Step Training Program in Chronic Stroke. Frontiers in Neurology, 2017, 8, 412.	1.1	16

#	Article	IF	Citations
55	The PD COMM trial: a protocol for the process evaluation of a randomised trial assessing the effectiveness of two types of SLT for people with Parkinson's disease. Trials, 2017, 18, 397.	0.7	6
56	Development of an evidence-based complex intervention for community rehabilitation of patients with hip fracture using realist review, survey and focus groups. BMJ Open, 2017, 7, e014362.	0.8	17
57	Developing a multidisciplinary rehabilitation package following hip fracture and testing in a randomised feasibility study: Fracture in the Elderly Multidisciplinary Rehabilitation (FEMuR). Health Technology Assessment, 2017, 21, 1-528.	1.3	18
58	Conductive Education as a Method of Stroke Rehabilitation: A Single Blinded Randomised Controlled Feasibility Study. Stroke Research and Treatment, 2016, 2016, 1-6.	0.5	4
59	Metronome Cueing of Walking Reduces Gait Variability after a Cerebellar Stroke. Frontiers in Neurology, 2016, 7, 84.	1.1	24
60	Fracture in the Elderly Multidisciplinary Rehabilitation (FEMuR): a phase II randomised feasibility study of a multidisciplinary rehabilitation package following hip fracture. BMJ Open, 2016, 6, e012422.	0.8	26
61	UK survey of occupational therapist's and physiotherapist's experiences and attitudes towards hip replacement precautions and equipment. BMC Musculoskeletal Disorders, 2016, 17, 228.	0.8	25
62	Assistive devices, hip precautions, environmental modifications and training to prevent dislocation and improve function after hip arthroplasty. The Cochrane Library, 2016, 2016, CD010815.	1.5	24
63	178 Does Physical Activity Change Following Hip and Knee Replacement? An Analysis of Data from the Osteoarthritis Initiative. Rheumatology, 2016, 55, i134-i134.	0.9	3
64	Community occupational therapy for people with dementia and family carers (COTiD-UK) versus treatment as usual (Valuing Active Life in Dementia [VALID] programme): study protocol for a randomised controlled trial. Trials, 2016, 17, 65.	0.7	31
65	A Randomised Controlled Trial of Treatment for Post-Stroke Homonymous Hemianopia: Screening and Recruitment. Neuro-Ophthalmology, 2016, 40, 1-7.	0.4	8
66	Physiotherapy and Occupational Therapy and Mild to Moderate Parkinson Diseaseâ€"Reply. JAMA Neurology, 2016, 73, 894.	4.5	3
67	Physiotherapy and Occupational Therapy vs No Therapy in Mild to Moderate Parkinson Disease. JAMA Neurology, 2016, 73, 291.	4.5	73
68	A feasibility randomised controlled trial of pre-operative occupational therapy to optimise recovery for patients undergoing primary total hip replacement for osteoarthritis (PROOF-THR). Clinical Rehabilitation, 2016, 30, 156-166.	1.0	12
69	The Hospital Anxiety and Depression Scale: a pilot study to examine its latent structure and the link between psychological state and symptom severity in transient ischaemic attack patients. Psychology, Health and Medicine, 2016, 21, 632-638.	1.3	6
70	An Occupational Therapy intervention for residents with stroke-related disabilities in UK Care Homes (OTCH): cluster randomised controlled trial with economic evaluation. Health Technology Assessment, 2016, 20, 1-138.	1.3	11
71	Clinical effectiveness and cost-effectiveness of physiotherapy and occupational therapy versus no therapy in mild to moderate Parkinson's disease: a large pragmatic randomised controlled trial (PD) Tj ETQq1	1 <b>:03</b> 7843	146ngBT /Ove
72	Improving patients' experience and outcome of total joint replacement: the RESTORE programme. Programme Grants for Applied Research, 2016, 4, 1-508.	0.4	18

#	Article	IF	CITATIONS
73	Fracture in the Elderly Multidisciplinary Rehabilitation (FEMuR): study protocol for a phase II randomised feasibility study of a multidisciplinary rehabilitation package following hip fracture [ISRCTN22464643]. Pilot and Feasibility Studies, 2015, 1, 13.	0.5	10
74	An occupational therapy intervention for residents with stroke living in care homes in the United Kingdom: A content analysis of occupational therapy records from the OTCH trial. British Journal of Occupational Therapy, 2015, 78, 422-430.	0.5	1
75	Subjective Experiences of Speech and Language Therapy in Patients with Parkinson's Disease: A Pilot Study. Rehabilitation Research and Practice, 2015, 2015, 1-11.	0.5	15
76	An occupational therapy intervention for residents with stroke related disabilities in UK care homes (OTCH): cluster randomised controlled trial. BMJ, The, 2015, 350, h468-h468.	3.0	47
77	Effectiveness of land-based physiotherapy exercise following hospital discharge following hip arthroplasty for osteoarthritis: an updated systematic review. Physiotherapy, 2015, 101, 252-265.	0.2	38
78	Effectiveness of physiotherapy exercise following total knee replacement: systematic review and meta-analysis. BMC Musculoskeletal Disorders, 2015, 16, 15.	0.8	178
79	Rehabilitation of Memory following Brain Injury (ReMemBrIn): study protocol for a randomised controlled trial. Trials, 2015, 16, 6.	0.7	9
80	Rehabilitation interventions for foot drop in neuromuscular disease. The Cochrane Library, 2015, , CD003908.	1.5	1
81	Feasibility and Preliminary Efficacy of Visual Cue Training to Improve Adaptability of Walking after Stroke: Multi-Centre, Single-Blind Randomised Control Pilot Trial. PLoS ONE, 2015, 10, e0139261.	1.1	36
82	Occupational Therapy for Care Home Residents with Stroke: What is Routine National Practice?. British Journal of Occupational Therapy, 2014, 77, 265-273.	0.5	4
83	A randomised controlled trial to compare the clinical and cost-effectiveness of prism glasses, visual search training and standard care in patients with hemianopia following stroke: a protocol. BMJ Open, 2014, 4, e005885-e005885.	0.8	11
84	Priority setting partnership to identify the top 10 research priorities for the management of Parkinson's disease. BMJ Open, 2014, 4, e006434.	0.8	120
85	Towards a programme theory for fidelity in the evaluation of complex interventions. Journal of Evaluation in Clinical Practice, 2014, 20, 445-452.	0.9	52
86	PD REHAB RCT OF PHYSIOTHERAPY AND OT IN PARKINSON'S DISEASE. Journal of Neurology, Neurosurgery and Psychiatry, 2014, 85, e4.124-e4.	0.9	0
87	Delayed Recovery of Leg Fatigue Symptoms Following a Maximal Exercise Session in People With Multiple Sclerosis. Neurorehabilitation and Neural Repair, 2014, 28, 139-148.	1.4	13
88	The DARS (Dopamine Augmented Rehabilitation in Stroke) trial: protocol for a randomised controlled trial of Co-careldopa treatment in addition to routine NHS occupational and physical therapy after stroke. Trials, 2014, 15, 316.	0.7	19
89	Physiotherapy for Parkinson's disease: a comparison of techniques. The Cochrane Library, 2014, , CD002815.	1.5	100
90	Lee Silverman voice treatment versus standard NHS speech and language therapy versus control in Parkinson's disease (PD COMM pilot): study protocol for a randomized controlled trial. Trials, 2014, 15, 213.	0.7	20

#	Article	IF	CITATIONS
91	Fatigue, psychological and cognitive impairment following transient ischaemic attack and minor stroke: a systematic review. European Journal of Neurology, 2014, 21, 1258-1267.	1.7	129
92	Attentional focus of feedback for improving performance of reach-to-grasp after stroke: a randomised crossover study. Physiotherapy, 2014, 100, 108-115.	0.2	37
93	Health-related quality of life and supportive care in patients with rare long-term neurological conditions. Quality of Life Research, 2013, 22, 1231-1238.	1.5	50
94	Letter to the Editor. Clinical Orthopaedics and Related Research, 2013, 471, 3706-3707.	0.7	2
95	Visual cue training to improve walking and turning after stroke: a study protocol for a multi-centre, single blind randomised pilot trial. Trials, 2013, 14, 276.	0.7	12
96	A systematic review investigating fatigue, psychological and cognitive impairment following TIA and minor stroke: protocol paper. Systematic Reviews, 2013, 2, 72.	2.5	9
97	Recovery of function following hip resurfacing arthroplasty: a randomized controlled trial comparing an accelerated versus standard physiotherapy rehabilitation programme. Clinical Rehabilitation, 2013, 27, 771-784.	1.0	26
98	Physiotherapy versus placebo or no intervention in Parkinson's disease. The Cochrane Library, 2013, , CD002817.	1.5	235
99	Metronome-Cued Stepping in Place after Hemiparetic Stroke: Comparison of a One- and Two-Tone Beat. ISRN Rehabilitation, 2013, 2013, 1-5.	0.6	20
100	Subjective experiences of transient ischaemic attack: a repertory grid approach. Disability and Rehabilitation, 2013, 35, 2205-2212.	0.9	11
101	Occupational therapy for care home residents with stroke. The Cochrane Library, 2013, , CD010116.	1.5	13
102	Comparison of postdischarge physiotherapy versus usual care following primary total knee arthroplasty for osteoarthritis: an exploratory pilot randomized clinical trial. Clinical Rehabilitation, 2012, 26, 629-641.	1.0	21
103	Speech and language therapy versus placebo or no intervention for speech problems in Parkinson's disease. The Cochrane Library, 2012, 2012, CD002812.	1.5	32
104	Comparison of speech and language therapy techniques for speech problems in Parkinson's disease. The Cochrane Library, 2012, , CD002814.	1.5	36
105	Practical research-based guidance for motor imagery practice in neurorehabilitation. Disability and Rehabilitation, 2012, 34, 2192-2200.	0.9	22
106	Reaching the †hard to reach': strategies to recruit black and minority ethnic service users with rare long-term neurological conditions. International Journal of Social Research Methodology: Theory and Practice, 2012, 15, 485-495.	2.3	11
107	Physiotherapy versus placebo or no intervention in Parkinson's disease., 2012, , CD002817.		72
108	Physiotherapy intervention in Parkinson's disease: systematic review and meta-analysis. BMJ, The, 2012, 345, e5004-e5004.	3.0	250

#	Article	IF	Citations
109	Home-based therapy programmes for upper limb functional recovery following stroke. The Cochrane Library, 2012, , CD006755.	1.5	52
110	Physiotherapy versus placebo or no intervention in Parkinson's disease., 2012,, CD002817.		40
111	Hydration and independence in activities of daily living in people with multiple sclerosis: a pilot investigation. Disability and Rehabilitation, 2011, 33, 1822-1825.	0.9	10
112	Huntington's Disease: Current Epidemiology and Pharmacological Management in UK Primary Care. Neuroepidemiology, 2011, 37, 216-221.	1.1	18
113	VISTA-Rehab: A Resource for Stroke Rehabilitation Trials. International Journal of Stroke, 2010, 5, 447-452.	2.9	8
114	Finding the right outcome measures for care home research. Age and Ageing, 2010, 39, 119-122.	0.7	5
115	Occupational Therapy to Optimise Independence in Parkinson's Disease: The Designing and Recording of a Randomised Controlled Trial Intervention. British Journal of Occupational Therapy, 2010, 73, 178-185.	0.5	11
116	A study of perceived facilitators to physical activity in neurological conditions. International Journal of Therapy and Rehabilitation, 2009, 16, 17-24.	0.1	51
117	Pedometer step counts in individuals with neurological conditions. Clinical Rehabilitation, 2009, 23, 171-175.	1.0	38
118	A Population-Based Study of the Prevalence of Fatigue After Transient Ischemic Attack and Minor Stroke. Stroke, 2009, 40, 757-761.	1.0	138
119	An exploratory cluster randomized controlled trial of group exercise on mobility and depression in care home residents. Clinical Rehabilitation, 2009, 23, 146-154.	1.0	28
120	Effectiveness of physiotherapy exercise following hip arthroplasty for osteoarthritis: a systematic review of clinical trials. BMC Musculoskeletal Disorders, 2009, 10, 98.	0.8	112
121	Use of information feedback and attentional focus of feedback in treating the person with a hemiplegic arm. Physiotherapy Research International, 2009, 14, 77-90.	0.7	91
122	Effects of a physiotherapy and occupational therapy intervention on mobility and activity in care home residents: a cluster randomised controlled trial. BMJ: British Medical Journal, 2009, 339, b3123-b3123.	2.4	51
123	Rehabilitation interventions for foot drop in neuromuscular disease. , 2009, , CD003908.		44
124	Treatment of chronic back pain by sensory discrimination training. A Phase I RCT of a novel device (FairMed) vs. TENS. BMC Musculoskeletal Disorders, 2008, 9, 97.	0.8	44
125	The Prevalence of Joint Contractures, Pressure Sores, Painful Shoulder, Other Pain, Falls, and Depression in the Year After a Severely Disabling Stroke. Stroke, 2008, 39, 3329-3334.	1.0	195
126	A phase II exploratory cluster randomized controlled trial of a group mobility training and staff education intervention to promote urinary continence in UK care homes. Clinical Rehabilitation, 2008, 22, 714-721.	1.0	30

#	ARTICLE	IF	CITATIONS
127	Short-term effects on mobility, activities of daily living and health-related quality of life of a Conductive Education programme for adults with multiple sclerosis, Parkinson's disease and stroke. Clinical Rehabilitation, 2008, 22, 329-337.	1.0	13
128	Occupational therapy for patients with problems in personal activities of daily living after stroke: systematic review of randomised trials. BMJ: British Medical Journal, 2007, 335, 922.	2.4	127
129	Effectiveness of physiotherapy exercise after knee arthroplasty for osteoarthritis: systematic review and meta-analysis of randomised controlled trials. BMJ: British Medical Journal, 2007, 335, 812.	2.4	225
130	Rehabilitation interventions for foot drop in neuromuscular disease., 2007,, CD003908.		17
131	Cluster Randomized Pilot Controlled Trial of an Occupational Therapy Intervention for Residents With Stroke in UK Care Homes. Stroke, 2006, 37, 2336-2341.	1.0	59
132	An investigation into the utility of the Stroke Aphasic Depression Questionnaire (SADQ) in care home settings. Clinical Rehabilitation, 2006, 20, 598-602.	1.0	11
133	Occupational Therapy in Nursing and Residential Care Settings: A Description of a Randomised Controlled Trial Intervention. British Journal of Occupational Therapy, 2004, 67, 104-110.	0.5	23
134	Stroke patients entering nursing home care: a content analysis of discharge letters. Clinical Rehabilitation, 2002, 16, 736-740.	1.0	12
135	Physiotherapy versus placebo or no intervention in Parkinson's disease., 2001,, CD002817.		37
136	Comparison of speech and language therapy techniques for speech problems in Parkinson's disease. , 2001, , CD002814.		17
137	Single blind randomized controlled trial of visual feedback after stroke: Effects on stance symmetry and function. Disability and Rehabilitation, 1997, 19, 536-546.	0.9	134
138	Physiotherapy treatment for stroke patients: A survey of current practice. Physiotherapy Theory and Practice, 1996, 12, 87-96.	0.6	82
139	Falls, sway, and symmetry of weight-bearing after stroke. International Disability Studies, 1991, 13, 1-4.	0.4	131
140	The relationships between weight-bearing asymmetry after stroke, motor function and activities of daily living. Physiotherapy Theory and Practice, 1990, 6, 179-185.	0.6	74