Yiping Chen

List of Publications by Citations

Source: https://exaly.com/author-pdf/6524782/yiping-chen-publications-by-citations.pdf

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

85
papers

3,363
citations

34
h-index

92
ext. papers

3,843
ext. citations

34
h-index

4.68
L-index

#	Paper	IF	Citations
85	Rescue of cleft palate inMsx1-deficient mice by transgenicBmp4reveals a network of BMP and Shh signaling in the regulation of mammalian palatogenesis. <i>Development (Cambridge)</i> , 2002 , 129, 4135-414	6.6	269
84	Rescue of cleft palate in Msx1-deficient mice by transgenic Bmp4 reveals a network of BMP and Shh signaling in the regulation of mammalian palatogenesis. <i>Development (Cambridge)</i> , 2002 , 129, 4135-	-6.6 -46	170
83	Wnt5a regulates directional cell migration and cell proliferation via Ror2-mediated noncanonical pathway in mammalian palate development. <i>Development (Cambridge)</i> , 2008 , 135, 3871-9	6.6	167
82	Shox2 is essential for the differentiation of cardiac pacemaker cells by repressing Nkx2-5. <i>Developmental Biology</i> , 2009 , 327, 376-85	3.1	166
81	Antagonistic signals between BMP4 and FGF8 define the expression of Pitx1 and Pitx2 in mouse tooth-forming anlage. <i>Developmental Biology</i> , 2000 , 217, 323-32	3.1	164
80	Shox2-deficient mice exhibit a rare type of incomplete clefting of the secondary palate. <i>Development (Cambridge)</i> , 2005 , 132, 4397-406	6.6	116
79	The cellular and molecular etiology of the cleft secondary palate in Fgf10 mutant mice. <i>Developmental Biology</i> , 2005 , 277, 102-13	3.1	107
78	Genetic interactions between Pax9 and Msx1 regulate lip development and several stages of tooth morphogenesis. <i>Developmental Biology</i> , 2010 , 340, 438-49	3.1	104
77	Pitx2-microRNA pathway that delimits sinoatrial node development and inhibits predisposition to atrial fibrillation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 9181-6	11.5	84
76	Modulation of BMP signaling by Noggin is required for the maintenance of palatal epithelial integrity during palatogenesis. <i>Developmental Biology</i> , 2010 , 347, 109-21	3.1	80
75	Transgenically ectopic expression of Bmp4 to the Msx1 mutant dental mesenchyme restores downstream gene expression but represses Shh and Bmp2 in the enamel knot of wild type tooth germ. <i>Mechanisms of Development</i> , 2000 , 99, 29-38	1.7	79
74	Shaping limbs by apoptosis>. <i>The Journal of Experimental Zoology</i> , 1998 , 282, 691-702		76
73	Msx1 is required for the induction of Patched by Sonic hedgehog in the mammalian tooth germ. <i>Developmental Dynamics</i> , 1999 , 215, 45-53	2.9	74
72	Epithelial Wnt/Etatenin signaling regulates palatal shelf fusion through regulation of TgfB expression. <i>Developmental Biology</i> , 2011 , 350, 511-9	3.1	66
71	Wnt5a regulates growth, patterning, and odontoblast differentiation of developing mouse tooth. <i>Developmental Dynamics</i> , 2011 , 240, 432-40	2.9	64
70	A common Shox2-Nkx2-5 antagonistic mechanism primes the pacemaker cell fate in the pulmonary vein myocardium and sinoatrial node. <i>Development (Cambridge)</i> , 2015 , 142, 2521-32	6.6	63
69	Hand2 is required in the epithelium for palatogenesis in mice. <i>Developmental Biology</i> , 2009 , 330, 131-41	3.1	59

(2003-2007)

68	Shox2 is required for chondrocyte proliferation and maturation in proximal limb skeleton. <i>Developmental Biology</i> , 2007 , 306, 549-59	3.1	59
67	Bmprla is required in mesenchymal tissue and has limited redundant function with Bmprlb in tooth and palate development. <i>Developmental Biology</i> , 2011 , 349, 451-61	3.1	58
66	Ectopic expression of Nkx2.5 suppresses the formation of the sinoatrial node in mice. <i>Developmental Biology</i> , 2011 , 356, 359-69	3.1	55
65	Shox2-deficiency leads to dysplasia and ankylosis of the temporomandibular joint in mice. <i>Mechanisms of Development</i> , 2008 , 125, 729-42	1.7	54
64	BMP-FGF signaling axis mediates Wnt-induced epidermal stratification in developing mammalian skin. <i>PLoS Genetics</i> , 2014 , 10, e1004687	6	51
63	Msx1/Bmp4 genetic pathway regulates mammalian alveolar bone formation via induction of Dlx5 and Cbfa1. <i>Mechanisms of Development</i> , 2003 , 120, 1469-79	1.7	48
62	Expression survey of genes critical for tooth development in the human embryonic tooth germ. <i>Developmental Dynamics</i> , 2007 , 236, 1307-12	2.9	46
61	Application of lentivirus-mediated RNAi in studying gene function in mammalian tooth development. <i>Developmental Dynamics</i> , 2006 , 235, 1334-44	2.9	45
60	Targeted misexpression of constitutively active BMP receptor-IB causes bifurcation, duplication, and posterior transformation of digit in mouse limb. <i>Developmental Biology</i> , 2000 , 220, 154-67	3.1	44
59	The non-canonical BMP and Wnt/Ecatenin signaling pathways orchestrate early tooth development. <i>Development (Cambridge)</i> , 2015 , 142, 128-39	6.6	42
58	Mice with Tak1 deficiency in neural crest lineage exhibit cleft palate associated with abnormal tongue development. <i>Journal of Biological Chemistry</i> , 2013 , 288, 10440-50	5.4	41
57	Induction of human keratinocytes into enamel-secreting ameloblasts. <i>Developmental Biology</i> , 2010 , 344, 795-9	3.1	38
56	Intra-epithelial requirement of canonical Wnt signaling for tooth morphogenesis. <i>Journal of Biological Chemistry</i> , 2013 , 288, 12080-9	5.4	37
55	Functional redundancy between human SHOX and mouse Shox2 genes in the regulation of sinoatrial node formation and pacemaking function. <i>Journal of Biological Chemistry</i> , 2011 , 286, 17029-3	8 ^{5.4}	37
54	Pten loss induces autocrine FGF signaling to promote skin tumorigenesis. <i>Cell Reports</i> , 2014 , 6, 818-26	10.6	36
53	Enhanced BMP signaling prevents degeneration and leads to endochondral ossification of MeckelS cartilage in mice. <i>Developmental Biology</i> , 2013 , 381, 301-11	3.1	35
52	Tissue interaction is required for glenoid fossa development during temporomandibular joint formation. <i>Developmental Dynamics</i> , 2011 , 240, 2466-73	2.9	35
51	Timing of odontogenic neural crest cell migration and tooth-forming capability in mice. Developmental Dynamics, 2003, 226, 713-8	2.9	34

50	Exploring the effects of gene dosage on mandible shape in mice as a model for studying the genetic basis of natural variation. <i>Development Genes and Evolution</i> , 2013 , 223, 279-87	1.8	30
49	Gsk3Is required in the epithelium for palatal elevation in mice. <i>Developmental Dynamics</i> , 2010 , 239, 3235-46	2.9	30
48	Chick Pcl2 regulates the left-right asymmetry by repressing Shh expression in Hensen's node. <i>Development (Cambridge)</i> , 2004 , 131, 4381-91	6.6	30
47	FGF signaling sustains the odontogenic fate of dental mesenchyme by suppressing Etatenin signaling. <i>Development (Cambridge)</i> , 2013 , 140, 4375-85	6.6	28
46	Generation of Shox2-Cre allele for tissue specific manipulation of genes in the developing heart, palate, and limb. <i>Genesis</i> , 2013 , 51, 515-22	1.9	27
45	Augmented BMPRIA-mediated BMP signaling in cranial neural crest lineage leads to cleft palate formation and delayed tooth differentiation. <i>PLoS ONE</i> , 2013 , 8, e66107	3.7	27
44	BMPRIA mediated signaling is essential for temporomandibular joint development in mice. <i>PLoS ONE</i> , 2014 , 9, e101000	3.7	26
43	Directed Bmp4 expression in neural crest cells generates a genetic model for the rare human bony syngnathia birth defect. <i>Developmental Biology</i> , 2014 , 391, 170-81	3.1	25
42	Mice with an anterior cleft of the palate survive neonatal lethality. <i>Developmental Dynamics</i> , 2008 , 237, 1509-16	2.9	24
41	An atypical canonical bone morphogenetic protein (BMP) signaling pathway regulates Msh homeobox 1 (Msx1) expression during odontogenesis. <i>Journal of Biological Chemistry</i> , 2014 , 289, 31492	2-502	23
40	The role of Shox2 in SAN development and function. <i>Pediatric Cardiology</i> , 2012 , 33, 882-9	2.1	23
39	FGF8 signaling sustains progenitor status and multipotency of cranial neural crest-derived mesenchymal cells in vivo and in vitro. <i>Journal of Molecular Cell Biology</i> , 2015 , 7, 441-54	6.3	22
38	Expression of SHH signaling molecules in the developing human primary dentition. <i>BMC Developmental Biology</i> , 2013 , 13, 11	3.1	22
38 37	, , , , , , , , , , , , , , , , , , , ,	3.1 5.4	22
	Developmental Biology, 2013, 13, 11 The short stature homeobox 2 (Shox2)-bone morphogenetic protein (BMP) pathway regulates dorsal mesenchymal protrusion development and its temporary function as a pacemaker during		
37	Developmental Biology, 2013, 13, 11 The short stature homeobox 2 (Shox2)-bone morphogenetic protein (BMP) pathway regulates dorsal mesenchymal protrusion development and its temporary function as a pacemaker during cardiogenesis. Journal of Biological Chemistry, 2015, 290, 2007-23 Genetic Regulation of Sinoatrial Node Development and Pacemaker Program in the Venous Pole.	5.4	21
37	Developmental Biology, 2013, 13, 11 The short stature homeobox 2 (Shox2)-bone morphogenetic protein (BMP) pathway regulates dorsal mesenchymal protrusion development and its temporary function as a pacemaker during cardiogenesis. Journal of Biological Chemistry, 2015, 290, 2007-23 Genetic Regulation of Sinoatrial Node Development and Pacemaker Program in the Venous Pole. Journal of Cardiovascular Development and Disease, 2015, 2, 282-298 Evidence for the differential regulation of Nkx-6.1 expression in the ventral spinal cord and foregut	5.4	21 18

(2014-2014)

32	Replacing Shox2 with human SHOX leads to congenital disc degeneration of the temporomandibular joint in mice. <i>Cell and Tissue Research</i> , 2014 , 355, 345-54	4.2	15
31	Expression and regulation of the chicken Nkx-6.2 homeobox gene suggest its possible involvement in the ventral neural patterning and cell fate specification. <i>Developmental Dynamics</i> , 1999 , 216, 459-68	2.9	15
30	Conditional deletion of Bmp2 in cranial neural crest cells recapitulates Pierre Robin sequence in mice. <i>Cell and Tissue Research</i> , 2019 , 376, 199-210	4.2	15
29	ISLET1-Dependent ECatenin/Hedgehog Signaling Is Required for Outgrowth of the Lower Jaw. <i>Molecular and Cellular Biology</i> , 2017 , 37,	4.8	12
28	defines a subpopulation of pacemaker cells and is essential for the physiological function of the sinoatrial node in mice. <i>Development (Cambridge)</i> , 2019 , 146,	6.6	12
27	Expression patterns of genes critical for BMP signaling pathway in developing human primary tooth germs. <i>Histochemistry and Cell Biology</i> , 2014 , 142, 657-65	2.4	12
26	Exogenous fibroblast growth factor 8 rescues development of mouse diastemal vestigial tooth ex vivo. <i>Developmental Dynamics</i> , 2011 , 240, 1344-53	2.9	11
25	Cellular and developmental basis of orofacial clefts. <i>Birth Defects Research</i> , 2020 , 112, 1558-1587	2.9	11
24	Regrowing a tooth: in vitro and in vivo approaches. Current Opinion in Cell Biology, 2019, 61, 126-131	9	10
23	A unique stylopod patterning mechanism by Shox2-controlled osteogenesis. <i>Development</i> (Cambridge), 2016 , 143, 2548-60	6.6	10
22	Bioengineering of a human whole tooth: progress and challenge. Cell Regeneration, 2014, 3, 8	2.5	10
21	Mouse embryonic diastema region is an ideal site for the development of ectopically transplanted tooth germ. <i>Developmental Dynamics</i> , 2008 , 237, 411-6	2.9	10
20	Efficient induction of functional ameloblasts from human keratinocyte stem cells. <i>Stem Cell Research and Therapy</i> , 2018 , 9, 126	8.3	9
19	Precise chronology of differentiation of developing human primary dentition. <i>Histochemistry and Cell Biology</i> , 2014 , 141, 221-7	2.4	9
18	Opposing roles of TCF7/LEF1 and TCF7L2 in cyclin D2 and Bmp4 expression and cardiomyocyte cell cycle control during late heart development. <i>Laboratory Investigation</i> , 2019 , 99, 807-818	5.9	9
17	LDL Receptor-Related Protein 6 Modulates Ret Proto-Oncogene Signaling in Renal Development and Cystic Dysplasia. <i>Journal of the American Society of Nephrology: JASN</i> , 2016 , 27, 417-27	12.7	8
16	Shox2 regulates osteogenic differentiation and pattern formation during hard palate development in mice. <i>Journal of Biological Chemistry</i> , 2019 , 294, 18294-18305	5.4	7
15	Phosphorylation of Shox2 is required for its function to control sinoatrial node formation. <i>Journal of the American Heart Association</i> , 2014 , 3, e000796	6	7

14	Overexpression of constitutively active BMP-receptor-IB in mouse skin causes an ichthyosis-vulgaris-like disease. <i>Cell and Tissue Research</i> , 2010 , 342, 401-10	4.2	7
13	Augmented Indian hedgehog signaling in cranial neural crest cells leads to craniofacial abnormalities and dysplastic temporomandibular joint in mice. <i>Cell and Tissue Research</i> , 2016 , 364, 105-	1 ⁴ 5 ⁻²	6
12	Temporomandibular Joint Development 2013 , 71-85		5
11	Persistent Noggin arrests cardiomyocyte morphogenesis and results in early in utero lethality. <i>Developmental Dynamics</i> , 2015 , 244, 457-67	2.9	4
10	Single-cell RNA sequencing deconvolutes the heterogeneity of human bone marrow-derived mesenchymal stem cells. <i>International Journal of Biological Sciences</i> , 2021 , 17, 4192-4206	11.2	4
9	The transcriptional regulator MEIS2 sets up the ground state for palatal osteogenesis in mice. Journal of Biological Chemistry, 2020 , 295, 5449-5460	5.4	3
8	Olig2 regulates terminal differentiation and maturation of peripheral olfactory sensory neurons. <i>Cellular and Molecular Life Sciences</i> , 2020 , 77, 3597-3609	10.3	3
7	Exogenous FGF8 signaling in osteocytes leads to mandibular hypoplasia in mice. <i>Oral Diseases</i> , 2020 , 26, 590-596	3.5	2
6	A systematic dissection of human primary osteoblasts at single-cell resolution. <i>Aging</i> , 2021 , 13, 20629-2	20,6650	2
5	Identification and analysis of a novel bmp4 enhancer in Fugu genome. <i>Archives of Oral Biology</i> , 2015 , 60, 540-5	2.8	1
4	Conjugated activation of myocardial-specific transcription of Gja5 by a pair of Nkx2-5-Shox2 co-responsive elements. <i>Developmental Biology</i> , 2020 , 465, 79-87	3.1	1
3	The Transcription Factor Shox2 Shapes Neuron Firing Properties and Suppresses Seizures by Regulation of Key Ion Channels in Thalamocortical Neurons. <i>Cerebral Cortex</i> , 2021 , 31, 3194-3212	5.1	1
2	Reply to Kelder et al.: Does the dorsal mesenchymal protrusion act as a temporary pacemaker during heart development?. <i>Journal of Biological Chemistry</i> , 2015 , 290, 8015	5.4	
1	Wnt5a regulates directional cell migration and cell proliferation via Ror2-mediated noncanonical pathway in mammalian palatogenesis. <i>FASEB Journal</i> , 2009 , 23, 308.4	0.9	