

Lieh-Bang Liou

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6531851/publications.pdf>

Version: 2024-02-01

13
papers

258
citations

1306789

7
h-index

1125271

13
g-index

13
all docs

13
docs citations

13
times ranked

343
citing authors

#	ARTICLE	IF	CITATIONS
1	$\hat{I}\pm$ -2,6-sialic acid/IgG anti-dsDNA ratios correlate with human lupus disease activity and possible mechanisms: A pilot study. <i>Lupus</i> , 2022, 31, 927-938.	0.8	2
2	De-sialylated and sialylated IgG anti-dsDNA antibodies respectively worsen and mitigate experimental mouse lupus proteinuria and possible mechanisms. <i>International Immunopharmacology</i> , 2022, 109, 108837.	1.7	2
3	Reverse expression of $\hat{I}\pm$ 2,6-sialic acid ratios on IgG, IgM, and IgG/IgM autoantibodies correlates with mouse arthritis and rheumatoid arthritis disease activity. <i>Journal of the Chinese Medical Association</i> , 2020, 83, 1079-1086.	0.6	8
4	A new laboratory surrogate (Monocyte Chemotactic Protein-1) for Disease Activity Score ₂₈ : a favourable indicator for remission in rheumatoid arthritis. <i>Scientific Reports</i> , 2020, 10, 8238.	1.6	5
5	Septicaemia is associated with increased disease activity and mortality in systemic lupus erythematosus: a retrospective analysis from Taiwan. <i>Lupus</i> , 2020, 29, 191-198.	0.8	4
6	$\hat{I}\pm$ -2,3-Sialyltransferase 1 and neuraminidase-3 from monocytes in patients with rheumatoid arthritis correlate with disease activity measures. <i>Journal of the Chinese Medical Association</i> , 2019, 82, 179-185.	0.6	7
7	Sialyltransferase and Neuraminidase Levels/Ratios and Sialic Acid Levels in Peripheral Blood B Cells Correlate with Measures of Disease Activity in Patients with Systemic Lupus Erythematosus and Rheumatoid Arthritis: A Pilot Study. <i>PLoS ONE</i> , 2016, 11, e0151669.	1.1	12
8	Blood Monocyte Chemotactic Protein-1 (MCP-1) and Adapted Disease Activity Score ₂₈ -MCP-1: Favorable Indicators for Rheumatoid Arthritis Activity. <i>PLoS ONE</i> , 2013, 8, e55346.	1.1	33
9	Cytomegalovirus infection causes morbidity and mortality in patients with autoimmune diseases, particularly systemic lupus: in a Chinese population in Taiwan. <i>Rheumatology International</i> , 2012, 32, 2901-2908.	1.5	43
10	Invasive fungal infection in systemic lupus erythematosus: an analysis of 15 cases and a literature review. <i>Rheumatology</i> , 2007, 46, 539-544.	0.9	95
11	Different monocyte reaction patterns in newly diagnosed, untreated rheumatoid arthritis and lupus patients probably confer disparate C-reactive protein levels. <i>Clinical and Experimental Rheumatology</i> , 2003, 21, 437-44.	0.4	12
12	Serum and in vitro production of IL-1 receptor antagonist correlate with C-reactive protein levels in newly diagnosed, untreated lupus patients. <i>Clinical and Experimental Rheumatology</i> , 2001, 19, 515-23.	0.4	34
13	Lymphadenopathy in an oriental with ankylosing spondylitis. <i>Clinical Rheumatology</i> , 1998, 17, 412-415.	1.0	1