

Vibeke Bertelsen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6537028/publications.pdf>

Version: 2024-02-01

14
papers

630
citations

840776

11
h-index

1058476

14
g-index

14
all docs

14
docs citations

14
times ranked

1050
citing authors

#	ARTICLE	IF	CITATIONS
1	Protein kinase C regulates ErbB3 turnover. <i>Experimental Cell Research</i> , 2019, 382, 111473.	2.6	7
2	Hypo-osmotic Stress Drives IL-33 Production in Human Keratinocytes An Epidermal Homeostatic Response. <i>Journal of Investigative Dermatology</i> , 2019, 139, 81-90.	0.7	10
3	ErbB3 interacts with Hrs and is sorted to lysosomes for degradation. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2017, 1864, 2241-2252.	4.1	8
4	A combination of two antibodies recognizing non-overlapping epitopes of HER2 induces kinase activity-dependent internalization of HER2. <i>Journal of Cellular and Molecular Medicine</i> , 2016, 20, 1999-2010.	3.6	21
5	Interaction with epsin 1 regulates the constitutive clathrin-dependent internalization of ErbB3. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2016, 1863, 1179-1188.	4.1	17
6	Epidermal Expression and Regulation of Interleukin-33 during Homeostasis and Inflammation: Strong Species Differences. <i>Journal of Investigative Dermatology</i> , 2015, 135, 1771-1780.	0.7	53
7	The Mysterious Ways of ErbB2/HER2 Trafficking. <i>Membranes</i> , 2014, 4, 424-446.	3.0	145
8	Preubiquitinated chimeric ErbB2 is constitutively endocytosed and subsequently degraded in lysosomes. <i>Experimental Cell Research</i> , 2013, 319, 32-45.	2.6	16
9	Pertuzumab counteracts the inhibitory effect of ErbB2 on degradation of ErbB3. <i>Carcinogenesis</i> , 2013, 34, 2031-2038.	2.8	22
10	The oncoprotein ErbB3 is endocytosed in the absence of added ligand in a clathrin-dependent manner. <i>Carcinogenesis</i> , 2012, 33, 1031-1039.	2.8	27
11	A Chimeric Preubiquitinated EGF Receptor is Constitutively Endocytosed in a Clathrin-Dependent, but Kinase-Independent Manner. <i>Traffic</i> , 2011, 12, 507-520.	2.7	37
12	Epsin 1 is Involved in Recruitment of Ubiquitinated EGF Receptors into Clathrin-Coated Pits. <i>Traffic</i> , 2009, 10, 235-245.	2.7	95
13	The Cbl-interacting protein TULA inhibits dynamin-dependent endocytosis. <i>Experimental Cell Research</i> , 2007, 313, 1696-1709.	2.6	27
14	Cbl-dependent Ubiquitination Is Required for Progression of EGF Receptors into Clathrin-coated Pits. <i>Molecular Biology of the Cell</i> , 2004, 15, 3591-3604.	2.1	145