List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6541665/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Analgesic use in adolescents with patellofemoral pain or Osgood–Schlatter Disease: a secondary cross-sectional analysis of 323 subjects. Scandinavian Journal of Pain, 2022, 22, 543-551.	0.5	5
2	Gold microâ€particles for knee osteoarthritis. European Journal of Pain, 2022, 26, 811-824.	1.4	8
3	The Role of Nuclear Medicine Imaging with 18F-FDG PET/CT, Combined 111In-WBC/99mTc-Nanocoll, and 99mTc-HDP SPECT/CT in the Evaluation of Patients with Chronic Problems after TKA or THA in a Prospective Study. Diagnostics, 2022, 12, 681.	1.3	5
4	Interactions between running volume and running pace on injury occurrence in recreational runners: A secondary analysis Journal of Athletic Training, 2021, , .	0.9	0
5	Biomechanical Evaluation of the Effect of Minimally Invasive Spine Surgery Compared with Traditional Approaches in Lifting Tasks. Frontiers in Bioengineering and Biotechnology, 2021, 9, 724854.	2.0	Ο
6	Influence of Ibuprofen on Bone Healing After Colles' Fracture: A Randomized Controlled Clinical Trial. Journal of Orthopaedic Research, 2020, 38, 545-554.	1.2	12
7	Cost-effectiveness of total knee replacement in addition to non-surgical treatment: a 2-year outcome from a randomised trial in secondary care in Denmark. BMJ Open, 2020, 10, e033495.	0.8	14
8	Support for mobilising medical students to join the COVID-19 pandemic emergency healthcare workforce: a cross-sectional questionnaire survey. BMJ Open, 2020, 10, e039082.	0.8	27
9	Multidisciplinary Diagnostic Algorithm for Evaluation of Patients Presenting with a Prosthetic Problem in the Hip or Knee: A Prospective Study. Diagnostics, 2020, 10, 98.	1.3	5
10	Cost-effectiveness of 12 weeks of supervised treatment compared to written advice in patients with knee osteoarthritis: a secondary analysis of the 2-year outcome from a randomized trial. Osteoarthritis and Cartilage, 2020, 28, 907-916.	0.6	12
11	Medical students for health-care staff shortages during the COVID-19 pandemic. Lancet, The, 2020, 395, e79-e80.	6.3	130
12	Neuromuscular Exercises Improve Shoulder Function More Than Standard Care Exercises in Patients With a Traumatic Anterior Shoulder Dislocation: A Randomized Controlled Trial. Orthopaedic Journal of Sports Medicine, 2020, 8, 232596711989610.	0.8	21
13	No influence of ibuprofen on bone healing after Colles' fracture – A randomized controlled clinical trial. Injury, 2019, 50, 1309-1317.	0.7	4
14	Patients with non-operated traumatic primary or recurrent anterior shoulder dislocation have equally poor self-reported and measured shoulder function: a cross-sectional study. BMC Musculoskeletal Disorders, 2019, 20, 59.	0.8	15
15	The Garmin-RUNSAFE Running Health Study on the aetiology of running-related injuries: rationale and design of an 18-month prospective cohort study including runners worldwide. BMJ Open, 2019, 9, e032627.	0.8	9
16	72 revision surgeries for aseptic failure after hip or knee arthroplasty: a prospective study with an extended diagnostic algorithm. BMC Musculoskeletal Disorders, 2019, 20, 600.	0.8	4
17	Response to Letter to Editor: â€~2-year outcome from two parallel randomized controlled trials. Reporting considerations'. Osteoarthritis and Cartilage, 2019, 27, e1-e2.	0.6	0
18	Increased postural stiffness during challenging postural tasks in patients with knee osteoarthritis with high pain sensitization. Clinical Biomechanics, 2019, 61, 129-135.	0.5	8

#	Article	IF	CITATIONS
19	Prediction of running-induced Achilles tendinopathy with pain sensitivity – a 1-year prospective study. Scandinavian Journal of Pain, 2019, 19, 139-146.	0.5	7
20	Changes in the running-related injury incidence rate ratio in a 1000-km explorative prospective cohort study involving two unspecific shoe changes. Footwear Science, 2019, 11, 63-70.	0.8	3
21	Contention over undergraduate medical curriculum content. International Journal of Medical Education, 2019, 10, 230-231.	0.6	6
22	Run Clever – No difference in risk of injury when comparing progression in running volume and running intensity in recreational runners: A randomised trial. BMJ Open Sport and Exercise Medicine, 2018, 4, e000333.	1.4	19
23	Differential Contributions of Specimen Types, Culturing, and 16S rRNA Sequencing in Diagnosis of Prosthetic Joint Infections. Journal of Clinical Microbiology, 2018, 56, .	1.8	22
24	NSAIDs are superior to paracetamol for osteoarthritic pain and function in a network meta-analysis. BMJ Evidence-Based Medicine, 2018, 23, 40-41.	1.7	6
25	Pain and sensitization after total knee replacement or nonsurgical treatment in patients with knee osteoarthritis: Identifying potential predictors of outcome at 12Âmonths. European Journal of Pain, 2018, 22, 1088-1102.	1.4	38
26	Total knee replacement and non-surgical treatment of knee osteoarthritis: 2-year outcome from two parallel randomized controlled trials. Osteoarthritis and Cartilage, 2018, 26, 1170-1180.	0.6	106
27	The association between eccentric hip abduction strength and hip and knee angular movements in recreational male runners: An explorative study. Scandinavian Journal of Medicine and Science in Sports, 2018, 28, 473-478.	1.3	11
28	How Do Novice Runners With Different Body Mass Indexes Begin a Self-chosen Running Regime?. Journal of Orthopaedic and Sports Physical Therapy, 2018, 48, 873-877.	1.7	3
29	Progression in Running Intensity or Running Volume and the Development of Specific Injuries in Recreational Runners: Run Clever, a Randomized Trial Using Competing Risks. Journal of Orthopaedic and Sports Physical Therapy, 2018, 48, 740-748.	1.7	21
30	THE START-TO-RUN DISTANCE AND RUNNING-RELATED INJURY AMONG OBESE NOVICE RUNNERS: A RANDOMIZED TRIAL. International Journal of Sports Physical Therapy, 2018, 13, 943-955.	0.5	10
31	Association Between the ABO Blood Types and Post-operative Pain. European Neurological Review, 2018, 13, 38.	0.5	5
32	THE START-TO-RUN DISTANCE AND RUNNING-RELATED INJURY AMONG OBESE NOVICE RUNNERS: A RANDOMIZED TRIAL. International Journal of Sports Physical Therapy, 2018, 13, 943-955.	0.5	1
33	The long-term outcome after treatment for patients with tibial fracture treated with intramedullary nailing is not influenced by time of day of surgery and surgeon experience. European Journal of Trauma and Emergency Surgery, 2017, 43, 221-226.	0.8	6
34	THE IMPACT OF RUNNING INJURIES IN RELATION TO OTHER REASONS FOR DISCONTINUED RUNNING IN NOVICE RUNNERS. British Journal of Sports Medicine, 2017, 51, 296.2-296.	3.1	2
35	A neuromuscular exercise programme versus standard care for patients with traumatic anterior shoulder instability: study protocol for a randomised controlled trial (the SINEX study). Trials, 2017, 18, 90.	0.7	12
36	Medial shoe-ground pressure and specific running injuries: A 1-year prospective cohort study. Journal of Science and Medicine in Sport, 2017, 20, 830-834.	0.6	27

#	Article	IF	CITATIONS
37	Variation in Prosthetic Joint Infection and treatment strategies during 4.5Âyears of follow-up after primary joint arthroplasty using administrative data of 41397 patients across Australian, European and United States hospitals. BMC Musculoskeletal Disorders, 2017, 18, 207.	0.8	31
38	Asymmetry in gait pattern following tibial shaft fractures – a prospective one-year follow-up study of 49 patients. Gait and Posture, 2017, 51, 47-51.	0.6	13
39	Effects of Chronic Musculoskeletal Pain on Fertility Potential in Lean and Overweight Male Patients. Pain Research and Management, 2017, 2017, 1-10.	0.7	0
40	Local and Widespread Hyperalgesia After Isolated Tibial Shaft Fractures Treated with Intramedullary Nailing. Pain Medicine, 2016, 17, pnv016.	0.9	3
41	The effects of total knee replacement and nonâ€surgical treatment on pain sensitization and clinical pain. European Journal of Pain, 2016, 20, 1612-1621.	1.4	23
42	A Pharmacokinetic-Pharmacodynamic Model of Morphine Exposure and Subsequent Morphine Consumption in Postoperative Pain. Pharmaceutical Research, 2016, 33, 1093-1103.	1.7	13
43	Enhanced function and quality of life following 5Âmonths of exercise therapy for patients with irreparable rotator cuff tears – an intervention study. BMC Musculoskeletal Disorders, 2016, 17, 252.	0.8	27
44	One-year development of QOL following orthopaedic polytrauma: a prospective observational cohort study of 53 patients. Archives of Orthopaedic and Trauma Surgery, 2016, 136, 1539-1546.	1.3	15
45	Decreased QOL and muscle strength are persistent 1Âyear after intramedullary nailing of a tibial shaft fracture: a prospective 1-year follow-up cohort study. Archives of Orthopaedic and Trauma Surgery, 2016, 136, 1395-1402.	1.3	12
46	Self-reported Recovery is Associated With Improvement in Localized Hyperalgesia Among Adolescent Females With Patellofemoral Pain. Clinical Journal of Pain, 2016, 32, 428-434.	0.8	15
47	The design of the run Clever randomized trial: running volume, â^'intensity and running-related injuries. BMC Musculoskeletal Disorders, 2016, 17, 177.	0.8	12
48	ls Knee Pain During Adolescence a Self-limiting Condition?. American Journal of Sports Medicine, 2016, 44, 1165-1171.	1.9	157
49	Effect of exercise therapy on neuromuscular activity and knee strength in female adolescents with patellofemoral pain—An ancillary analysis of a cluster randomized trial. Clinical Biomechanics, 2016, 34, 22-29.	0.5	17
50	Criteria used when deciding on eligibility for total knee arthroplasty — Between thinking and doing. Knee, 2016, 23, 300-305.	0.8	29
51	The efficacy of non-surgical treatment on pain and sensitization in patients with knee osteoarthritis: a pre-defined ancillary analysis from a randomized controlled trial. Osteoarthritis and Cartilage, 2016, 24, 108-116.	0.6	21
52	The outcome after lateral tibial plateau fracture treated with percutaneus screw fixation show a tendency towards worse functional outcome compared with a reference population. European Journal of Trauma and Emergency Surgery, 2016, 42, 177-184.	0.8	14
53	High degree of patient satisfaction after percutaneous treatment of lateral tibia plateau fractures. Danish Medical Journal, 2016, 63, A5174.	0.5	5
54	Population-Based Epidemiology of Tibial Plateau Fractures. Orthopedics, 2015, 38, e780-6.	0.5	158

#	Article	IF	CITATIONS
55	Repeated Time-to-event Analysis of Consecutive Analgesic Events in Postoperative Pain. Anesthesiology, 2015, 123, 1411-1419.	1.3	28
56	Tourniquet induced ischemia and changes in metabolism during TKA: a randomized study using microdialysis. BMC Musculoskeletal Disorders, 2015, 16, 326.	0.8	32
57	Examination of Muscle Strength and Pressure Pain Thresholds in Knee Osteoarthritis. Journal of Geriatric Physical Therapy, 2015, 38, 141-147.	0.6	27
58	High Eccentric Hip Abduction Strength Reduces the Risk of Developing Patellofemoral Pain Among Novice Runners Initiating a Self-Structured Running Program: A 1-Year Observational Study. Journal of Orthopaedic and Sports Physical Therapy, 2015, 45, 153-161.	1.7	36
59	The efficacy of 12 weeks non-surgical treatment for patients not eligible for total knee replacement: a randomized controlled trial with 1-year follow-up. Osteoarthritis and Cartilage, 2015, 23, 1465-1475.	0.6	70
60	Decreased muscle strength is associated with impaired long-term functional outcome after intramedullary nailing of femoral shaft fracture. European Journal of Trauma and Emergency Surgery, 2015, 41, 673-681.	0.8	14
61	Progressive high-load strength training compared with general low-load exercises in patients with rotator cuff tendinopathy: study protocol for a randomised controlled trial. Trials, 2015, 16, 27.	0.7	19
62	Incidence of Running-Related Injuries Per 1000Âh of running in Different Types of Runners: A Systematic Review and Meta-Analysis. Sports Medicine, 2015, 45, 1017-1026.	3.1	283
63	Incidence and epidemiology of tibial shaft fractures. Injury, 2015, 46, 746-750.	0.7	161
64	Chronic Postoperative Pain After Primary and Revision Total Knee Arthroplasty. Clinical Journal of Pain, 2015, 31, 1-6.	0.8	94
65	A Randomized, Controlled Trial of Total Knee Replacement. New England Journal of Medicine, 2015, 373, 1597-1606.	13.9	498
66	Knee Confidence as It Relates to Self-reported and Objective Correlates of Knee Osteoarthritis: A Cross-sectional Study of 220 Patients. Journal of Orthopaedic and Sports Physical Therapy, 2015, 45, 765-771.	1.7	13
67	Absence of a Tourniquet Does Not Affect Fixation of Cemented TKA: A Randomized RSA Study of 70 Patients. Journal of Arthroplasty, 2015, 30, 2128-2132.	1.5	43
68	Exercise during school hours when added to patient education improves outcome for 2â€years in adolescent patellofemoral pain: a cluster randomised trial. British Journal of Sports Medicine, 2015, 49, 406-412.	3.1	113
69	Successful conservative treatment of patients with MRI-verified meniscal lesions. Knee Surgery, Sports Traumatology, Arthroscopy, 2015, 23, 178-183.	2.3	11
70	A Prospective Study on Time to Recovery in 254 Injured Novice Runners. PLoS ONE, 2014, 9, e99877.	1.1	80
71	Blockade of Glutamate Release by Botulinum Neurotoxin Type A in Humans: A Dermal Microdialysis Study. Pain Research and Management, 2014, 19, 126-132.	0.7	22
72	Facilitation of pain sensitization in knee osteoarthritis and persistent postâ€operative pain: A crossâ€sectional study. European Journal of Pain, 2014, 18, 1024-1031.	1.4	61

#	Article	IF	CITATIONS
73	The orthopaedic research scene and strategies to improve it. Bone and Joint Journal, 2014, 96-B, 1578-1585.	1.9	6
74	Restrictions in Quality of Life After Intramedullary Nailing of Tibial Shaft Fracture. Journal of Orthopaedic Trauma, 2014, 28, 507-512.	0.7	30
75	Foot pronation is not associated with increased injury risk in novice runners wearing a neutral shoe: a 1-year prospective cohort study. British Journal of Sports Medicine, 2014, 48, 440-447.	3.1	93
76	Comparisons of increases in knee and ankle joint moments following an increase in running speed from 8 to 12 to 16km·hâ^'1. Clinical Biomechanics, 2014, 29, 959-964.	0.5	31
77	â€~All in a box' a concept for optimizing microbiological diagnostic sampling in prosthetic joint infections. BMC Research Notes, 2014, 7, 418.	0.6	9
78	Excessive Progression in Weekly Running Distance and Risk of Running-Related Injuries: An Association Which Varies According to Type of Injury. Journal of Orthopaedic and Sports Physical Therapy, 2014, 44, 739-747.	1.7	114
79	Faster recovery without the use of a tourniquet in total knee arthroplasty. Monthly Notices of the Royal Astronomical Society: Letters, 2014, 85, 422-426.	1.2	130
80	Impaired postural balance in the morning in patients with knee osteoarthritis. Gait and Posture, 2014, 39, 1040-1044.	0.6	20
81	Normative values of eccentric hip abduction strength in novice runners: an equation adjusting for age and gender. International Journal of Sports Physical Therapy, 2014, 9, 68-75.	0.5	9
82	Running more than three kilometers during the first week of a running regimen may be associated with increased risk of injury in obese novice runners. International Journal of Sports Physical Therapy, 2014, 9, 338-45.	0.5	19
83	Care-seeking behaviour of adolescents with knee pain: a population-based study among 504 adolescents. BMC Musculoskeletal Disorders, 2013, 14, 225.	0.8	50
84	High prevalence of daily and multi-site pain – a cross-sectional population-based study among 3000 Danish adolescents. BMC Pediatrics, 2013, 13, 191.	0.7	106
85	Widespread sensitization in patients with chronic pain after revision total knee arthroplasty. Pain, 2013, 154, 1588-1594.	2.0	121
86	Footstrike patterns among novice runners wearing a conventional, neutral running shoe. Gait and Posture, 2013, 38, 354-356.	0.6	47
87	Predictors of Running-Related Injuries Among 930 Novice Runners. Orthopaedic Journal of Sports Medicine, 2013, 1, 232596711348731.	0.8	67
88	Degenerative Spondylolisthesis Is Associated with Low Spinal Bone Density: A Comparative Study between Spinal Stenosis and Degenerative Spondylolisthesis. BioMed Research International, 2013, 2013, 1-8.	0.9	18
89	Neuromuscular Activity and Knee Kinematics in Adolescents with Patellofemoral Pain. Medicine and Science in Sports and Exercise, 2013, 45, 1730-1739.	0.2	43
90	Lower Mechanical Pressure Pain Thresholds in Female Adolescents With Patellofemoral Pain Syndrome. Journal of Orthopaedic and Sports Physical Therapy, 2013, 43, 414-421.	1.7	92

#	Article	lF	CITATIONS
91	Can GPS Be Used to Detect Deleterious Progression in Training Volume Among Runners?. Journal of Strength and Conditioning Research, 2013, 27, 1471-1478.	1.0	56
92	Hip and Knee Strength Is Not Affected in 12-16 Year Old Adolescents with Patellofemoral Pain - A Cross-Sectional Population-Based Study. PLoS ONE, 2013, 8, e79153.	1.1	58
93	Weekly running volume and risk of running-related injuries among marathon runners. International Journal of Sports Physical Therapy, 2013, 8, 111-20.	0.5	41
94	Classifying running-related injuries based upon etiology, with emphasis on volume and pace. International Journal of Sports Physical Therapy, 2013, 8, 172-9.	0.5	28
95	Half of 12-15-year-olds with knee pain still have pain after one year. Danish Medical Journal, 2013, 60, A4725.	0.5	30
96	Efficacy of multimodal, systematic non-surgical treatment of knee osteoarthritis for patients not eligible for a total knee replacement: a study protocol of a randomised controlled trial. BMJ Open, 2012, 2, e002168.	0.8	23
97	Total knee replacement plus physical and medical therapy or treatment with physical and medical therapy alone: a randomised controlled trial in patients with knee osteoarthritis (the MEDIC-study). BMC Musculoskeletal Disorders, 2012, 13, 67.	0.8	18
98	Autologous Blood Transfusion after Local Infiltration Analgesia with Ropivacaine in Total Knee and Hip Arthroplasty. Anesthesiology Research and Practice, 2012, 2012, 1-6.	0.2	2
99	Early intervention for adolescents with Patellofemoral Pain Syndrome - a pragmatic cluster randomised controlled trial. BMC Musculoskeletal Disorders, 2012, 13, 9.	0.8	34
100	Training errors and running related injuries: a systematic review. International Journal of Sports Physical Therapy, 2012, 7, 58-75.	0.5	115
101	Fusion mass bone quality after uninstrumented spinal fusion in older patients. European Spine Journal, 2010, 19, 2200-2208.	1.0	33
102	Microdialysis of paraspinal muscle in healthy volunteers and patients underwent posterior lumbar fusion surgery. European Spine Journal, 2009, 18, 1604-1609.	1.0	10
103	The Effect of Electrical Stimulation on Lumbar Spinal Fusion in Older Patients: A Randomized, Controlled, Multi-Center Trial. Spine, 2009, 34, 2248-2253.	1.0	27
104	The Effect of Electrical Stimulation on Lumbar Spinal Fusion in Older Patients: A Randomized, Controlled, Multi-Center Trial. Spine, 2009, 34, 2241-2247.	1.0	18
105	Shockwave therapy for chronic Achilles tendinopathy: A double-blind, randomized clinical trial of efficacy. Monthly Notices of the Royal Astronomical Society: Letters, 2008, 79, 249-256.	1.2	136
106	Epidural Steroid Following Discectomy for Herniated Lumbar Disc Reduces Neurological Impairment and Enhances Recovery. Spine, 2008, 33, 2028-2033.	1.0	53
107	Management of nerves during leg amputation – a neglected area in our understanding of the pathogenesis of phantom limb pain. Acta Anaesthesiologica Scandinavica, 2007, 51, 1115-1116.	0.7	18
108	Increased flexion and reduced hospital stay with continuous intraarticular morphine and ropivacaine after primary total knee replacementOpen intervention study of efficacy and safety in 154 patients. Acta Orthopaedica, 2004, 75, 606-609.	1.4	72

#	Article	IF	CITATIONS
109	Effects of metoprolol CR/XL on mortality and hospitalizations in patients with heart failure and history of hypertension. Journal of Cardiac Failure, 2002, 8, 8-14.	0.7	22
110	Combined intra-articular glucocorticoid, bupivacaine and morphine reduces pain and convalescence after diagnostic knee arthroscopy. Acta Orthopaedica, 2002, 73, 175-178.	1.4	36
111	Arthroscopic treatment of impingement of the ankle reduces pain and enhances function. Scandinavian Journal of Medicine and Science in Sports, 2002, 12, 69-72.	1.3	50
112	Intraarticular glucocorticoid, morphine and bupivacaine reduces pain and convalescence after arthroscopic ankle surgery: A randomized study of 36 patients. Acta Orthopaedica, 2000, 71, 301-304.	1.4	25
113	Intra-articular glucocorticoid, bupivacaine and morphine reduces pain, inflammatory response and convalescence after arthroscopic meniscectomy. Pain, 1998, 78, 131-134.	2.0	67
114	Being overweight and multiple fractures are indications for operative treatment of humeral shaft fractures. Injury, 1995, 26, 263-264.	0.7	26
115	Observer variation in the Lauge-Hansen classification of ankle fractures: Precision improved by instruction. Acta Orthopaedica, 1993, 64, 693-694.	1.4	56
116	The clinical effect of naproxen sodium after arthroscopy of the knee: A randomized, double-blind, prospective study. Arthroscopy - Journal of Arthroscopic and Related Surgery, 1993, 9, 375-380.	1.3	28
117	Displaced proximal humeral fractures: results of conservative treatment. Injury, 1992, 23, 41-43.	0.7	97
118	Closed Longitudinal Splitting of the Carpus. Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery, 1990, 24, 167-169.	0.6	2
119	Post-operative infections of osteoplastic compared with free bone flaps. British Journal of Neurosurgery, 1990, 4, 493-495.	0.4	27
120	Effects of amiloride on plasma and total body potassium, blood pressure, and the renin-angiotensin-aldosterone system in thiazide-treated hypertensive patients. Clinical Pharmacology and Therapeutics, 1983, 34, 448-453.	2.3	7
121	DELAYED HYPERSENSITIVITY AND BORDERLINE ESSENTIAL HYPERTENSION Delayed Hypersensitivity Against Arterial Wall Components in Patients suffering from Borderline Essential Hypertension. Acta Pathologica Et Microbiologica Scandinavica Section C, Immunology, 1977, 85C, 196-198.	0.0	1
122	Care-seeking behaviour of adolescents with patellofemoral pain: a retrospective cohort study. F1000Research, 0, 11, 161.	0.8	1