

Hongtu Zheng

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6542004/publications.pdf>

Version: 2024-02-01

16
papers

338
citations

840776

11
h-index

940533

16
g-index

16
all docs

16
docs citations

16
times ranked

643
citing authors

#	ARTICLE	IF	CITATIONS
1	HIF1 α promotes tumor chemoresistance via recruiting GDF15-producing TAMs in colorectal cancer. <i>Experimental Cell Research</i> , 2021, 398, 112394.	2.6	17
2	UPF1 promotes chemoresistance to oxaliplatin through regulation of TOP2A activity and maintenance of stemness in colorectal cancer. <i>Cell Death and Disease</i> , 2021, 12, 519.	6.3	19
3	Hypoxia Induces Growth Differentiation Factor 15 to Promote the Metastasis of Colorectal Cancer via PERK-eIF2 α Signaling. <i>BioMed Research International</i> , 2020, 2020, 1-12.	1.9	17
4	Prognosis of three histological subtypes of colorectal adenocarcinoma: A retrospective analysis of 8005 Chinese patients. <i>Cancer Medicine</i> , 2019, 8, 3411-3419.	2.8	25
5	Laparoscopic surgery may decrease the risk of clinical anastomotic leakage and a nomogram to predict anastomotic leakage after anterior resection for rectal cancer. <i>International Journal of Colorectal Disease</i> , 2019, 34, 319-328.	2.2	35
6	Silencing of long non-coding RNA SBDSP1 suppresses tumor growth and invasion in colorectal cancer. <i>Biomedicine and Pharmacotherapy</i> , 2017, 85, 355-361.	5.6	11
7	A study of peritoneal metastatic xenograft model of colorectal cancer in the treatment of hyperthermic intraperitoneal chemotherapy with Raltitrexed. <i>Biomedicine and Pharmacotherapy</i> , 2017, 92, 149-156.	5.6	10
8	Temporary Diverting Stoma Improves Recovery of Anastomotic Leakage after Anterior Resection for Rectal Cancer. <i>Scientific Reports</i> , 2017, 7, 15930.	3.3	20
9	Comparative study of single-center patients with thyroid metastases from colorectal cancer and previously reported cases in the literature. <i>World Journal of Surgical Oncology</i> , 2017, 15, 88.	1.9	14
10	Rectovaginal fistula after low anterior resection in Chinese patients with colorectal cancer. <i>Oncotarget</i> , 2017, 8, 73123-73132.	1.8	20
11	High preoperative peripheral blood neutrophil predicts poor outcome in rectal cancer treated with neoadjuvant chemoradiation therapy. <i>International Journal of Clinical and Experimental Pathology</i> , 2017, 10, 7718-7725.	0.5	1
12	Prognostic risk factors in patients with bone metastasis from colorectal cancer. <i>Tumor Biology</i> , 2016, 37, 16127-16134.	1.8	20
13	Neutrophil-to-lymphocyte and platelet-to-lymphocyte ratios predict chemotherapy outcomes and prognosis in patients with colorectal cancer and synchronous liver metastasis. <i>World Journal of Surgical Oncology</i> , 2016, 14, 289.	1.9	71
14	CapOX as neoadjuvant chemotherapy for locally advanced operable colon cancer patients: a prospective single-arm phase II trial. <i>Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association, Beijing Institute for Cancer Research</i> , 2016, 28, 589-597.	2.2	35
15	Pathological Features and Survival Outcomes of Young Patients with Operable Colon Cancer: Are They Homogeneous?. <i>PLoS ONE</i> , 2014, 9, e102004.	2.5	10
16	Risk Factors of Synchronous Inguinal Lymph Nodes Metastasis for Lower Rectal Cancer Involving the Anal Canal. <i>PLoS ONE</i> , 2014, 9, e111770.	2.5	13