

Raffaella Rastaldo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6546234/publications.pdf>

Version: 2024-02-01

45
papers

3,357
citations

304368

22
h-index

233125

45
g-index

46
all docs

46
docs citations

46
times ranked

3873
citing authors

#	ARTICLE	IF	CITATIONS
1	Silica Nanoparticle Internalization Improves Chemotactic Behaviour of Human Mesenchymal Stem Cells Acting on the SDF1 \pm /CXCR4 Axis. <i>Biomedicines</i> , 2022, 10, 336.	1.4	6
2	Advanced Nanotechnology for Enhancing Immune Checkpoint Blockade Therapy. <i>Nanomaterials</i> , 2021, 11, 661.	1.9	23
3	Targeting Cancer Cells Overexpressing Folate Receptors with New Terpolymer-Based Nanocapsules: Toward a Novel Targeted DNA Delivery System for Cancer Therapy. <i>Biomedicines</i> , 2021, 9, 1275.	1.4	24
4	Therapeutic Acellular Scaffolds for Limiting Left Ventricular Remodelling-Current Status and Future Directions. <i>International Journal of Molecular Sciences</i> , 2021, 22, 13054.	1.8	5
5	Nanoengineering in Cardiac Regeneration: Looking Back and Going Forward. <i>Nanomaterials</i> , 2020, 10, 1587.	1.9	14
6	Dual Role of Autophagy in Regulation of Mesenchymal Stem Cell Senescence. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 276.	1.8	36
7	Cardioprotection of PLGA/gelatine cardiac patches functionalised with adenosine in a large animal model of ischaemia and reperfusion injury: A feasibility study. <i>Journal of Tissue Engineering and Regenerative Medicine</i> , 2019, 13, 1253-1264.	1.3	22
8	Apelin α -induced cardioprotection against ischaemia/reperfusion injury: roles of epidermal growth factor and Src. <i>Acta Physiologica</i> , 2018, 222, e12924.	1.8	22
9	Silica nanoparticles actively engage with mesenchymal stem cells in improving acute functional cardiac integration. <i>Nanomedicine</i> , 2018, 13, 1121-1138.	1.7	21
10	Role of three adipokines in metabolic syndrome. <i>Polish Archives of Internal Medicine</i> , 2016, 126, 219-221.	0.3	2
11	Alpha-linolenic acid protects against cardiac injury and remodelling induced by beta-adrenergic overstimulation. <i>Food and Function</i> , 2015, 6, 2231-2239.	2.1	21
12	Effects of apelin on the cardiovascular system. <i>Heart Failure Reviews</i> , 2015, 20, 505-518.	1.7	73
13	Preconditioning cardioprotection and exercise performance: a radical point of view. <i>Sport Sciences for Health</i> , 2015, 11, 137-151.	0.4	6
14	The effect of bioartificial constructs that mimic myocardial structure and biomechanical properties on stem cell commitment towards cardiac lineage. <i>Biomaterials</i> , 2014, 35, 92-104.	5.7	27
15	Injured cardiomyocytes promote dental pulp mesenchymal stem cell homing. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2014, 1840, 2152-2161.	1.1	15
16	Balance of Nitric Oxide and Reactive Oxygen Species in Myocardial Reperfusion Injury and Protection. <i>Journal of Cardiovascular Pharmacology</i> , 2013, 62, 567-575.	0.8	22
17	A Lipophilic Nitric Oxide Donor and a Lipophilic Antioxidant Compound Protect Rat Heart Against Ischemia α Reperfusion Injury if Given as Hybrid Molecule but Not as a Mixture. <i>Journal of Cardiovascular Pharmacology</i> , 2012, 59, 241-248.	0.8	8
18	Effect of Apelin-Apelin Receptor System in Postischaemic Myocardial Protection: A Pharmacological Postconditioning Tool?. <i>Antioxidants and Redox Signaling</i> , 2011, 14, 909-922.	2.5	31

#	ARTICLE	IF	CITATIONS
19	Apelin-13 limits infarct size and improves cardiac postischemic mechanical recovery only if given after ischemia. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2011, 300, H2308-H2315.	1.5	68
20	Low concentrations of an nitric oxide-donor combined with a liposoluble antioxidant compound enhance protection against reperfusion injury in isolated rat hearts. <i>Journal of Physiology and Pharmacology</i> , 2010, 61, 21-7.	1.1	6
21	Effects of Nitric Oxide Donor Antioxidants Containing the Phenol Vitamin E Substructure and a Furoxan Moiety on Ischemia/Reperfusion Injury. <i>Arzneimittelforschung</i> , 2009, 59, 111-116.	0.5	1
22	Cardioprotection: A radical view. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 2009, 1787, 781-793.	0.5	176
23	Postconditioning cardioprotection against infarct size and post-ischemic systolic dysfunction is influenced by gender. <i>Basic Research in Cardiology</i> , 2009, 104, 390-402.	2.5	70
24	Omega 3 has a beneficial effect on ischemia/reperfusion injury, but cannot reverse the effect of stressful forced exercise. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2009, 19, 20-26.	1.1	17
25	HYPEROXIA CONFERS MYOCARDIAL PROTECTION IN MECHANICALLY VENTILATED RATS THROUGH THE GENERATION OF FREE RADICALS AND OPENING OF MITOCHONDRIAL ATP-SENSITIVE POTASSIUM CHANNELS. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2008, 35, 64-71.	0.9	13
26	Early homing of adult mesenchymal stem cells in normal and infarcted isolated beating hearts. <i>Journal of Cellular and Molecular Medicine</i> , 2008, 12, 507-521.	1.6	25
27	Local Activation or Implantation of Cardiac Progenitor Cells Rescues Scarred Infarcted Myocardium Improving Cardiac Function. <i>Circulation Research</i> , 2008, 103, 107-116.	2.0	266
28	Intermittent activation of bradykinin B2 receptors and mitochondrial KATP channels trigger cardiac postconditioning through redox signaling. <i>Cardiovascular Research</i> , 2007, 75, 168-177.	1.8	128
29	Nitric oxide and cardiac function. <i>Life Sciences</i> , 2007, 81, 779-793.	2.0	188
30	Human recombinant chromogranin A-derived vasostatin-1 mimics preconditioning via an adenosine/nitric oxide signaling mechanism. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2007, 293, H719-H727.	1.5	60
31	Effect of endothelins on the cardiovascular system. <i>Journal of Cardiovascular Medicine</i> , 2006, 7, 645-652.	0.6	44
32	Postconditioning reduces infarct size in the isolated rat heart: Role of coronary flow and pressure and the nitric oxide/cGMP pathway. <i>Basic Research in Cardiology</i> , 2006, 101, 168-179.	2.5	118
33	Postconditioning induced cardioprotection requires signaling through a redox-sensitive mechanism, mitochondrial ATP-sensitive K ⁺ channel and protein kinase C activation. <i>Basic Research in Cardiology</i> , 2006, 101, 180-189.	2.5	222
34	Exogenous High-Mobility Group Box 1 Protein Induces Myocardial Regeneration After Infarction via Enhanced Cardiac C-Kit + Cell Proliferation and Differentiation. <i>Circulation Research</i> , 2005, 97, e73-83.	2.0	256
35	Cardiac Stem Cells Possess Growth Factor-Receptor Systems That After Activation Regenerate the Infarcted Myocardium, Improving Ventricular Function and Long-Term Survival. <i>Circulation Research</i> , 2005, 97, 663-673.	2.0	494
36	Cardiac stem cells delivered intravascularly traverse the vessel barrier, regenerate infarcted myocardium, and improve cardiac function. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 3766-3771.	3.3	458

#	ARTICLE	IF	CITATIONS
37	FOF1ATP synthase activity is differently modulated by coronary reactive hyperemia before and after ischemic preconditioning in the goat. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2004, 287, H2192-H2200.	1.5	25
38	Endothelial cytochrome P450 contributes to the acetylcholine-induced cardiodepression in isolated rat hearts. <i>Acta Physiologica Scandinavica</i> , 2004, 182, 11-20.	2.3	3
39	Nitroxyl affords thiol-sensitive myocardial protective effects akin to early preconditioning. <i>Free Radical Biology and Medicine</i> , 2003, 34, 33-43.	1.3	193
40	Ischemic preconditioning changes the pattern of coronary reactive hyperemia regardless of mitochondrial ATP-sensitive K ⁺ channel blockade. <i>Life Sciences</i> , 2002, 71, 2299-2309.	2.0	5
41	Fatty acids are important for the Frank-Starling mechanism and Gregg effect but not for catecholamine response in isolated rat hearts. <i>Acta Physiologica Scandinavica</i> , 2002, 176, 167-176.	2.3	8
42	Ischemic preconditioning. <i>Life Sciences</i> , 2001, 69, 1-15.	2.0	91
43	Comparison between the effects of pentobarbital or ketamine/nitrous oxide anesthesia on metabolic and endothelial components of coronary reactive hyperemia. <i>Life Sciences</i> , 2001, 69, 729-738.	2.0	14
44	Cytochrome P-450 metabolite of arachidonic acid mediates bradykinin-induced negative inotropic effect. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2001, 280, H2823-H2832.	1.5	26
45	Mitochondrial ATP-sensitive channel opener does not induce vascular preconditioning, but potentiates the effect of a preconditioning ischemia on coronary reactive hyperemia in the anesthetized goat. <i>Pflugers Archiv European Journal of Physiology</i> , 2001, 443, 166-174.	1.3	4