

Yuping Wang

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6547590/yuping-wang-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

75
papers

2,215
citations

25
h-index

45
g-index

76
ext. papers

2,489
ext. citations

3.4
avg, IF

4.82
L-index

#	Paper	IF	Citations
75	Preeclampsia Status Controls Interleukin-6 and Soluble IL-6 Receptor Release from Neutrophils and Endothelial Cells: Relevance to Increased Inflammatory Responses.. <i>Pathophysiology</i> , 2021 , 28, 202-211	1.8	2
74	Downregulation of miR-126-3p expression contributes to increased inflammatory response in placental trophoblasts in preeclampsia. <i>Journal of Reproductive Immunology</i> , 2021 , 144, 103281	4.2	5
73	Upregulation of histone H3K9 methylation in fetal endothelial cells from preeclamptic pregnancies. <i>Journal of Cellular Physiology</i> , 2021 , 236, 1866-1874	7	2
72	Upregulation of METTL3 expression and m6A RNA methylation in placental trophoblasts in preeclampsia. <i>Placenta</i> , 2021 , 103, 43-49	3.4	9
71	Association of fetal gender and the onset and severity of hypertensive disorders of pregnancy. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2020 , 1-6	2	
70	Maternal soluble PD-1 levels are significantly increased in women with preeclampsia. <i>American Journal of Reproductive Immunology</i> , 2020 , 83, e13193	3.8	4
69	Prolonged Fetal Heart Rate Decelerations in Labor: Can We Reduce Unplanned Primary Cesarean Sections in This Group?. <i>Advances in Therapy</i> , 2020 , 37, 4325-4335	4.1	1
68	Downregulation of vitamin D receptor and miR-126-3p expression contributes to increased endothelial inflammatory response in preeclampsia. <i>American Journal of Reproductive Immunology</i> , 2019 , 82, e13172	3.8	18
67	Human Placental Stem Cell Therapy in Stroke: Endothelial / Smooth Muscle Mechanisms Underlying Protection?. <i>FASEB Journal</i> , 2019 , 33, 524.1	0.9	
66	3D Printing for Bio-Synthetic Biliary Stents. <i>Bioengineering</i> , 2019 , 6,	5.3	19
65	Histone deacetylase inhibition disturbs the balance between ACE and chymase expression in endothelial cells: a potential mechanism of chymase activation in preeclampsia. <i>Hypertension Research</i> , 2019 , 42, 155-164	4.7	5
64	Aberrant pro-atrial natriuretic peptide/corin/natriuretic peptide receptor signaling is present in maternal vascular endothelium in preeclampsia. <i>Pregnancy Hypertension</i> , 2018 , 11, 1-6	2.6	16
63	Reduced CD200 expression is associated with altered Th1/Th2 cytokine production in placental trophoblasts from preeclampsia. <i>American Journal of Reproductive Immunology</i> , 2018 , 79, e12763	3.8	21
62	Loss of slit protein nephrin is associated with reduced antioxidant superoxide dismutase expression in podocytes shed from women with preeclampsia. <i>Physiological Reports</i> , 2018 , 6, e13785	2.6	8
61	Human placental derived stem cells protection in stroke injury.. <i>FASEB Journal</i> , 2018 , 32, 740.1	0.9	
60	Human Placental Stem Cell Therapy in Stroke: Endothelial/Smooth Muscle Mechanisms Underlying Protection. <i>FASEB Journal</i> , 2018 , 32, 575.2	0.9	
59	High-throughput scaffold-free microtissues through 3D printing. <i>3D Printing in Medicine</i> , 2018 , 4, 9	5	12

58	Notch ligand Jagged1 promotes mesenchymal stromal cell-based cartilage repair. <i>Experimental and Molecular Medicine</i> , 2018 , 50, 1-10	12.8	11
57	IL-1 β reduces cardiac lymphatic muscle contraction via COX-2 and PGE induction: Potential role in myocarditis. <i>Biomedicine and Pharmacotherapy</i> , 2018 , 107, 1591-1600	7.5	11
56	Vitamin D Reduces Oxidative Stress-Induced Procaspase-3/ROCK1 Activation and MP Release by Placental Trophoblasts. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017 , 102, 2100-2110	5.6	19
55	Vitamin D suppresses oxidative stress-induced microparticle release by human umbilical vein endothelial cells. <i>Biology of Reproduction</i> , 2017 , 96, 199-210	3.9	6
54	Vitamin D suppresses oxidative stress-induced microparticle release by human umbilical vein endothelial cells. <i>Biology of Reproduction</i> , 2017 , 96, 199-210	3.9	10
53	Upregulation of cathepsin C expression contributes to endothelial chymase activation in preeclampsia. <i>Hypertension Research</i> , 2017 , 40, 976-981	4.7	5
52	Up-regulation of miR-203 expression induces endothelial inflammatory response: Potential role in preeclampsia. <i>American Journal of Reproductive Immunology</i> , 2016 , 76, 482-490	3.8	21
51	1,25(OH) $_2$ D $_3$ Induces Placental Vascular Smooth Muscle Cell Relaxation by Phosphorylation of Myosin Phosphatase Target Subunit 1Ser507: Potential Beneficial Effects of Vitamin D on Placental Vasculature in Humans. <i>Biology of Reproduction</i> , 2016 , 94, 116	3.9	6
50	Expectant management of mild preeclampsia versus superimposed preeclampsia up to 37 weeks. <i>American Journal of Obstetrics and Gynecology</i> , 2015 , 212, 515.e1-8	6.4	16
49	Increased urinary levels of podocyte glycoproteins, matrix metalloproteinases, inflammatory cytokines, and kidney injury biomarkers in women with preeclampsia. <i>American Journal of Physiology - Renal Physiology</i> , 2015 , 309, F1009-17	4.3	24
48	Activation of vitamin D receptor promotes VEGF and CuZn-SOD expression in endothelial cells. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2014 , 140, 56-62	5.1	69
47	Down-regulation of TIMP3 leads to increase in TACE expression and TNF α production by placental trophoblast cells. <i>American Journal of Reproductive Immunology</i> , 2014 , 71, 427-33	3.8	5
46	Differential miRNA expression profiles between the first and third trimester human placentas. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2013 , 304, E836-43	6	101
45	Role of chymase in preeclampsia. <i>Current Vascular Pharmacology</i> , 2013 , 11, 606-15	3.3	2
44	Increased urinary excretion of nephrin, podocalyxin, and Ig-h3 in women with preeclampsia. <i>American Journal of Physiology - Renal Physiology</i> , 2012 , 302, F1084-9	4.3	50
43	Digoxin immune fab protects endothelial cells from ouabain-induced barrier injury. <i>American Journal of Reproductive Immunology</i> , 2012 , 67, 66-72	3.8	3
42	Expressions of vitamin D metabolic components VDBP, CYP2R1, CYP27B1, CYP24A1, and VDR in placentas from normal and preeclamptic pregnancies. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2012 , 303, E928-35	6	89
41	Elevated maternal soluble Gp130 and IL-6 levels and reduced Gp130 and SOCS-3 expressions in women complicated with preeclampsia. <i>Hypertension</i> , 2011 , 57, 336-42	8.5	22

40	Factors derived from preeclamptic placentas perturb polarity protein PARD-3 expression and distribution in endothelial cells. <i>Reproductive Sciences</i> , 2011 , 18, 164-71	3	5
39	Isolation and Culture of Human Umbilical Vein Endothelial Cells 2011 , 163-169		0
38	Altered nephrin and podoplanin distribution is associated with disturbed polarity protein PARD-3 and PARD-6 expressions in podocytes from preeclampsia. <i>Reproductive Sciences</i> , 2011 , 18, 772-80	3	19
37	Analysis of endothelial barrier function in vitro. <i>Methods in Molecular Biology</i> , 2011 , 763, 253-64	1.4	12
36	Elevated plasma chymotrypsin-like protease (chymase) activity in women with preeclampsia. <i>Hypertension in Pregnancy</i> , 2010 , 29, 253-61	2	23
35	Vascular Biology of the Placenta. <i>Colloquium Series on Integrated Systems Physiology From Molecule To Function</i> , 2010 , 2, 1-98		58
34	Placenta-derived chymotrypsin-like protease (CLP) disturbs endothelial junctional structure in preeclampsia. <i>Reproductive Sciences</i> , 2009 , 16, 479-88	3	10
33	Chymotrypsin-like protease (chymase) mediates endothelial activation by factors derived from preeclamptic placentas. <i>Reproductive Sciences</i> , 2009 , 16, 905-13	3	8
32	Decreased nephrin and GLEPP-1, but increased VEGF, Flt-1, and nitrotyrosine, expressions in kidney tissue sections from women with preeclampsia. <i>Reproductive Sciences</i> , 2009 , 16, 970-9	3	48
31	Maternal circulating TNF-alpha levels are highly correlated with IL-10 levels, but not IL-6 and IL-8 levels, in women with pre-eclampsia. <i>American Journal of Reproductive Immunology</i> , 2009 , 62, 269-74	3.8	28
30	Prostacyclin and thromboxane levels in women with severe preeclampsia undergoing magnesium sulfate therapy during antepartum and postpartum periods. <i>Hypertension in Pregnancy</i> , 2008 , 27, 17-27	2	3
29	Placental productions and expressions of soluble endoglin, soluble fms-like tyrosine kinase receptor-1, and placental growth factor in normal and preeclamptic pregnancies. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 260-6	5.6	143
28	Elevated maternal IL-16 levels, enhanced IL-16 expressions in endothelium and leukocytes, and increased IL-16 production by placental trophoblasts in women with preeclampsia. <i>Journal of Immunology</i> , 2008 , 181, 4418-22	5.3	28
27	Endothelial angiotensin II generation induced by placenta-derived factors from preeclampsia. <i>Reproductive Sciences</i> , 2008 , 15, 932-8	3	7
26	Endothelial barrier function in preeclampsia. <i>Frontiers in Bioscience - Landmark</i> , 2007 , 12, 2412-24	2.8	20
25	Proteomic analysis of human cerebral endothelial cells activated by MS serum and IFN B-1b. <i>FASEB Journal</i> , 2007 , 21, A849	0.9	
24	Reduced cellular glutathione reductase activity and increased adhesion molecule expression in endothelial cells cultured with maternal plasma from women with preeclampsia. <i>Journal of the Society for Gynecologic Investigation</i> , 2006 , 13, 412-7		10
23	Increased superoxide generation and decreased stress protein Hsp90 expression in human umbilical cord vein endothelial cells (HUVECs) from pregnancies complicated by preeclampsia. <i>Hypertension in Pregnancy</i> , 2006 , 25, 169-82	2	29

22	Effects of peroxynitrite and superoxide radicals on endothelial monolayer permeability: potential role of peroxynitrite in preeclampsia. <i>Journal of the Society for Gynecologic Investigation</i> , 2005 , 12, 586-92		25
21	Placental tissue levels of nonesterified polyunsaturated fatty acids in normal and preeclamptic pregnancies. <i>Hypertension in Pregnancy</i> , 2005 , 24, 235-45	2	33
20	Increased phospholipase A2 and thromboxane but not prostacyclin production by placental trophoblast cells from normal and preeclamptic pregnancies cultured under hypoxia condition. <i>Placenta</i> , 2005 , 26, 402-9	3.4	41
19	Hypoxia promotes interleukin-6 and -8 but reduces interleukin-10 production by placental trophoblast cells from preeclamptic pregnancies. <i>Journal of the Society for Gynecologic Investigation</i> , 2005 , 12, 428-32		46
18	Placental trophoblast-derived factors diminish endothelial barrier function. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004 , 89, 2421-8	5.6	33
17	Evidence of endothelial dysfunction in preeclampsia: decreased endothelial nitric oxide synthase expression is associated with increased cell permeability in endothelial cells from preeclampsia. <i>American Journal of Obstetrics and Gynecology</i> , 2004 , 190, 817-24	6.4	90
16	Increased endothelial monolayer permeability is induced by serum from women with preeclampsia but not by serum from women with normal pregnancy or that are not pregnant. <i>Hypertension in Pregnancy</i> , 2003 , 22, 99-108	2	31
15	Antioxidant Superoxide Dismutase Attenuates Increased Endothelial Permeability Induced By Platelet-Activating Factor. <i>Journal of the Society for Gynecologic Investigation</i> , 2003 , 10, 5-10		1
14	Antioxidant superoxide dismutase attenuates increased endothelial permeability induced by platelet-activating factor. <i>Journal of the Society for Gynecologic Investigation</i> , 2003 , 10, 5-10		3
13	Endothelial junctional protein redistribution and increased monolayer permeability in human umbilical vein endothelial cells isolated during preeclampsia. <i>American Journal of Obstetrics and Gynecology</i> , 2002 , 186, 214-20	6.4	56
12	Expression of thrombin receptors in endothelial cells and neutrophils from normal and preeclamptic pregnancies. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2002 , 87, 3728-34	5.6	33
11	Maternal Perfusion with Low-Dose Aspirin Preferentially Inhibits Placental Thromboxane While Sparing Prostacyclin. <i>Hypertension in Pregnancy</i> , 1998 , 17, 203-215	2	13
10	The Ratio of Thromboxane to Prostacyclin is Increased by Peroxide in a Dose-Dependent Manner, Along with Increased Vasoconstriction in the Human Placenta. <i>Hypertension in Pregnancy</i> , 1998 , 17, 1-11 ²		6
9	Activation of Endothelial Cells in Preeclampsia: Increased Neutrophil-Endothelial Adhesion Correlates With Up-regulation of Adhesion Molecule P-selectin in Human Umbilical Vein Endothelial Cells Isolated From Preeclampsia. <i>Journal of the Society for Gynecologic Investigation</i> , 1998 , 5, 237-243		3
8	Vitamin E Attenuates Peroxide-induced Vasoconstriction in the Human Placenta. <i>Hypertension in Pregnancy</i> , 1997 , 16, 389-401	2	2
7	Placental Production of Lipid Peroxides, Thromboxane, And Prostacyclin in Preeclampsia. <i>Hypertension in Pregnancy</i> , 1996 , 15, 101-111	2	30
6	Antioxidant Activities and mRNA Expression of Superoxide Dismutase, Catalase, and Glutathione Peroxidase in Normal and Preeclamptic Placentas. <i>Journal of the Society for Gynecologic Investigation</i> , 1996 , 3, 179-184		61
5	TNF alpha concentrations and mRNA expression are increased in preeclamptic placentas. <i>Journal of Reproductive Immunology</i> , 1996 , 32, 157-69	4.2	175

4	Placental Production of Nitric Oxide and Endothelin in Normal and Preeclamptic Pregnancies. <i>Hypertension in Pregnancy</i> , 1994 , 13, 171-178	2	23
3	Placental lipid peroxides and thromboxane are increased and prostacyclin is decreased in women with preeclampsia. <i>American Journal of Obstetrics and Gynecology</i> , 1992 , 167, 946-9	6.4	177
2	The imbalance between thromboxane and prostacyclin in preeclampsia is associated with an imbalance between lipid peroxides and vitamin E in maternal blood. <i>American Journal of Obstetrics and Gynecology</i> , 1991 , 165, 1695-700	6.4	230
1	Decreased levels of polyunsaturated fatty acids in preeclampsia. <i>American Journal of Obstetrics and Gynecology</i> , 1991 , 164, 812-8	6.4	60