

Camilla Steen Jensen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6550964/publications.pdf>

Version: 2024-02-01

13
papers

291
citations

1162889

8
h-index

1372474

10
g-index

13
all docs

13
docs citations

13
times ranked

628
citing authors

#	ARTICLE	IF	CITATIONS
1	Exercise as a potential modulator of inflammation in patients with Alzheimer's disease measured in cerebrospinal fluid and plasma. <i>Experimental Gerontology</i> , 2019, 121, 91-98.	1.2	72
2	Biochemical Markers of Physical Exercise on Mild Cognitive Impairment and Dementia: Systematic Review and Perspectives. <i>Frontiers in Neurology</i> , 2015, 6, 187.	1.1	49
3	Cerebrospinal Fluid Amyloid Beta and Tau Concentrations Are Not Modulated by 16 Weeks of Moderate- to High-Intensity Physical Exercise in Patients with Alzheimer Disease. <i>Dementia and Geriatric Cognitive Disorders</i> , 2016, 42, 146-158.	0.7	40
4	Patients with Alzheimer's disease who carry the <i>APOE</i> ϵ 4 allele benefit more from physical exercise. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2019, 5, 99-106.	1.8	40
5	Effect of physical exercise on markers of neuronal dysfunction in cerebrospinal fluid in patients with Alzheimer's disease. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2017, 3, 284-290.	1.8	23
6	Lactoferrin in cerebrospinal fluid and saliva is not a diagnostic biomarker for Alzheimer's disease in a mixed memory clinic population. <i>EBioMedicine</i> , 2021, 67, 103361.	2.7	23
7	Variants of the <i>ADRB2</i> Gene in COPD: Systematic Review and Meta-Analyses of Disease Risk and Treatment Response. <i>COPD: Journal of Chronic Obstructive Pulmonary Disease</i> , 2017, 14, 451-460.	0.7	22
8	Txl1 and Txc1 Are Co-Factors of the 26S Proteasome in Fission Yeast. <i>Antioxidants and Redox Signaling</i> , 2011, 14, 1601-1608.	2.5	18
9	Physical Exercise May Increase Plasma Concentration of High-Density Lipoprotein-Cholesterol in Patients With Alzheimer's Disease. <i>Frontiers in Neuroscience</i> , 2020, 14, 532.	1.4	3
10	Cerebrospinal fluid and saliva lactoferrin as a diagnostic biomarker for Alzheimer's disease in a mixed memory clinic population. <i>Alzheimer's and Dementia</i> , 2021, 17, .	0.4	1
11	Pharmacogenetics of COPD: a role for the β 2-adrenergic receptor gene?. <i>Lancet Respiratory Medicine</i> , 2014, 2, 3-4.	5.2	0
12	P1-069: Moderate-to-high intensity physical training does not alter cerebrospinal amyloid- β 1-42 levels in patients with Alzheimer's disease. , 2015, 11, P364-P365.		0
13	[P3-174]: EFFECT OF PHYSICAL EXERCISE ON MARKERS OF NEURONAL DYSFUNCTION IN CEREBROSPINAL FLUID IN PATIENTS WITH ALZHEIMER'S DISEASE. <i>Alzheimer's and Dementia</i> , 2017, 13, P1000.	0.4	0