Yutaka J Matsuoka

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6554329/publications.pdf

Version: 2024-02-01

109264 133188 4,417 143 35 59 citations g-index h-index papers 152 152 152 6060 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Nutritional medicine as mainstream in psychiatry. Lancet Psychiatry, the, 2015, 2, 271-274.	3.7	375
2	Smaller regional volumes of brain gray and white matter demonstrated in breast cancer survivors exposed to adjuvant chemotherapy. Cancer, 2007, 109, 146-156.	2.0	298
3	A systematic review of predictors of posttraumatic stress disorder (PTSD) for adult road traffic crash survivors. Injury, 2013, 44, 1413-1422.	0.7	153
4	Estimating the risk of PTSD in recent trauma survivors: results of the International Consortium to Predict PTSD (ICPP). World Psychiatry, 2019, 18, 77-87.	4.8	126
5	Reliability and validity of the Japanese version of the Resilience Scale and its short version. BMC Research Notes, 2010, 3, 310.	0.6	118
6	International Society for Nutritional Psychiatry Research Practice Guidelines for Omega-3 Fatty Acids in the Treatment of Major Depressive Disorder. Psychotherapy and Psychosomatics, 2019, 88, 263-273.	4.0	114
7	Posttraumatic growth, posttraumatic stress disorder and resilience of motor vehicle accident survivors. BioPsychoSocial Medicine, 2010, 4, 7.	0.9	103
8	Association of Delirium Response and Safety of Pharmacological Interventions for the Management and Prevention of Delirium. JAMA Psychiatry, 2019, 76, 526.	6.0	98
9	Effect of bifidobacterium breve A-1 on anxiety and depressive symptoms in schizophrenia: A proof-of-concept study. Journal of Affective Disorders, 2019, 245, 377-385.	2.0	90
10	Omega-3 Polyunsaturated Fatty Acids in Prevention of Mood and Anxiety Disorders. Clinical Psychopharmacology and Neuroscience, 2015, 13, 129-137.	0.9	85
11	A volumetric study of amygdala in cancer survivors with intrusive recollections. Biological Psychiatry, 2003, 54, 736-743.	0.7	82
12	Association of Use of Omega-3 Polyunsaturated Fatty Acids With Changes in Severity of Anxiety Symptoms. JAMA Network Open, 2018, 1, e182327.	2.8	82
13	<scp>International $<$ scp>Society for $<$ scp>Nutritional $<$ scp>Psychiatry $<$ scp>Research consensus position statement: nutritional medicine in modern psychiatry. World Psychiatry, 2015, 14, 370-371.	4.8	81
14	Peritraumatic Distress Inventory as a predictor of postâ€traumatic stress disorder after a severe motor vehicle accident. Psychiatry and Clinical Neurosciences, 2010, 64, 149-156.	1.0	73
15	Structure of orbitofrontal cortex and its longitudinal course in cancer-related post-traumatic stress disorder. Neuroscience Research, 2007, 59, 383-389.	1.0	71
16	Relationship Between Distressing Cancer-Related Recollections and Hippocampal Volume in Cancer Survivors. American Journal of Psychiatry, 2002, 159, 2087-2093.	4.0	67
17	Peritraumatic Distress, Watching Television, and Posttraumatic Stress Symptoms among Rescue Workers after the Great East Japan Earthquake. PLoS ONE, 2012, 7, e35248.	1.1	67
18	Association between adjuvant regional radiotherapy and cognitive function in breast cancer patients treated with conservation therapy. Cancer Medicine, 2014, 3, 702-709.	1.3	65

#	Article	IF	Citations
19	Dietary fish, n-3 polyunsaturated fatty acid consumption, and depression risk in Japan: a population-based prospective cohort study. Translational Psychiatry, 2017, 7, e1242-e1242.	2.4	62
20	Concern over radiation exposure and psychological distress among rescue workers following the Great East Japan Earthquake Concern over radiation exposure and distress. BMC Public Health, 2012, 12, 249.	1.2	60
21	No adverse effects of adjuvant chemotherapy on hippocampal volumein Japanese breast cancer survivors. Breast Cancer Research and Treatment, 2005, 92, 81-84.	1.1	54
22	Associations between Disaster Exposures, Peritraumatic Distress, and Posttraumatic Stress Responses in Fukushima Nuclear Plant Workers following the 2011 Nuclear Accident: The Fukushima NEWS Project Study. PLoS ONE, 2014, 9, e87516.	1.1	51
23	Clinical utility and validation of the Japanese version of Memorial Delirium Assessment Scale in a psychogeriatric inpatient setting. General Hospital Psychiatry, 2001, 23, 36-40.	1.2	49
24	Possibility of Polyunsaturated Fatty Acids for the Prevention and Treatment of Neuropsychiatric Illnesses. Journal of Pharmacological Sciences, 2014, 124, 294-300.	1.1	49
25	Cancer-Related Intrusive Thoughts as an Indicator of Poor Psychological Adjustment at 3 or More Years After Breast Surgery: A Preliminary Study. Breast Cancer Research and Treatment, 2002, 76, 117-124.	1.1	47
26	Resilience, postâ€traumatic growth, and work engagement among health care professionals after the Great East Japan Earthquake: A 4â€year prospective followâ€up study. Journal of Occupational Health, 2016, 58, 347-353.	1.0	47
27	Occurrence of fatigue and associated factors in disease-free breast cancer patients without depression. Supportive Care in Cancer, 2005, 13, 628-636.	1.0	44
28	Incidence and prediction of psychiatric morbidity after a motor vehicle accident in Japan: The Tachikawa Cohort of Motor Vehicle Accident Study. Critical Care Medicine, 2008, 36, 74-80.	0.4	44
29	Omega-3 Fatty Acids for Secondary Prevention of Posttraumatic Stress Disorder After Accidental Injury. Journal of Clinical Psychopharmacology, 2010, 30, 217-219.	0.7	43
30	Preoperative Anxiety as a Predictor of Delirium in Cancer Patients: A Prospective Observational Cohort Study. World Journal of Surgery, 2019, 43, 134-142.	0.8	43
31	Differentiating PTSD from anxiety and depression: Lessons from the ICDâ€11 PTSD diagnostic criteria. Depression and Anxiety, 2019, 36, 490-498.	2.0	43
32	Relationship of psychological characteristics and selfâ€efficacy in gastrointestinal cancer survivors. Psycho-Oncology, 2010, 19, 71-76.	1.0	40
33	Reliability and validity of the Japanese version of the Peritraumatic Distress Inventory. General Hospital Psychiatry, 2009, 31, 75-79.	1.2	38
34	Prefrontal Cortex and Amygdala Volume in First Minor or Major Depressive Episode After Cancer Diagnosis. Biological Psychiatry, 2006, 59, 707-712.	0.7	37
35	Hippocampal Volume and First Major Depressive Episode After Cancer Diagnosis in Breast Cancer Survivors. American Journal of Psychiatry, 2004, 161, 2263-2270.	4.0	36
36	Potential Role of Brain-Derived Neurotrophic Factor in Omega–3 Fatty Acid Supplementation to Prevent Posttraumatic Distress after Accidental Injury: An Open-Label Pilot Study. Psychotherapy and Psychosomatics, 2011, 80, 310-312.	4.0	36

#	Article	IF	Citations
37	Fish Oil for Attenuating Posttraumatic Stress Symptoms among Rescue Workers after the Great East Japan Earthquake: A Randomized Controlled Trial. Psychotherapy and Psychosomatics, 2012, 81, 315-317.	4.0	36
38	Culturally sensitive and universal measure of resilience for ⟨scp⟩J⟨ scp⟩apanese populations: ⟨scp⟩T⟨ scp⟩achikawa ⟨scp⟩R⟨ scp⟩esilience ⟨scp⟩S⟨ scp⟩cale in comparison with ⟨scp⟩R⟨ scp⟩esilience ⟨scp⟩S⟨ scp⟩cale 14â€item version. Psychiatry and Clinical Neurosciences, 2013, 67, 174-181.	1.0	36
39	Longitudinal Changes in Serum Brain-Derived Neurotrophic Factor in Accident Survivors with Posttraumatic Stress Disorder. Neuropsychobiology, 2013, 68, 44-50.	0.9	36
40	Brief mindfulness-based stress management program for a better mental state in working populations - Happy Nurse Project: A randomized controlled trial✰✰. Journal of Affective Disorders, 2019, 251, 186-194.	2.0	36
41	Clearance of fear memory from the hippocampus through neurogenesis by omega-3 fatty acids: A novel preventive strategy for posttraumatic stress disorder?. BioPsychoSocial Medicine, 2011, 5, 3.	0.9	35
42	Docosahexaenoic Acid for Selective Prevention of Posttraumatic Stress Disorder Among Severely Injured Patients. Journal of Clinical Psychiatry, 2015, 76, e1015-e1022.	1.1	34
43	Probability of major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic interviews controlling for Hospital Anxiety and Depression Scale – Depression subscale scores: An individual participant data meta-analysis of 73 primary studies. Journal of Psychosomatic Research, 2020, 129, 109892.	1.2	33
44	Association between regular physical exercise and depressive symptoms mediated through social support and resilience in Japanese company workers: a cross-sectional study. BMC Public Health, 2016, 16, 553.	1.2	32
45	Impact of chemotherapy on the association between fear of cancer recurrence and the gut microbiota in breast cancer survivors. Brain, Behavior, and Immunity, 2020, 85, 186-191.	2.0	32
46	Posttraumatic stress disorder symptom trajectories within the first year following emergency department admissions: pooled results from the International Consortium to predict PTSD. Psychological Medicine, 2021, 51, 1129-1139.	2.7	32
47	Clinical implications of the proposed ICD-11 PTSD diagnostic criteria. Psychological Medicine, 2019, 49, 483-490.	2.7	31
48	The Tachikawa cohort of motor vehicle accident study investigating psychological distress: design, methods and cohort profiles. Social Psychiatry and Psychiatric Epidemiology, 2009, 44, 333-340.	1.6	30
49	Secondary traumatic stress and associated factors among Japanese nurses working in hospitals. International Journal of Nursing Practice, 2012, 18, 155-163.	0.8	30
50	Serum pro-BDNF/BDNF as a treatment biomarker for response to docosahexaenoic acid in traumatized people vulnerable to developing psychological distress: a randomized controlled trial. Translational Psychiatry, 2015, 5, e596-e596.	2.4	29
51	Regional cerebral glucose metabolism in patients with secondary depressive episodes after fatal pancreatic cancer diagnosis. Journal of Affective Disorders, 2007, 99, 231-236.	2.0	28
52	Data Validation and Verification Using Blockchain in a Clinical Trial for Breast Cancer: Regulatory Sandbox. Journal of Medical Internet Research, 2020, 22, e18938.	2.1	28
53	Effects of omega-3 polyunsaturated fatty acids on psychophysiological symptoms of post-traumatic stress disorder in accident survivors: A randomized, double-blind, placebo-controlled trial. Journal of Affective Disorders, 2017, 224, 27-31.	2.0	27
54	Cardiorespiratory fitness in breast cancer survivors: a randomised controlled trial of home-based smartphone supported high intensity interval training. BMJ Supportive and Palliative Care, 2022, 12, 33-37.	0.8	27

#	Article	IF	CITATIONS
55	A randomized controlled multicenter trial of post-suicide attempt case management for the prevention of further attempts in Japan (ACTION-J). BMC Public Health, 2009, 9, 364.	1.2	26
56	Fish consumption and resilience to depression in Japanese company workers: a cross-sectional study. Lipids in Health and Disease, 2015, 14, 51.	1.2	26
57	A mindfulness-based stress management program and treatment with omega-3 fatty acids to maintain a healthy mental state in hospital nurses (Happy Nurse Project): study protocol for a randomized controlled trial. Trials, 2015, 16, 36.	0.7	26
58	Glutamatergic system abnormalities in posttraumatic stress disorder. Psychopharmacology, 2015, 232, 4261-4268.	1.5	26
59	Effectiveness of assertive case management on repeat self-harm in patients admitted for suicide attempt: Findings from ACTION-J study. Journal of Affective Disorders, 2018, 225, 460-465.	2.0	25
60	Serum Oxytocin, Posttraumatic Coping and C-Reactive Protein in Motor Vehicle Accident Survivors by Gender. Neuropsychobiology, 2015, 71, 196-201.	0.9	24
61	Limited effect of omega-3 fatty acids on the quality of life in survivors of traumatic injury: A randomized, placebo-controlled trial. Prostaglandins Leukotrienes and Essential Fatty Acids, 2017, 127, 1-5.	1.0	22
62	Plasma estradiol levels and antidepressant effects of omega-3 fatty acids in pregnant women. Brain, Behavior, and Immunity, 2020, 85, 29-34.	2.0	22
63	Hippocampal and Amygdalar Volumes in Breast Cancer Survivors with Posttraumatic Stress Disorder. Journal of Neuropsychiatry and Clinical Neurosciences, 2008, 20, 302-308.	0.9	19
64	Attenuating posttraumatic distress with omega-3 polyunsaturated fatty acids among disaster medical assistance team members after the Great East Japan Earthquake: The APOP randomized controlled trial. BMC Psychiatry, 2011, 11, 132.	1.1	19
65	Potential impact of propofol immediately after motor vehicle accident on later symptoms of posttraumatic stress disorder at 6-month follow up: a retrospective cohort study. Critical Care, 2012, 16, R196.	2.5	18
66	The longitudinal mental health impact of Fukushima nuclear disaster exposures and public criticism among power plant workers: the Fukushima NEWS Project study. Psychological Medicine, 2016, 46, 3117-3125.	2.7	18
67	Application of data pooling to longitudinal studies of early post-traumatic stress disorder (PTSD): the International Consortium to Predict PTSD (ICPP) project. HA¶gre Utbildning, 2018, 9, 1476442.	1.4	18
68	The Efficacy of Omega-3 Fatty Acids for Depressive Symptoms among Pregnant Women in Japan and Taiwan: A Randomized, Double-Blind, Placebo-Controlled Trial (SYNCHRO; NCT01948596). Psychotherapy and Psychosomatics, 2019, 88, 122-124.	4.0	18
69	Prevalence and related factors of common mental disorders during pregnancy in Japan: a cross-sectional study. BioPsychoSocial Medicine, 2016, 10, 17.	0.9	17
70	Yokukansan for the treatment of preoperative anxiety and postoperative delirium in colorectal cancer patients: a retrospective study. Japanese Journal of Clinical Oncology, 2017, 47, 844-848.	0.6	17
71	Relationship between polyunsaturated fatty acid and eating disorders: Systematic review and meta-analysis. Prostaglandins Leukotrienes and Essential Fatty Acids, 2019, 142, 11-19.	1.0	17
72	Lipid and Energy Metabolism of the Gut Microbiota Is Associated with the Response to Probiotic Bifidobacterium breve Strain for Anxiety and Depressive Symptoms in Schizophrenia. Journal of Personalized Medicine, 2021, 11, 987.	1.1	17

#	Article	IF	CITATIONS
73	Incidence and Prediction of Post-Traumatic Stress Disorder at 6 Months After Motor Vehicle Accidents in Japan. Psychosomatics, 2013, 54, 263-271.	2.5	16
74	Serum Levels of Polyunsaturated Fatty Acids and the Risk of Posttraumatic Stress Disorder. Psychotherapy and Psychosomatics, 2013, 82, 408-410.	4.0	15
75	Omega-3 fatty acids for a better mental state in working populations - Happy Nurse Project: A 52-week randomized controlled trial. Journal of Psychiatric Research, 2018, 102, 72-80.	1.5	15
76	Diet quality and depression risk in a Japanese population: the Japan Public Health Center (JPHC)-based Prospective Study. Scientific Reports, 2019, 9, 7150.	1.6	15
77	Fish consumption and cardiovascular response during mental stress. BMC Research Notes, 2012, 5, 288.	0.6	14
78	Tachikawa project for prevention of posttraumatic stress disorder with polyunsaturated fatty acid (TPOP): study protocol for a randomized controlled trial. BMC Psychiatry, 2013, 13, 8.	1.1	14
79	Explicit prognostic disclosure to Asian women with breast cancer: A randomized, scripted videoâ€vignette study (J‣UPPORT1601). Cancer, 2019, 125, 3320-3329.	2.0	14
80	Expectations of and recommendations for a cancer survivorship guideline in Japan: a literature review of guidelines for cancer survivorship. Japanese Journal of Clinical Oncology, 2019, 49, 812-822.	0.6	14
81	Association Between Dietary Fish and PUFA Intake in Midlife and Dementia in Later Life: The JPHC Saku Mental Health Study. Journal of Alzheimer's Disease, 2021, 79, 1091-1104.	1.2	14
82	Pharmacologic and hormonal treatments for menopausal sleep disturbances: A network meta-analysis of 43 randomized controlled trials and 32,271 menopausal women. Sleep Medicine Reviews, 2021, 57, 101469.	3.8	14
83	Post-Traumatic Stress Symptoms and Burnout Among Medical Rescue Workers 4 Years After the Great East Japan Earthquake: A Longitudinal Study. Disaster Medicine and Public Health Preparedness, 2016, 10, 848-853.	0.7	14
84	Biomedical and Psychosocial Determinants of Intrusive Recollections in Breast Cancer Survivors. Psychosomatics, 2005, 46, 203-211.	2.5	13
85	Are patients after severe injury who drop out of a longitudinal study at high risk of mental disorder?. Comprehensive Psychiatry, 2008, 49, 393-398.	1.5	13
86	Omegaâ€3 fatty acid supplementation for expectant mothers with depressive symptoms in Japan and Taiwan: An open″abel trial. Psychiatry and Clinical Neurosciences, 2016, 70, 253-254.	1.0	13
87	Change in blood levels of eicosapentaenoic acid and posttraumatic stress symptom: A secondary analysis of data from a placebo-controlled trial of omega3 supplements. Journal of Affective Disorders, 2016, 205, 289-291.	2.0	13
88	High-intensity interval training in breast cancer survivors: a systematic review. BMC Cancer, 2021, 21, 184.	1.1	13
89	Serum neuropeptide Y in accident survivors with depression or posttraumatic stress disorder. Neuroscience Research, 2014, 83, 8-12.	1.0	12
90	A multi-national, multi-disciplinary Delphi consensus study on using omega-3 polyunsaturated fatty acids (n-3 PUFAs) for the treatment of major depressive disorder. Journal of Affective Disorders, 2020, 265, 233-238.	2.0	12

#	Article	IF	CITATIONS
91	A nationally representative crossâ€sectional survey on health information access for consumers in Japan: A protocol for the INFORM Study. World Medical and Health Policy, 2022, 14, 225-275.	0.9	11
92	The synchronized trial on expectant mothers with depressive symptoms by omega-3 PUFAs (SYNCHRO): Study protocol for a randomized controlled trial. BMC Psychiatry, 2016, 16, 321.	1.1	10
93	Mechanisms underlying the effects of n-3 polyunsaturated fatty acids on fear memory processing and their hypothetical effects on fear of cancer recurrence in cancer survivors. Prostaglandins Leukotrienes and Essential Fatty Acids, 2018, 131, 14-23.	1.0	10
94	Naturally absorbed polyunsaturated fatty acids, lithium, and suicide-related behaviors: A case-controlled study. Journal of Affective Disorders, 2018, 241, 200-205.	2.0	10
95	Effect of home-based high-intensity interval training and behavioural modification using information and communication technology on cardiorespiratory fitness and exercise habits among sedentary breast cancer survivors: habit-B study protocol for a randomised controlled trial. BMJ Open, 2019, 9, e030911.	0.8	10
96	Evaluating a screener to quantify PTSD risk using emergency care information: a proof of concept study. BMC Emergency Medicine, 2020, 20, 16.	0.7	10
97	Towards an Explanation of Inconsistent Rates of Posttraumatic Stress Disorder across Different Countries: Infant Mortality Rate as a Marker of Social Circumstances and Basic Population Health. Psychotherapy and Psychosomatics, 2010, 79, 56-57.	4.0	9
98	Association between frequency of fried food consumption and resilience to depression in Japanese company workers: a cross-sectional study. Lipids in Health and Disease, 2016, 15, 156.	1.2	9
99	Developing the structure of Japan's cancer survivorship guidelines using an expert panel and modified Delphi method. Journal of Cancer Survivorship, 2020, 14, 273-283.	1.5	9
100	Yokukansan for Treatment of Preoperative Anxiety and Prevention of Postoperative Delirium in Cancer Patients Undergoing Highly Invasive Surgery. J-SUPPORT 1605 (ProD Study): A Randomized, Double-Blind, Placebo-Controlled Trial. Journal of Pain and Symptom Management, 2021, 61, 71-80.	0.6	9
101	Impact of psychiatric morbidity on quality of life after motor vehicle accident at 1â€month follow up. Psychiatry and Clinical Neurosciences, 2009, 63, 235-237.	1.0	8
102	Long-term influence of adjuvant breast radiotherapy on cognitive function in breast cancer patients treated with conservation therapy. International Journal of Clinical Oncology, 2019, 24, 68-77.	1.0	8
103	Left Hippocampal Volume Inversely Correlates With Enhanced Emotional Memory in Healthy Middle-Aged Women. Journal of Neuropsychiatry and Clinical Neurosciences, 2007, 19, 335-338.	0.9	7
104	The role of high-density lipoprotein cholesterol in risk for posttraumatic stress disorder: Taking a nutritional approach toward universal prevention. European Psychiatry, 2014, 29, 408-413.	0.1	7
105	Association between blood polyunsaturated fatty acid levels and depressive symptoms in breast cancer survivors. Prostaglandins Leukotrienes and Essential Fatty Acids, 2018, 139, 9-13.	1.0	7
106	Differential Effect of Two Mental Stress Tasks on Arterial Stiffness. Japanese Psychological Research, 2019, 61, 249-261.	0.4	7
107	Relationship between heart rate and emotional memory in subjects with a past history of post-traumatic stress disorder. Psychiatry and Clinical Neurosciences, 2007, 61, 441-443.	1.0	6
108	The Effect of Omega-3 Fatty Acids on Psychophysiological Assessment for the Secondary Prevention of Posttraumatic Stress Disorder: An Open-Label Pilot Study. Global Journal of Health Science, 2011, 4, 3-9.	0.1	6

#	Article	IF	CITATIONS
109	Serum polyunsaturated fatty acids and risk of psychiatric disorder after acute coronary syndrome: A prospective cohort study. Journal of Affective Disorders, 2017, 218, 306-312.	2.0	6
110	Midlife cancer/diabetes and risk of dementia and mild cognitive impairment: A populationâ€based prospective cohort study in Japan. Psychiatry and Clinical Neurosciences, 2019, 73, 597-599.	1.0	6
111	Fear of cancer recurrence among breast cancer survivors could be controlled by prudent dietary modification with polyunsaturated fatty acids. Journal of Affective Disorders, 2019, 245, 1114-1118.	2.0	6
112	Death by suicide, other externally caused injuries and cardiovascular diseases within 6Âmonths of cancer diagnosis (J-SUPPORT 1902). Japanese Journal of Clinical Oncology, 2021, 51, 744-752.	0.6	6
113	Comorbid insomnia among breast cancer survivors and its prediction using machine learning: a nationwide study in Japan. Japanese Journal of Clinical Oncology, 2021, , .	0.6	6
114	Lack of association between suicidal ideation and enhanced platelet 5-HT2A receptor-mediated calcium mobilization in cancer patients with depression. Biological Psychiatry, 2002, 52, 1159-1165.	0.7	5
115	Effect of cortisol diurnal rhythm on emotional memory in healthy young adults. Scientific Reports, 2017, 7, 10158.	1.6	5
116	Changes in depressive symptoms among family caregivers of patients with cancer after bereavement and their association with resilience: A prospective cohort study. Psycho-Oncology, 2021, , .	1.0	5
117	The Safety of Fish Oils for Those Whose Risk of Injury is High. Military Medicine, 2014, 179, 134-137.	0.4	4
118	Serum polyunsaturated fatty acids and risk of psychiatric disorder at 6 months after acute coronary syndrome: A prospective cohort study. Prostaglandins Leukotrienes and Essential Fatty Acids, 2019, 149, 18-23.	1.0	4
119	Differences between Japan and Taiwan in the treatment of pregnant women with depressive symptoms by omega-3 fatty acids: An open-label pilot study. Nutritional Neuroscience, 2019, 22, 63-71.	1.5	4
120	Study protocol for a nationwide questionnaire survey of physical activity among breast cancer survivors in Japan. BMJ Open, 2020, 10, e032871.	0.8	4
121	Different emotional memory consolidation in cancer survivors with and those without a history of intrusive recollection. Journal of Traumatic Stress, 2007, 20, 727-736.	1.0	3
122	Peritraumatic distress after an earthquake: a bridge between neuroimaging and epidemiology. Molecular Psychiatry, 2013, 18, 743-744.	4.1	3
123	Plasma phospholipid n-3 polyunsaturated fatty acids and major depressive disorder in Japanese elderly: the Japan Public Health Center-based Prospective Study. Scientific Reports, 2021, 11, 4003.	1.6	3
124	Intrusion in Women with Breast Cancer. , 2006, , 169-178.		3
125	Polyunsaturated Fatty Acids, Exercise, and Cancer-Related Fatigue in Breast Cancer Survivors. Frontiers in Physiology, 2021, 12, 759280.	1.3	3
126	Generalizability of Findings from a Randomized Controlled Trial of Fish Oil Supplementation for Attenuating Posttraumatic Stress Symptoms among Rescue Workers in Japan. Psychotherapy and Psychosomatics, 2018, 87, 114-115.	4.0	2

#	Article	IF	CITATIONS
127	Suicide, other externally caused injuries and cardiovascular death following a cancer diagnosis: study protocol for a nationwide population-based study in Japan (J-SUPPORT 1902). BMJ Open, 2019, 9, e030681.	0.8	2
128	The Association Between Postoperative Recovery and Psychosocial Factors in Cancer Patients. Journal of Surgery (New York, NY), 2020, 8, 9.	0.1	2
129	Oncology care providers' awareness and practice related to physical activity promotion for breast cancer survivors and barriers and facilitators to such promotion: a nationwide cross-sectional web-based survey. Supportive Care in Cancer, 2022, 30, 3105-3118.	1.0	2
130	Cavum septi pellucidi and intrusive recollections in cancer survivors. Neuroscience Research, 2006, 56, 344-346.	1.0	1
131	Yokukansan for perioperative psychiatric symptoms in cancer patients undergoing high invasive surgery. J-SUPPORT 1605 (ProD Study): study protocol for a randomized controlled trial. Trials, 2019, 20, 110.	0.7	1
132	Potential beneficial effects of Bifidobacterium breve A1 on cognitive impairment and psychiatric disorders., 2021,, 497-504.		1
133	Impact of peritraumatic distress on posttraumatic stress disorder symptoms at 6 months after acute coronary syndrome: a prospective cohort study. Högre Utbildning, 2021, 12, 1854511.	1.4	1
134	Application of the Unified Protocol for a Japanese Patient with Post-Traumatic Stress Disorder and Multiple Comorbidities: A Single-Case Study. International Journal of Environmental Research and Public Health, 2021, 18, 11644.	1.2	1
135	Accuracy of exercise-based tests for estimating cardiorespiratory fitness and muscle strength in early-stage breast cancer survivors in Japan. Supportive Care in Cancer, 2022, 30, 3857.	1.0	1
136	Association between cortisol diurnal rhythm and emotional memory in healthy young volunteers. Psychoneuroendocrinology, 2017, 83, 68.	1.3	0
137	The Association Between Fear of Cancer Recurrence and the Gut Microbiota in Breast Cancer Survivors (P05-030-19). Current Developments in Nutrition, 2019, 3, nzz030.P05-030-19.	0.1	0
138	Role of resilience for the association between trait hostility and depressive symptoms in Japanese company workers. Current Psychology, 2021, 40, 2301-2308.	1.7	0
139	Perilla Oil and Bifidobacterium for Alleviating Fear of Cancer Recurrence in Breast Cancer Survivors: Study Protocol for a Three-Arm Phase II Randomized Controlled Study (POB Study). Methods and Protocols, 2021, 4, 46.	0.9	0
140	Attitude to suicide prevention and suicide intervention skills among oncology professionals: An online crossâ€sectional survey in Japan. Psychiatry and Clinical Neurosciences, 2021, 75, 401-402.	1.0	0
141	Pharmacologic Interventions for Kawasaki Disease in Children: A Network Meta-Analysis of 56 Randomized Controlled Trials. SSRN Electronic Journal, 0, , .	0.4	0
142	Abstract OT3-12-01: Effect of home-based high-intensity interval training and behavioral modification using information and communication technology on cardiorespiratory fitness and exercise habits among sedentary breast cancer survivors: The habit-B randomized controlled trial in progress. , 2020,		0
143	BPI20-018: Direction of Developing Cancer Survivorship Guidelines in Japan: An Expert Panel's Consensus. Journal of the National Comprehensive Cancer Network: JNCCN, 2020, 18, BPI20-018.	2.3	0