Yutaka J Matsuoka

List of Publications by Citations

Source: https://exaly.com/author-pdf/6554329/yutaka-j-matsuoka-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

136 papers

3,260 citations

33 h-index 52 g-index

152 ext. papers

3,888 ext. citations

5.1 avg, IF

5.16 L-index

#	Paper	IF	Citations
136	Nutritional medicine as mainstream in psychiatry. <i>Lancet Psychiatry,the</i> , 2015 , 2, 271-4	23.3	268
135	Smaller regional volumes of brain gray and white matter demonstrated in breast cancer survivors exposed to adjuvant chemotherapy. <i>Cancer</i> , 2007 , 109, 146-56	6.4	248
134	A systematic review of predictors of posttraumatic stress disorder (PTSD) for adult road traffic crash survivors. <i>Injury</i> , 2013 , 44, 1413-22	2.5	118
133	Reliability and validity of the Japanese version of the Resilience Scale and its short version. <i>BMC Research Notes</i> , 2010 , 3, 310	2.3	88
132	Estimating the risk of PTSD in recent trauma survivors: results of the International Consortium to Predict PTSD (ICPP). <i>World Psychiatry</i> , 2019 , 18, 77-87	14.4	87
131	Posttraumatic growth, posttraumatic stress disorder and resilience of motor vehicle accident survivors. <i>BioPsychoSocial Medicine</i> , 2010 , 4, 7	2.8	83
130	The Association Between Fear of Cancer Recurrence and the Gut Microbiota in Breast Cancer Survivors (P05-030-19). <i>Current Developments in Nutrition</i> , 2019 , 3,	0.4	78
129	A volumetric study of amygdala in cancer survivors with intrusive recollections. <i>Biological Psychiatry</i> , 2003 , 54, 736-43	7.9	71
128	International Society for Nutritional Psychiatry Research Practice Guidelines for Omega-3 Fatty Acids in the Treatment of Major Depressive Disorder. <i>Psychotherapy and Psychosomatics</i> , 2019 , 88, 263-	2 973	66
127	Omega-3 Polyunsaturated Fatty Acids in Prevention of Mood and Anxiety Disorders. <i>Clinical Psychopharmacology and Neuroscience</i> , 2015 , 13, 129-37	3.4	65
126	Relationship between distressing cancer-related recollections and hippocampal volume in cancer survivors. <i>American Journal of Psychiatry</i> , 2002 , 159, 2087-93	11.9	63
125	Structure of orbitofrontal cortex and its longitudinal course in cancer-related post-traumatic stress disorder. <i>Neuroscience Research</i> , 2007 , 59, 383-9	2.9	62
124	Effect of bifidobacterium breve A-1 on anxiety and depressive symptoms in schizophrenia: A proof-of-concept study. <i>Journal of Affective Disorders</i> , 2019 , 245, 377-385	6.6	60
123	Association of Delirium Response and Safety of Pharmacological Interventions for the Management and Prevention of Delirium: A Network Meta-analysis. <i>JAMA Psychiatry</i> , 2019 , 76, 526-535	14.5	59
122	International Society for Nutritional Psychiatry Research consensus position statement: nutritional medicine in modern psychiatry. <i>World Psychiatry</i> , 2015 , 14, 370-1	14.4	55
121	Peritraumatic Distress Inventory as a predictor of post-traumatic stress disorder after a severe motor vehicle accident. <i>Psychiatry and Clinical Neurosciences</i> , 2010 , 64, 149-56	6.2	53
120	Peritraumatic distress, watching television, and posttraumatic stress symptoms among rescue workers after the Great East Japan earthquake. <i>PLoS ONE</i> , 2012 , 7, e35248	3.7	53

(2011-2014)

119	Association between adjuvant regional radiotherapy and cognitive function in breast cancer patients treated with conservation therapy. <i>Cancer Medicine</i> , 2014 , 3, 702-9	4.8	48
118	No adverse effects of adjuvant chemotherapy on hippocampal volume in Japanese breast cancer survivors. <i>Breast Cancer Research and Treatment</i> , 2005 , 92, 81-4	4.4	46
117	Cancer-related intrusive thoughts as an indicator of poor psychological adjustment at 3 or more years after breast surgery: a preliminary study. <i>Breast Cancer Research and Treatment</i> , 2002 , 76, 117-24	4.4	45
116	Concern over radiation exposure and psychological distress among rescue workers following the Great East Japan Earthquake. <i>BMC Public Health</i> , 2012 , 12, 249	4.1	43
115	Association of Use of Omega-3 Polyunsaturated Fatty Acids With Changes in Severity of Anxiety Symptoms: A Systematic Review and Meta-analysis. <i>JAMA Network Open</i> , 2018 , 1, e182327	10.4	43
114	Associations between disaster exposures, peritraumatic distress, and posttraumatic stress responses in Fukushima nuclear plant workers following the 2011 nuclear accident: the Fukushima NEWS Project study. <i>PLoS ONE</i> , 2014 , 9, e87516	3.7	42
113	Occurrence of fatigue and associated factors in disease-free breast cancer patients without depression. <i>Supportive Care in Cancer</i> , 2005 , 13, 628-36	3.9	41
112	Dietary fish, n-3 polyunsaturated fatty acid consumption, and depression risk in Japan: a population-based prospective cohort study. <i>Translational Psychiatry</i> , 2017 , 7, e1242	8.6	40
111	Possibility of polyunsaturated fatty acids for the prevention and treatment of neuropsychiatric illnesses. <i>Journal of Pharmacological Sciences</i> , 2014 , 124, 294-300	3.7	39
110	Clinical utility and validation of the Japanese version of Memorial Delirium Assessment Scale in a psychogeriatric inpatient setting. <i>General Hospital Psychiatry</i> , 2001 , 23, 36-40	5.6	39
109	Relationship of psychological characteristics and self-efficacy in gastrointestinal cancer survivors. <i>Psycho-Oncology</i> , 2010 , 19, 71-6	3.9	37
108	Omega-3 fatty acids for secondary prevention of posttraumatic stress disorder after accidental injury: an open-label pilot study. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 217-9	1.7	36
107	Incidence and prediction of psychiatric morbidity after a motor vehicle accident in Japan: the Tachikawa Cohort of Motor Vehicle Accident Study. <i>Critical Care Medicine</i> , 2008 , 36, 74-80	1.4	36
106	Prefrontal cortex and amygdala volume in first minor or major depressive episode after cancer diagnosis. <i>Biological Psychiatry</i> , 2006 , 59, 707-12	7.9	35
105	Resilience, post-traumatic growth, and work engagement among health care professionals after the Great East Japan Earthquake: A 4-year prospective follow-up study. <i>Journal of Occupational Health</i> , 2016 , 58, 347-53	2.3	34
104	Fish oil for attenuating posttraumatic stress symptoms among rescue workers after the great east Japan earthquake: a randomized controlled trial. <i>Psychotherapy and Psychosomatics</i> , 2012 , 81, 315-7	9.4	33
103	Hippocampal volume and first major depressive episode after cancer diagnosis in breast cancer survivors. <i>American Journal of Psychiatry</i> , 2004 , 161, 2263-70	11.9	33
102	Potential role of brain-derived neurotrophic factor in omega-3 Fatty Acid supplementation to prevent posttraumatic distress after accidental injury: an open-label pilot study. <i>Psychotherapy and Psychosomatics</i> , 2011 , 80, 310-2	9.4	32

101	Reliability and validity of the Japanese version of the Peritraumatic Distress Inventory. <i>General Hospital Psychiatry</i> , 2009 , 31, 75-9	5.6	32
100	Longitudinal changes in serum brain-derived neurotrophic factor in accident survivors with posttraumatic stress disorder. <i>Neuropsychobiology</i> , 2013 , 68, 44-50	4	29
99	Culturally sensitive and universal measure of resilience for Japanese populations: Tachikawa Resilience Scale in comparison with Resilience Scale 14-item version. <i>Psychiatry and Clinical Neurosciences</i> , 2013 , 67, 174-81	6.2	28
98	Docosahexaenoic acid for selective prevention of posttraumatic stress disorder among severely injured patients: a randomized, placebo-controlled trial. <i>Journal of Clinical Psychiatry</i> , 2015 , 76, e1015-	22 ^{4.6}	28
97	Clearance of fear memory from the hippocampus through neurogenesis by omega-3 fatty acids: a novel preventive strategy for posttraumatic stress disorder?. <i>BioPsychoSocial Medicine</i> , 2011 , 5, 3	2.8	26
96	Effects of omega-3 polyunsaturated fatty acids on psychophysiological symptoms of post-traumatic stress disorder in accident survivors: A randomized, double-blind, placebo-controlled trial. <i>Journal of Affective Disorders</i> , 2017 , 224, 27-31	6.6	25
95	Regional cerebral glucose metabolism in patients with secondary depressive episodes after fatal pancreatic cancer diagnosis. <i>Journal of Affective Disorders</i> , 2007 , 99, 231-6	6.6	25
94	Differentiating PTSD from anxiety and depression: Lessons from the ICD-11 PTSD diagnostic criteria. <i>Depression and Anxiety</i> , 2019 , 36, 490-498	8.4	24
93	Serum pro-BDNF/BDNF as a treatment biomarker for response to docosahexaenoic acid in traumatized people vulnerable to developing psychological distress: a randomized controlled trial. <i>Translational Psychiatry</i> , 2015 , 5, e596	8.6	24
92	The Tachikawa cohort of motor vehicle accident study investigating psychological distress: design, methods and cohort profiles. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009 , 44, 333-40	4.5	24
91	Association between regular physical exercise and depressive symptoms mediated through social support and resilience in Japanese company workers: a cross-sectional study. <i>BMC Public Health</i> , 2016 , 16, 553	4.1	23
90	Probability of major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic interviews controlling for Hospital Anxiety and Depression Scale - Depression subscale scores: An individual participant data meta-analysis of 73 primary studies. <i>Journal of Psychosomatic</i>	4.1	22
89	Clinical implications of the proposed ICD-11 PTSD diagnostic criteria. <i>Psychological Medicine</i> , 2019 , 49, 483-490	6.9	21
88	Secondary traumatic stress and associated factors among Japanese nurses working in hospitals. <i>International Journal of Nursing Practice</i> , 2012 , 18, 155-63	2.3	21
87	A randomized controlled multicenter trial of post-suicide attempt case management for the prevention of further attempts in Japan (ACTION-J). <i>BMC Public Health</i> , 2009 , 9, 364	4.1	21
86	Preoperative Anxiety as a Predictor of Delirium in Cancer Patients: A Prospective Observational Cohort Study. <i>World Journal of Surgery</i> , 2019 , 43, 134-142	3.3	20
85	A mindfulness-based stress management program and treatment with omega-3 fatty acids to maintain a healthy mental state in hospital nurses (Happy Nurse Project): study protocol for a randomized controlled trial. <i>Trials</i> , 2015 , 16, 36	2.8	19
84	Serum oxytocin, posttraumatic coping and C-reactive protein in motor vehicle accident survivors by gender. <i>Neuropsychobiology</i> , 2015 , 71, 196-201	4	19

(2020-2018)

83	Effectiveness of assertive case management on repeat self-harm in patients admitted for suicide attempt: Findings from ACTION-J study. <i>Journal of Affective Disorders</i> , 2018 , 225, 460-465	6.6	19	
82	Glutamatergic system abnormalities in posttraumatic stress disorder. <i>Psychopharmacology</i> , 2015 , 232, 4261-8	4.7	18	
81	Fish consumption and resilience to depression in Japanese company workers: a cross-sectional study. <i>Lipids in Health and Disease</i> , 2015 , 14, 51	4.4	17	
80	Posttraumatic stress disorder symptom trajectories within the first year following emergency department admissions: pooled results from the International Consortium to predict PTSD. <i>Psychological Medicine</i> , 2021 , 51, 1129-1139	6.9	17	
79	Brief mindfulness-based stress management program for a better mental state in working populations - Happy Nurse Project: A randomized controlled trial. <i>Journal of Affective Disorders</i> , 2019 , 251, 186-194	6.6	16	
78	Attenuating posttraumatic distress with omega-3 polyunsaturated fatty acids among disaster medical assistance team members after the Great East Japan Earthquake: the APOP randomized controlled trial. <i>BMC Psychiatry</i> , 2011 , 11, 132	4.2	15	
77	Limited effect of omega-3 fatty acids on the quality of life in survivors of traumatic injury: A randomized, placebo-controlled trial. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2017 , 127, 1-5	2.8	14	
76	The longitudinal mental health impact of Fukushima nuclear disaster exposures and public criticism among power plant workers: the Fukushima NEWS Project study. <i>Psychological Medicine</i> , 2016 , 46, 3117	7-3725	14	
75	The Efficacy of Omega-3 Fatty Acids for Depressive Symptoms among Pregnant Women in Japan and Taiwan: A Randomized, Double-Blind, Placebo-Controlled Trial (SYNCHRO; NCT01948596). <i>Psychotherapy and Psychosomatics</i> , 2019 , 88, 122-124	9.4	14	
74	Plasma estradiol levels and antidepressant effects of omega-3 fatty acids in pregnant women. Brain, Behavior, and Immunity, 2020 , 85, 29-34	16.6	14	
73	Fish consumption and cardiovascular response during mental stress. <i>BMC Research Notes</i> , 2012 , 5, 288	2.3	13	
72	Biomedical and psychosocial determinants of intrusive recollections in breast cancer survivors. <i>Psychosomatics</i> , 2005 , 46, 203-11	2.6	13	
71	Impact of chemotherapy on the association between fear of cancer recurrence and the gut microbiota in breast cancer survivors. <i>Brain, Behavior, and Immunity,</i> 2020 , 85, 186-191	16.6	13	
70	Tachikawa project for prevention of posttraumatic stress disorder with polyunsaturated fatty acid (TPOP): study protocol for a randomized controlled trial. <i>BMC Psychiatry</i> , 2013 , 13, 8	4.2	12	
69	Serum neuropeptide Y in accident survivors with depression or posttraumatic stress disorder. Neuroscience Research, 2014 , 83, 8-12	2.9	12	
68	Potential impact of propofol immediately after motor vehicle accident on later symptoms of posttraumatic stress disorder at 6-month follow up: a retrospective cohort study. <i>Critical Care</i> , 2012 , 16, R196	10.8	12	
67	Hippocampal and amygdalar volumes in breast cancer survivors with posttraumatic stress disorder. Journal of Neuropsychiatry and Clinical Neurosciences, 2008 , 20, 302-8	2.7	12	
66	Data Validation and Verification Using Blockchain in a Clinical Trial for Breast Cancer: Regulatory Sandbox. <i>Journal of Medical Internet Research</i> , 2020 , 22, e18938	7.6	12	

65	Application of data pooling to longitudinal studies of early post-traumatic stress disorder (PTSD): the International Consortium to Predict PTSD (ICPP) project. <i>Hgre Utbildning</i> , 2018 , 9, 1476442	5	12
64	Prevalence and related factors of common mental disorders during pregnancy in Japan: a cross-sectional study. <i>BioPsychoSocial Medicine</i> , 2016 , 10, 17	2.8	11
63	Yokukansan for the treatment of preoperative anxiety and postoperative delirium in colorectal cancer patients: a retrospective study. <i>Japanese Journal of Clinical Oncology</i> , 2017 , 47, 844-848	2.8	10
62	Omega-3 fatty acids for a better mental state in working populations - Happy Nurse Project: A 52-week randomized controlled trial. <i>Journal of Psychiatric Research</i> , 2018 , 102, 72-80	5.2	10
61	Change in blood levels of eicosapentaenoic acid and posttraumatic stress symptom: A secondary analysis of data from a placebo-controlled trial of omega3 supplements. <i>Journal of Affective Disorders</i> , 2016 , 205, 289-291	6.6	10
60	Incidence and prediction of post-traumatic stress disorder at 6 months after motor vehicle accidents in Japan. <i>Psychosomatics</i> , 2013 , 54, 263-71	2.6	10
59	Are patients after severe injury who drop out of a longitudinal study at high risk of mental disorder?. <i>Comprehensive Psychiatry</i> , 2008 , 49, 393-8	7.3	10
58	Omega-3 fatty acid supplementation for expectant mothers with depressive symptoms in Japan and Taiwan: An open-label trial. <i>Psychiatry and Clinical Neurosciences</i> , 2016 , 70, 253-4	6.2	10
57	Diet quality and depression risk in a Japanese population: the Japan Public Health Center (JPHC)-based Prospective Study. <i>Scientific Reports</i> , 2019 , 9, 7150	4.9	9
56	Expectations of and recommendations for a cancer survivorship guideline in Japan: a literature review of guidelines for cancer survivorship. <i>Japanese Journal of Clinical Oncology</i> , 2019 , 49, 812-822	2.8	9
55	Mechanisms underlying the effects of n-3 polyunsaturated fatty acids on fear memory processing and their hypothetical effects on fear of cancer recurrence in cancer survivors. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2018 , 131, 14-23	2.8	9
54	Serum levels of polyunsaturated fatty acids and the risk of posttraumatic stress disorder. <i>Psychotherapy and Psychosomatics</i> , 2013 , 82, 408-10	9.4	9
53	Post-Traumatic Stress Symptoms and Burnout Among Medical Rescue Workers 4 Years After the Great East Japan Earthquake: A Longitudinal Study. <i>Disaster Medicine and Public Health Preparedness</i> , 2016 , 10, 848-853	2.8	9
52	A multi-national, multi-disciplinary Delphi consensus study on using omega-3 polyunsaturated fatty acids (n-3 PUFAs) for the treatment of major depressive disorder. <i>Journal of Affective Disorders</i> , 2020 , 265, 233-238	6.6	8
51	Pharmacologic and hormonal treatments for menopausal sleep disturbances: A network meta-analysis of 43 randomized controlled trials and 32,271 menopausal women. <i>Sleep Medicine Reviews</i> , 2021 , 57, 101469	10.2	8
50	Relationship between polyunsaturated fatty acid and eating disorders: Systematic review and meta-analysis. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2019 , 142, 11-19	2.8	8
49	Explicit prognostic disclosure to Asian women with breast cancer: A randomized, scripted video-vignette study (J-SUPPORT1601). <i>Cancer</i> , 2019 , 125, 3320-3329	6.4	7
48	The synchronized trial on expectant mothers with depressive symptoms by omega-3 PUFAs (SYNCHRO): Study protocol for a randomized controlled trial. <i>BMC Psychiatry</i> , 2016 , 16, 321	4.2	7

(2007-2010)

Towards an explanation of inconsistent rates of posttraumatic stress disorder across different countries: infant mortality rate as a marker of social circumstances and basic population health. <i>Psychotherapy and Psychosomatics</i> , 2010 , 79, 56-7	9.4	7	
Impact of psychiatric morbidity on quality of life after motor vehicle accident at 1-month follow up. <i>Psychiatry and Clinical Neurosciences</i> , 2009 , 63, 235-7	6.2	7	
The effect of omega-3 fatty acids on psychophysiological assessment for the secondary prevention of posttraumatic stress disorder: an open-label pilot study. <i>Global Journal of Health Science</i> , 2011 , 4, 3-9	1.3	6	
Left hippocampal volume inversely correlates with enhanced emotional memory in healthy middle-aged women. <i>Journal of Neuropsychiatry and Clinical Neurosciences</i> , 2007 , 19, 335-8	2.7	6	
Naturally absorbed polyunsaturated fatty acids, lithium, and suicide-related behaviors: A case-controlled study. <i>Journal of Affective Disorders</i> , 2018 , 241, 200-205	6.6	5	
The role of high-density lipoprotein cholesterol in risk for posttraumatic stress disorder: taking a nutritional approach toward universal prevention. <i>European Psychiatry</i> , 2014 , 29, 408-13	6	5	
Developing the structure of Japan's cancer survivorship guidelines using an expert panel and modified Delphi method. <i>Journal of Cancer Survivorship</i> , 2020 , 14, 273-283	5.1	5	
Effect of home-based high-intensity interval training and behavioural modification using information and communication technology on cardiorespiratory fitness and exercise habits among sedentary breast cancer survivors: habit-B study protocol for a randomised controlled trial. <i>BMJ</i>	3	5	
Differential Effect of Two Mental Stress Tasks on Arterial Stiffness. <i>Japanese Psychological Research</i> , 2019 , 61, 249-261	1.1	5	
Serum polyunsaturated fatty acids and risk of psychiatric disorder after acute coronary syndrome: A prospective cohort study. <i>Journal of Affective Disorders</i> , 2017 , 218, 306-312	6.6	4	
Evaluating a screener to quantify PTSD risk using emergency care information: a proof of concept study. <i>BMC Emergency Medicine</i> , 2020 , 20, 16	2.4	4	
Relationship between heart rate and emotional memory in subjects with a past history of post-traumatic stress disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2007 , 61, 441-3	6.2	4	
Association between frequency of fried food consumption and resilience to depression in Japanese company workers: a cross-sectional study. <i>Lipids in Health and Disease</i> , 2016 , 15, 156	4.4	4	
Cardiorespiratory fitness in breast cancer survivors: a randomised controlled trial of home-based smartphone supported high intensity interval training. <i>BMJ Supportive and Palliative Care</i> , 2021 ,	2.2	4	
Lipid and Energy Metabolism of the Gut Microbiota Is Associated with the Response to Probiotic Strain for Anxiety and Depressive Symptoms in Schizophrenia. <i>Journal of Personalized Medicine</i> , 2021 , 11,	3.6	4	
Midlife cancer/diabetes and risk of dementia and mild cognitive impairment: A population-based prospective cohort study in Japan. <i>Psychiatry and Clinical Neurosciences</i> , 2019 , 73, 597-599	6.2	3	
Effect of cortisol diurnal rhythm on emotional memory in healthy young adults. <i>Scientific Reports</i> , 2017 , 7, 10158	4.9	3	
Different emotional memory consolidation in cancer survivors with and those without a history of intrusive recollection. <i>Journal of Traumatic Stress</i> , 2007 , 20, 727-36	3.8	3	
	countries: infant mortality rate as a marker of social circumstances and basic population health. Psychotherapy and Psychosomatics, 2010, 79, 56-7 Impact of psychiatric morbidity on quality of life after motor vehicle accident at 1-month follow up. Psychiatry and Clinical Neurosciences, 2009, 63, 235-7 The effect of omega-3 fatty acids on psychophysiological assessment for the secondary prevention of posttraumatic stress disorder: an open-label pilot study. Global Journal of Health Science, 2011, 4, 3-9 Left hippocampal volume inversely correlates with enhanced emotional memory in healthy middle-aged women. Journal of Neuropsychiatry and Clinical Neurosciences, 2007, 19, 335-8 Naturally absorbed polyunsaturated fatty acids, lithium, and suicide-related behaviors: A case-controlled study. Journal of Affective Disorders, 2018, 241, 200-205 The role of high-density lipoprotein cholesterol in risk for posttraumatic stress disorder: taking a nutritional approach toward universal prevention. European Psychiatry, 2014, 29, 408-13 Developing the structure of Japan's cancer survivorship guidelines using an expert panel and modified Delphi method. Journal of Cancer Survivorship, 2020, 14, 273-283 Effect of home-based high-intensity interval training and behaviory fitness and exercise habits among sedentary brast cancer survivors: habit-B study protocol for a randomised controlled trial. BMJ Differential Effect of Two Mental Stress Tasks on Arterial Stiffness. Japanese Psychological Research, 2019, 61, 249-261 Serum polyunsaturated fatty acids and risk of psychiatric disorder after acute coronary syndrome: A prospective cohort study. Journal of Affective Disorders, 2017, 218, 306-312 Evaluating a screener to quantify PTSD risk using emergency care information: a proof of concept study. BMC Emergency Medicine, 2020, 20, 16 Relationship between heart rate and emotional memory in subjects with a past history of post-traumatic stress disorder. Psychiatry and Clinical Neurosciences, 2016, 15, 156 Cardiorespirator	countries: infant mortality rate as a marker of social circumstances and basic population health. <i>Psychotherapy and Psychosomatics</i> , 2010, 79, 56-7 Impact of psychiatric morbidity on quality of life after motor vehicle accident at 1-month follow up. <i>Psychiatry and Clinical Neurosciences</i> , 2009, 63, 235-7 The effect of omega-3 fatty acids on psychophysiological assessment for the secondary prevention of posttraumatic stress disorder: an open-label pilot study. <i>Global Journal of Health Science</i> , 2011, 4, 3-9 Left hippocampal volume inversely correlates with enhanced emotional memory in healthy middle-aged women. <i>Journal of Neuropsychiatry and Clinical Neurosciences</i> , 2007, 19, 335-8 Naturally absorbed polyunsaturated fatty acids, lithium, and suicide-related behaviors: A case-controlled study. <i>Journal of Affective Disorders</i> , 2018, 241, 200-205 Attractional approach toward universal prevention. <i>European Psychiatry</i> , 2014, 29, 408-13 Developing the structure of Japan's cancer survivorship guidelines using an expert panel and modified Delphi method. <i>Journal of Cancer Survivorship</i> , 2020, 14, 273-283 Effect of home-based high-intensity interval training and behavioural modification using information and communication technology on cardiorespiratory fitness and exercise habits among sedentary breast cancer survivors: habit-B study protocol for a randomised controlled trial. <i>BMJ</i> Differential Effect of Two Mental Stress Tasks on Arterial Stiffness. <i>Japanese Psychological Research</i> , 2019, 61, 249-261 Serum polyunsaturated fatty acids and risk of psychiatric disorder after acute coronary syndrome: A prospective cohort study. <i>Journal of Affective Disorders</i> , 2017, 218, 306-312 Evaluating a screener to quantify PTSD risk using emergency care information: a proof of concept study. <i>BMC Emergency Medicine</i> , 2020, 20, 16 Relationship between heart rate and emotional memory in subjects with a past history of post-traumatic stress disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2007, 61, 441-3 Associat	countries: Infant mortality rate as a marker of social circumstances and basic population health. Psychotherapy and Psychosomatics, 2010, 79, 56-7 Impact of psychiatric morbidity on quality of life after motor vehicle accident at 1-month follow up. Psychiatry and Clinical Neurosciences, 2009, 63, 235-7 The effect of omega-3 fatty acids on psychophysiological assessment for the secondary prevention of posttraumatic stress disorder: an open-label pilot study. Clobal Journal of Health Science, 2011, 4, 3-9 Left hippocampal volume inversely correlates with enhanced emotional memory in healthy middle-aged women. Journal of Neuropsychiatry and Clinical Neurosciences, 2007, 19, 335-8 Naturally absorbed polyunsaturated fatty acids, lithium, and suicide-related behaviors: A case-controlled study. Journal of Affective Disorders, 2018, 241, 200-205 The role of high-density lipoprotein cholesterol in risk for posttraumatic stress disorder: taking a nutritional approach toward universal prevention. European Psychiatry, 2014, 29, 408-13 Developing the structure of Japan's cancer survivorship guidelines using an expert panel and modified Delphi method. Journal of Cancer Survivorship, 2020, 14, 273-283 Developing the structure of Japan's cancer survivorship, 2020, 14, 273-283 Effect of home-based high-intensity interval training and behavioural modification using information and communication technology on cardiorespiratory fitness and exercise habits among sedentary breast cancer survivors: habit B study protocol for a randomised controlled trial. BMJ Differential Effect of Two Mental Stress Tasks on Arterial Stiffness. Japanese Psychological Research, 2019, 61, 249-261 Serum polyunsaturated fatty acids and risk of psychiatric disorder after acute coronary syndrome: A prospective cohort study. Journal of Affective Disorders, 2017, 218, 306-312 Evaluating a screener to quantify PTSD risk using emergency care information: a proof of concept study. BMC Emergency Medicine, 2020, 20, 16 Relationship between heart rate

29	Lack of association between suicidal ideation and enhanced platelet 5-HT2A receptor-mediated calcium mobilization in cancer patients with depression. <i>Biological Psychiatry</i> , 2002 , 52, 1159-65	7.9	3
28	Long-term influence of adjuvant breast radiotherapy on cognitive function in breast cancer patients treated with conservation therapy. <i>International Journal of Clinical Oncology</i> , 2019 , 24, 68-77	4.2	3
27	Differences between Japan and Taiwan in the treatment of pregnant women with depressive symptoms by omega-3 fatty acids: An open-label pilot study. <i>Nutritional Neuroscience</i> , 2019 , 22, 63-71	3.6	3
26	Yokukansan for Treatment of Preoperative Anxiety and Prevention of Postoperative Delirium in Cancer Patients Undergoing Highly Invasive Surgery. J-SUPPORT 1605 (ProD Study): A Randomized, Double-Blind, Placebo-Controlled Trial. <i>Journal of Pain and Symptom Management</i> , 2021 , 61, 71-80	4.8	3
25	High-intensity interval training in breast cancer survivors: a systematic review. <i>BMC Cancer</i> , 2021 , 21, 184	4.8	3
24	Association between blood polyunsaturated fatty acid levels and depressive symptoms in breast cancer survivors. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2018 , 139, 9-13	2.8	3
23	Serum polyunsaturated fatty acids and risk of psychiatric disorder at 6 months after acute coronary syndrome: A prospective cohort study. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2019 , 149, 18-23	2.8	2
22	Peritraumatic distress after an earthquake: a bridge between neuroimaging and epidemiology. <i>Molecular Psychiatry</i> , 2013 , 18, 743-4	15.1	2
21	The Association Between Postoperative Recovery and Psychosocial Factors in Cancer Patients. Journal of Surgery (New York, NY), 2020 , 8, 9	0.6	2
20	Fear of cancer recurrence among breast cancer survivors could be controlled by prudent dietary modification with polyunsaturated fatty acids. <i>Journal of Affective Disorders</i> , 2019 , 245, 1114-1118	6.6	2
19	Association Between Dietary Fish and PUFA Intake in Midlife and Dementia in Later Life: The JPHC Saku Mental Health Study. <i>Journal of Alzheimerjs Disease</i> , 2021 , 79, 1091-1104	4.3	2
18	Death by suicide, other externally caused injuries and cardiovascular diseases within 6 months of cancer diagnosis (J-SUPPORT 1902). <i>Japanese Journal of Clinical Oncology</i> , 2021 , 51, 744-752	2.8	2
17	Intrusion in Women with Breast Cancer 2006 , 169-178		2
16	Generalizability of Findings from a Randomized Controlled Trial of Fish Oil Supplementation for Attenuating Posttraumatic Stress Symptoms among Rescue Workers in Japan. <i>Psychotherapy and Psychosomatics</i> , 2018 , 87, 114-115	9.4	1
15	Suicide, other externally caused injuries and cardiovascular death following a cancer diagnosis: study protocol for a nationwide population-based study in Japan (J-SUPPORT 1902). <i>BMJ Open</i> , 2019 , 9, e030681	3	1
14	The safety of fish oils for those whose risk of injury is high. <i>Military Medicine</i> , 2014 , 179, 134-7	1.3	1
13	Cavum septi pellucidi and intrusive recollections in cancer survivors. <i>Neuroscience Research</i> , 2006 , 56, 344-6	2.9	1
12	Study protocol for a nationwide questionnaire survey of physical activity among breast cancer survivors in Japan. <i>BMJ Open</i> , 2020 , 10, e032871	3	1

LIST OF PUBLICATIONS

11	Accuracy of exercise-based tests for estimating cardiorespiratory fitness and muscle strength in early-stage breast cancer survivors in Japan <i>Supportive Care in Cancer</i> , 2022 , 30, 3857	3.9	О
10	Application of the Unified Protocol for a Japanese Patient with Post-Traumatic Stress Disorder and Multiple Comorbidities: A Single-Case Study. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	O
9	Oncology care providers' awareness and practice related to physical activity promotion for breast cancer survivors and barriers and facilitators to such promotion: a nationwide cross-sectional web-based survey. Supportive Care in Cancer, 2021, 1	3.9	0
8	Polyunsaturated Fatty Acids, Exercise, and Cancer-Related Fatigue in Breast Cancer Survivors. <i>Frontiers in Physiology</i> , 2021 , 12, 759280	4.6	0
7	Plasma phospholipid n-3 polyunsaturated fatty acids and major depressive disorder in Japanese elderly: the Japan Public Health Center-based Prospective Study. <i>Scientific Reports</i> , 2021 , 11, 4003	4.9	О
6	Yokukansan for perioperative psychiatric symptoms in cancer patients undergoing high invasive surgery. J-SUPPORT 1605 (ProD Study): study protocol for a randomized controlled trial. <i>Trials</i> , 2019 , 20, 110	2.8	
5	BPI20-018: Direction of Developing Cancer Survivorship Guidelines in Japan: An Expert Panel Consensus. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , 2020 , 18, BPI20-018	7.3	
4	Role of resilience for the association between trait hostility and depressive symptoms in Japanese company workers. <i>Current Psychology</i> , 2021 , 40, 2301-2308	1.4	
3	Potential beneficial effects of Bifidobacterium breve A1 on cognitive impairment and psychiatric disorders 2021 , 497-504		
2	Attitude to suicide prevention and suicide intervention skills among oncology professionals: An online cross-sectional survey in Japan. <i>Psychiatry and Clinical Neurosciences</i> , 2021 , 75, 401-402	6.2	
1	Impact of peritraumatic distress on posttraumatic stress disorder symptoms at 6 months after acute coronary syndrome: a prospective cohort study. Higre Utbildning, 2021, 12, 1854511	5	