

JÃ¼rgen Zeier

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6560376/publications.pdf>

Version: 2024-02-01

51
papers

6,226
citations

76326

40
h-index

182427

51
g-index

52
all docs

52
docs citations

52
times ranked

6443
citing authors

#	ARTICLE	IF	CITATIONS
1	Natural variation in temperature-modulated immunity uncovers transcription factor bHLH059 as a thermoresponsive regulator in <i>Arabidopsis thaliana</i> . <i>PLoS Genetics</i> , 2021, 17, e1009290.	3.5	23
2	UGT76B1, a promiscuous hub of small molecule-based immune signaling, glucosylates N-hydroxypipecolic acid, and balances plant immunity. <i>Plant Cell</i> , 2021, 33, 714-734.	6.6	47
3	The mobile SAR signal N-hydroxypipecolic acid induces NPR1-dependent transcriptional reprogramming and immune priming. <i>Plant Physiology</i> , 2021, 186, 1679-1705.	4.8	39
4	Metabolic regulation of systemic acquired resistance. <i>Current Opinion in Plant Biology</i> , 2021, 62, 102050.	7.1	69
5	Insect eggs trigger systemic acquired resistance against a fungal and an oomycete pathogen. <i>New Phytologist</i> , 2021, 232, 2491-2505.	7.3	9
6	Inducible biosynthesis and immune function of the systemic acquired resistance inducer N-hydroxypipecolic acid in monocotyledonous and dicotyledonous plants. <i>Journal of Experimental Botany</i> , 2020, 71, 6444-6459.	4.8	36
7	Putrescine elicits ROS-dependent activation of the salicylic acid pathway in <i>Arabidopsis thaliana</i> . <i>Plant, Cell and Environment</i> , 2020, 43, 2755-2768.	5.7	40
8	Root-specific camalexin biosynthesis controls the plant growth-promoting effects of multiple bacterial strains. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 15735-15744.	7.1	134
9	Fluctuating Light Interacts with Time of Day and Leaf Development Stage to Reprogram Gene Expression. <i>Plant Physiology</i> , 2019, 179, 1632-1657.	4.8	53
10	Nitrite and nitric oxide are important in the adjustment of primary metabolism during the hypersensitive response in tobacco. <i>Journal of Experimental Botany</i> , 2019, 70, 4571-4582.	4.8	10
11	N-hydroxypipecolic acid and salicylic acid: a metabolic duo for systemic acquired resistance. <i>Current Opinion in Plant Biology</i> , 2019, 50, 44-57.	7.1	107
12	A Role for Tocopherol Biosynthesis in <i>Arabidopsis</i> Basal Immunity to Bacterial Infection. <i>Plant Physiology</i> , 2019, 181, 1008-1028.	4.8	49
13	A critical role for <i>Arabidopsis</i> MILDEW RESISTANCE LOCUS O2 in systemic acquired resistance. <i>Plant Journal</i> , 2018, 94, 1064-1082.	5.7	28
14	Flavin Monooxygenase-Generated N-Hydroxypipecolic Acid Is a Critical Element of Plant Systemic Immunity. <i>Cell</i> , 2018, 173, 456-469.e16.	28.9	297
15	A MPK3/6-WRKY33-ALD1-Pipecolic Acid Regulatory Loop Contributes to Systemic Acquired Resistance. <i>Plant Cell</i> , 2018, 30, 2480-2494.	6.6	119
16	Chemical Activation of EDS1/PAD4 Signaling Leading to Pathogen Resistance in <i>Arabidopsis</i> . <i>Plant and Cell Physiology</i> , 2018, 59, 1592-1607.	3.1	31
17	Lysine metabolism to N-hydroxypipecolic acid: an integral immune-activating pathway in plants. <i>Plant Journal</i> , 2018, 96, 5-21.	5.7	88
18	Biochemical Principles and Functional Aspects of Pipecolic Acid Biosynthesis in Plant Immunity. <i>Plant Physiology</i> , 2017, 174, 124-153.	4.8	111

#	ARTICLE	IF	CITATIONS
19	<i>Botrytis cinerea</i> B05.10 promotes disease development in <i>Arabidopsis</i> by suppressing WRKY33-mediated host immunity. <i>Plant, Cell and Environment</i> , 2017, 40, 2189-2206.	5.7	60
20	Regulatory and Functional Aspects of Indolic Metabolism in Plant Systemic Acquired Resistance. <i>Molecular Plant</i> , 2016, 9, 662-681.	8.3	62
21	Pipecolic Acid Orchestrates Plant Systemic Acquired Resistance and Defense Priming via Salicylic Acid-Dependent and -Independent Pathways. <i>Plant Cell</i> , 2016, 28, 102-129.	6.6	246
22	Insect eggs induce a systemic acquired resistance in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2014, 80, 1085-1094.	5.7	73
23	Spatial H ₂ O ₂ Signaling Specificity: H ₂ O ₂ from Chloroplasts and Peroxisomes Modulates the Plant Transcriptome Differentially. <i>Molecular Plant</i> , 2014, 7, 1191-1210.	8.3	167
24	Long-distance communication and signal amplification in systemic acquired resistance. <i>Frontiers in Plant Science</i> , 2013, 4, 30.	3.6	268
25	New insights into the regulation of plant immunity by amino acid metabolic pathways. <i>Plant, Cell and Environment</i> , 2013, 36, 2085-2103.	5.7	296
26	Copper and herbivory lead to priming and synergism in phytohormones and plant volatiles in the absence of salicylate-jasmonate antagonism. <i>Plant Signaling and Behavior</i> , 2013, 8, e24264.	2.4	10
27	The form of nitrogen nutrition affects resistance against <i>Pseudomonas syringae</i> pv. <i>phaseolicola</i> in tobacco. <i>Journal of Experimental Botany</i> , 2013, 64, 553-568.	4.8	116
28	Reprogramming of plants during systemic acquired resistance. <i>Frontiers in Plant Science</i> , 2013, 4, 252.	3.6	100
29	Pipecolic Acid, an Endogenous Mediator of Defense Amplification and Priming, Is a Critical Regulator of Inducible Plant Immunity. <i>Plant Cell</i> , 2013, 24, 5123-5141.	6.6	525
30	Pipecolic acid enhances resistance to bacterial infection and primes salicylic acid and nicotine accumulation in tobacco. <i>Plant Signaling and Behavior</i> , 2013, 8, e26366.	2.4	68
31	Heavy metal stress can prime for herbivore-induced plant volatile emission. <i>Plant, Cell and Environment</i> , 2012, 35, 1287-1298.	5.7	47
32	Cytokinins Mediate Resistance against <i>Pseudomonas syringae</i> in Tobacco through Increased Antimicrobial Phytoalexin Synthesis Independent of Salicylic Acid Signaling. <i>Plant Physiology</i> , 2011, 157, 815-830.	4.8	178
33	A role for sitosterol to stigmasterol conversion in plant pathogen interactions. <i>Plant Journal</i> , 2010, 63, 254-268.	5.7	134
34	Post-Translational Derepression of Invertase Activity in Source Leaves via Down-Regulation of Invertase Inhibitor Expression Is Part of the Plant Defense Response. <i>Molecular Plant</i> , 2010, 3, 1037-1048.	8.3	105
35	Methyl Salicylate Production and Jasmonate Signaling Are Not Essential for Systemic Acquired Resistance in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2009, 21, 954-971.	6.6	208
36	Light Regulation and Daytime Dependency of Inducible Plant Defenses in <i>Arabidopsis</i> : Phytochrome Signaling Controls Systemic Acquired Resistance Rather Than Local Defense. <i>Plant Physiology</i> , 2008, 147, 790-801.	4.8	236

#	ARTICLE	IF	CITATIONS
37	<i>Pseudomonas syringae</i> Elicits Emission of the Terpenoid (E,E)-4,8,12-Trimethyl-1,3,7,11-Tridecatetraene in <i>Arabidopsis</i> Leaves Via Jasmonate Signaling and Expression of the Terpene Synthase TPS4. <i>Molecular Plant-Microbe Interactions</i> , 2008, 21, 1482-1497.	2.6	45
38	A Central Role of Abscisic Acid in Drought Stress Protection of <i>Agrobacterium</i> -Induced Tumors on <i>Arabidopsis</i> . <i>Plant Physiology</i> , 2007, 145, 853-862.	4.8	74
39	Expression of a nitric oxide degrading enzyme induces a senescence programme in <i>Arabidopsis</i> . <i>Plant, Cell and Environment</i> , 2007, 30, 39-52.	5.7	138
40	Bacterial non-host resistance: interactions of <i>Arabidopsis</i> with non-adapted <i>Pseudomonas syringae</i> strains. <i>Physiologia Plantarum</i> , 2007, 131, 448-461.	5.2	49
41	Pathogen-associated molecular pattern recognition rather than development of tissue necrosis contributes to bacterial induction of systemic acquired resistance in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2007, 50, 500-513.	5.7	347
42	Nitric oxide (NO) as an intermediate in the cryptogein-induced hypersensitive response - a critical re-evaluation. <i>Plant, Cell and Environment</i> , 2006, 29, 59-69.	5.7	62
43	The <i>Arabidopsis</i> Flavin-Dependent Monooxygenase FMO1 Is an Essential Component of Biologically Induced Systemic Acquired Resistance. <i>Plant Physiology</i> , 2006, 141, 1666-1675.	4.8	229
44	Flavo-haemoglobin HmpX from <i>Erwinia chrysanthemi</i> confers nitrosative stress tolerance and affects the plant hypersensitive reaction by intercepting nitric oxide produced by the host. <i>Plant Journal</i> , 2005, 43, 226-237.	5.7	67
45	Age-dependent variations of local and systemic defence responses in <i>Arabidopsis</i> leaves towards an avirulent strain of <i>Pseudomonas syringae</i> . <i>Physiological and Molecular Plant Pathology</i> , 2005, 66, 30-39.	2.5	36
46	<i>Arabidopsis</i> Nonsymbiotic Hemoglobin AHb1 Modulates Nitric Oxide Bioactivity. <i>Plant Cell</i> , 2004, 16, 2785-2794.	6.6	332
47	Genetic Elucidation of Nitric Oxide Signaling in Incompatible Plant-Pathogen Interactions. <i>Plant Physiology</i> , 2004, 136, 2875-2886.	4.8	165
48	Light conditions influence specific defence responses in incompatible plant-pathogen interactions: uncoupling systemic resistance from salicylic acid and PR-1 accumulation. <i>Planta</i> , 2004, 219, 673-83.	3.2	190
49	Chemical analysis and immunolocalisation of lignin and suberin in endodermal and hypodermal/rhizodermal cell walls of developing maize (<i>Zea mays</i> L.) primary roots. <i>Planta</i> , 1999, 209, 1-12.	3.2	112
50	Fourier transform infrared-spectroscopic characterisation of isolated endodermal cell walls from plant roots: chemical nature in relation to anatomical development. <i>Planta</i> , 1999, 209, 537-542.	3.2	62
51	Comparative investigation of primary and tertiary endodermal cell walls isolated from the roots of five monocotyledonous species: chemical composition in relation to fine structure. <i>Planta</i> , 1998, 206, 349-361.	3.2	125