

Elena V Filatova

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6560958/publications.pdf>

Version: 2024-02-01

18
papers

236
citations

1307366

7
h-index

1058333

14
g-index

18
all docs

18
docs citations

18
times ranked

345
citing authors

#	ARTICLE	IF	CITATIONS
1	miRNA expression is highly sensitive to a drug therapy in Parkinson's disease. <i>Parkinsonism and Related Disorders</i> , 2015, 21, 72-74.	1.1	58
2	Major Depression: One Brain, One Disease, One Set of Intertwined Processes. <i>Cells</i> , 2021, 10, 1283.	1.8	47
3	Involvement of Endocytosis and Alternative Splicing in the Formation of the Pathological Process in the Early Stages of Parkinson's Disease. <i>BioMed Research International</i> , 2014, 2014, 1-6.	0.9	33
4	Transcriptome Profile Changes in Mice with MPTP-Induced Early Stages of Parkinson's Disease. <i>Molecular Neurobiology</i> , 2017, 54, 6775-6784.	1.9	20
5	VCP expression decrease as a biomarker of preclinical and early clinical stages of Parkinson's disease. <i>Scientific Reports</i> , 2020, 10, 827.	1.6	15
6	GABA, Selank, and Olanzapine Affect the Expression of Genes Involved in GABAergic Neurotransmission in IMR-32 Cells. <i>Frontiers in Pharmacology</i> , 2017, 8, 89.	1.6	14
7	Potential Biomarkers of the Earliest Clinical Stages of Parkinson's Disease. <i>Parkinson's Disease</i> , 2015, 2015, 1-6.	0.6	11
8	Expression analysis of suppression of tumorigenicity 13 gene in patients with Parkinson's disease. <i>Neuroscience Letters</i> , 2010, 473, 257-259.	1.0	7
9	Targeted exome analysis of Russian patients with hypertrophic cardiomyopathy. <i>Molecular Genetics & Genomic Medicine</i> , 2021, 9, e1808.	0.6	7
10	Peptides semax and selank affect the behavior of rats with 6-OHDA induced PD-like parkinsonism. <i>Doklady Biological Sciences</i> , 2017, 474, 106-109.	0.2	6
11	Housekeeping Genes for Parkinson's Disease in Humans and Mice. <i>Cells</i> , 2021, 10, 2252.	1.8	6
12	Polymorphisms in the SNCA Gene: Association with the Risk of Development of the Sporadic Form of Parkinson's Disease and the Level of SNCA Gene Expression in Peripheral Blood of Patients from Russia. <i>Neuroscience and Medicine</i> , 2013, 04, 208-214.	0.2	5
13	Expression Analysis of Genes Involved in Transport Processes in Mice with MPTP-Induced Model of Parkinson's Disease. <i>Life</i> , 2022, 12, 751.	1.1	5
14	Transcriptome profiling of 6-OHDA model of Parkinson's disease. <i>Advances in Bioscience and Biotechnology (Print)</i> , 2013, 04, 28-35.	0.3	2
15	3.007 ANALYSIS OF GENE EXPRESSION IN PERIPHERAL BLOOD IN PATIENTS WITH PARKINSON'S DISEASE. <i>Parkinsonism and Related Disorders</i> , 2012, 18, S171.	1.1	0
16	Analysis of known point mutations and SNPs in genes responsible for monogenic Parkinson's disease in Russian patients. <i>Advances in Parkinson's Disease</i> , 2013, 02, 28-30.	0.2	0
17	Synthetic Peptides Affect the Expression of α -Synuclein and Gdnf Receptors in Rats with 6-OHDA-Induced PD-Like Parkinsonism. <i>World Journal of Neuroscience</i> , 2016, 06, 243-259.	0.1	0
18	Differences in Relative Levels of 88 microRNAs in Various Regions of the Normal Adult Human Brain. <i>MicroRNA (Sharjah, United Arab Emirates)</i> , 2017, 6, 125-135.	0.6	0