Jan Johansson

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/6567995/jan-johansson-publications-by-year.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

176
papers

8,154
citations

h-index

85
g-index

185
ext. papers

9,227
ext. citations

85
g-index

L-index

#	Paper	IF	Citations
176	A Novel Approach for the Production of Aggregation-Prone Proteins Using the Spidroin-Derived NT* Tag <i>Methods in Molecular Biology</i> , 2022 , 2406, 113-130	1.4	1
175	A "spindle and thread" mechanism unblocks p53 translation by modulating N-terminal disorder <i>Structure</i> , 2022 ,	5.2	1
174	The dimerization mechanism of the N-terminal domain of spider silk proteins is conserved despite extensive sequence divergence <i>Journal of Biological Chemistry</i> , 2022 , 101913	5.4	1
173	Artificial spider silk supports and guides neurite extension in vitro. FASEB Journal, 2021, 35, e21896	0.9	1
172	AA amyloid in human food chain is a possible biohazard. <i>Scientific Reports</i> , 2021 , 11, 21069	4.9	2
171	Tyrosine residues mediate supercontraction in biomimetic spider silk. <i>Communications Materials</i> , 2021 , 2,	6	6
170	Ablation of p75 signaling strengthens gamma-theta rhythm interaction and counteracts Allnduced degradation of neuronal dynamics in mouse hippocampus in vitro. <i>Translational Psychiatry</i> , 2021 , 11, 212	8.6	3
169	The synthesis and characterization of Bri2 BRICHOS coated magnetic particles and their application to protein fishing: Identification of novel binding proteins. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2021 , 198, 113996	3.5	0
168	In Vitro Study of Human Immune Responses to Hyaluronic Acid Hydrogels, Recombinant Spidroins and Human Neural Progenitor Cells of Relevance to Spinal Cord Injury Repair. <i>Cells</i> , 2021 , 10,	7.9	2
167	Citrullination Alters the Antibacterial and Anti-Inflammatory Functions of the Host Defense Peptide Canine Cathelicidin K9CATH In Vitro. <i>Journal of Immunology</i> , 2021 , 207, 974-984	5.3	
166	Expression of the human molecular chaperone domain Bri2 BRICHOS on a gram per liter scale with an E. coli fed-batch culture. <i>Microbial Cell Factories</i> , 2021 , 20, 150	6.4	2
165	Impact of synthetic surfactant CHF5633 with SP-B and SP-C analogues on lung function and inflammation in rabbit model of acute respiratory distress syndrome. <i>Physiological Reports</i> , 2021 , 9, e14	700	6
164	Native-like Flow Properties of an Artificial Spider Silk Dope. <i>ACS Biomaterials Science and Engineering</i> , 2021 , 7, 462-471	5.5	4
163	Smallest Secondary Nucleation Competent Alaggregates Probed by an ATP-Independent Molecular Chaperone Domain. <i>Biochemistry</i> , 2021 , 60, 678-688	3.2	3
162	High-yield production of a super-soluble miniature spidroin for biomimetic high-performance materials. <i>Materials Today</i> , 2021 ,	21.8	5
161	Bri2 BRICHOS chaperone rescues impaired fast-spiking interneuron behavior and neuronal network dynamics in an AD mouse model in vitro. <i>Neurobiology of Disease</i> , 2021 , 159, 105514	7.5	0
160	Ion mobility-mass spectrometry shows stepwise protein unfolding under alkaline conditions. <i>Chemical Communications</i> , 2021 , 57, 1450-1453	5.8	1

159	Doing What Spiders Cannot-A Road Map to Supreme Artificial Silk Fibers. ACS Nano, 2021, 15, 1952-195	9 16.7	16
158	Recombinant BRICHOS chaperone domains delivered to mouse brain parenchyma by focused ultrasound and microbubbles are internalized by hippocampal and cortical neurons. <i>Molecular and Cellular Neurosciences</i> , 2020 , 105, 103498	4.8	7
157	Structure-Function Relationship of Artificial Spider Silk Fibers Produced by Straining Flow Spinning. <i>Biomacromolecules</i> , 2020 , 21, 2116-2124	6.9	16
156	Tensile properties of synthetic pyriform spider silk fibers depend on the number of repetitive units as well as the presence of N- and C-terminal domains. <i>International Journal of Biological Macromolecules</i> , 2020 , 154, 765-772	7.9	11
155	Recombinant Bri3 BRICHOS domain is a molecular chaperone with effect against amyloid formation and non-fibrillar protein aggregation. <i>Scientific Reports</i> , 2020 , 10, 9817	4.9	2
154	High-yield Production of Amyloid-Peptide Enabled by a Customized Spider Silk Domain. <i>Scientific Reports</i> , 2020 , 10, 235	4.9	16
153	Augmentation of Bri2 molecular chaperone activity against amyloid-Ireduces neurotoxicity in mouse hippocampus in vitro. <i>Communications Biology</i> , 2020 , 3, 32	6.7	11
152	High intracellular stability of the spidroin N-terminal domain in spite of abundant amyloidogenic segments revealed by in-cell hydrogen/deuterium exchange mass spectrometry. <i>FEBS Journal</i> , 2020 , 287, 2823-2833	5.7	4
151	Modulation of Kv3.1/Kv3.2 promotes gamma oscillations by rescuing Allnduced desynchronization of fast-spiking interneuron firing in an AD mouse model in vitro. <i>Journal of Physiology</i> , 2020 , 598, 3711-3	3725	7
150	Functionalization of amyloid fibrils via the Bri2 BRICHOS domain. <i>Scientific Reports</i> , 2020 , 10, 21765	4.9	5
149	Blood-brain and blood-cerebrospinal fluid passage of BRICHOS domains from two molecular chaperones in mice. <i>Journal of Biological Chemistry</i> , 2019 , 294, 2606-2615	5.4	6
148	BRICHOS: a chaperone with different activities depending on quaternary structure and cellular location?. <i>Amyloid: the International Journal of Experimental and Clinical Investigation: the Official Journal of the International Society of Amyloidosis</i> , 2019 , 26, 152-153	2.7	4
147	Synthetic surfactant with a recombinant surfactant protein C analogue improves lung function and attenuates inflammation in a model of acute respiratory distress syndrome in adult rabbits. <i>Respiratory Research</i> , 2019 , 20, 245	7.3	10
146	Efficient delipidation of a recombinant lung surfactant lipopeptide analogue by liquid-gel chromatography. <i>PLoS ONE</i> , 2019 , 14, e0226072	3.7	Ο
145	Disulphide Bridges in Surfactant Protein B Analogues Affect Their Activity in Synthetic Surfactant Preparations. <i>Neonatology</i> , 2019 , 115, 134-141	4	0
144	Synthetic surfactants with SP-B and SP-C analogues to enable worldwide treatment of neonatal respiratory distress syndrome and other lung diseases. <i>Journal of Internal Medicine</i> , 2019 , 285, 165-186	10.8	27
143	BRICHOS domain of Bri2 inhibits islet amyloid polypeptide (IAPP) fibril formation and toxicity in human beta cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, E2752-E2761	11.5	28
142	Production and Properties of Triple Chimeric Spidroins. <i>Biomacromolecules</i> , 2018 , 19, 2825-2833	6.9	14

141	A spidroin-derived solubility tag enables controlled aggregation of a designed amyloid protein. <i>FEBS Journal</i> , 2018 , 285, 1873-1885	5.7	22
140	Mass Spectrometry Reveals the Direct Action of a Chemical Chaperone. <i>Journal of Physical Chemistry Letters</i> , 2018 , 9, 4082-4086	6.4	16
139	The Bri2 and Bri3 BRICHOS Domains Interact Differently with Aland Alzheimer Amyloid Plaques. Journal of Alzheimers Disease Reports, 2018 , 2, 27-39	3.3	11
138	Novel expression of a functional trimeric fragment of human SP-A with efficacy in neutralisation of RSV. <i>Immunobiology</i> , 2017 , 222, 111-118	3.4	16
137	Biomimetic spinning of artificial spider silk from a chimeric minispidroin. <i>Nature Chemical Biology</i> , 2017 , 13, 262-264	11.7	143
136	Mass spectrometry captures structural intermediates in protein fiber self-assembly. <i>Chemical Communications</i> , 2017 , 53, 3319-3322	5.8	12
135	BRICHOS - an anti-amyloid chaperone: evaluation of blood-brain barrier permeability of Bri2 BRICHOS. <i>Amyloid: the International Journal of Experimental and Clinical Investigation: the Official Journal of the International Society of Amyloidosis</i> , 2017 , 24, 7-8	2.7	О
134	Efficient protein production inspired by how spiders make silk. <i>Nature Communications</i> , 2017 , 8, 15504	17.4	48
133	Degree of Biomimicry of Artificial Spider Silk Spinning Assessed by NMR Spectroscopy. <i>Angewandte Chemie</i> , 2017 , 129, 12745-12749	3.6	3
132	Degree of Biomimicry of Artificial Spider Silk Spinning Assessed by NMR Spectroscopy. <i>Angewandte Chemie - International Edition</i> , 2017 , 56, 12571-12575	16.4	15
131	Systemic AA amyloidosis in the red fox (Vulpes vulpes). <i>Protein Science</i> , 2017 , 26, 2312-2318	6.3	2
130	Bri2 BRICHOS client specificity and chaperone activity are governed by assembly state. <i>Nature Communications</i> , 2017 , 8, 2081	17.4	35
129	Transthyretin and BRICHOS: The Paradox of Amyloidogenic Proteins with Anti-Amyloidogenic Activity for Alin the Central Nervous System. <i>Frontiers in Neuroscience</i> , 2017 , 11, 119	5.1	18
128	TGF- 1 I-induced EMT promotes targeted migration of breast cancer cells through the lymphatic system by the activation of CCR7/CCL21-mediated chemotaxis. <i>Oncogene</i> , 2016 , 35, 748-60	9.2	194
127	Dementia-related Bri2 BRICHOS is a versatile molecular chaperone that efficiently inhibits A 2 toxicity in Drosophila. <i>Biochemical Journal</i> , 2016 , 473, 3683-3704	3.8	28
126	BRICHOS binds to a designed amyloid-forming Eprotein and reduces proteasomal inhibition and aggresome formation. <i>Biochemical Journal</i> , 2016 , 473, 167-78	3.8	14
125	Phospholipid Composition in Synthetic Surfactants Is Important for Tidal Volumes and Alveolar Stability in Surfactant-Treated Preterm Newborn Rabbits. <i>Neonatology</i> , 2016 , 109, 177-85	4	11
124	Silk Spinning in Silkworms and Spiders. <i>International Journal of Molecular Sciences</i> , 2016 , 17,	6.3	78

123	Kinetic analysis reveals the diversity of microscopic mechanisms through which molecular chaperones suppress amyloid formation. <i>Nature Communications</i> , 2016 , 7, 10948	17.4	153
122	Amyloid: a multifaceted player in human health and disease. <i>Journal of Internal Medicine</i> , 2016 , 280, 13	6-18 0.8	2
121	Current and future treatment of amyloid diseases. <i>Journal of Internal Medicine</i> , 2016 , 280, 177-202	10.8	65
120	Transmissible amyloid. <i>Journal of Internal Medicine</i> , 2016 , 280, 153-63	10.8	16
119	Dissociation of a BRICHOS trimer into monomers leads to increased inhibitory effect on AII2 fibril formation. <i>Biochimica Et Biophysica Acta - Proteins and Proteomics</i> , 2015 , 1854, 835-43	4	15
118	Toward spinning artificial spider silk. <i>Nature Chemical Biology</i> , 2015 , 11, 309-15	11.7	210
117	Specific chaperones and regulatory domains in control of amyloid formation. <i>Journal of Biological Chemistry</i> , 2015 , 290, 26430-6	5.4	27
116	Folding and Intramembraneous BRICHOS Binding of the Prosurfactant Protein C Transmembrane Segment. <i>Journal of Biological Chemistry</i> , 2015 , 290, 17628-41	5.4	7
115	BRI2 ectodomain affects AB2 fibrillation and tau truncation in human neuroblastoma cells. <i>Cellular and Molecular Life Sciences</i> , 2015 , 72, 1599-611	10.3	9
114	Diversified Structural Basis of a Conserved Molecular Mechanism for pH-Dependent Dimerization in Spider Silk N-Terminal Domains. <i>ChemBioChem</i> , 2015 , 16, 1720-4	3.8	29
113	Carbonic anhydrase generates a pH gradient in Bombyx mori silk glands. <i>Insect Biochemistry and Molecular Biology</i> , 2015 , 65, 100-6	4.5	30
112	A molecular chaperone breaks the catalytic cycle that generates toxic Albligomers. <i>Nature Structural and Molecular Biology</i> , 2015 , 22, 207-213	17.6	268
111	Separate molecular determinants in amyloidogenic and antimicrobial peptides. <i>Journal of Molecular Biology</i> , 2014 , 426, 2159-66	6.5	16
110	Amyloid-Induced action potential desynchronization and degradation of hippocampal gamma oscillations is prevented by interference with peptide conformation change and aggregation. <i>Journal of Neuroscience</i> , 2014 , 34, 11416-25	6.6	61
109	Recombinant spider silk genetically functionalized with affinity domains. <i>Biomacromolecules</i> , 2014 , 15, 1696-706	6.9	45
108	Spider silk for xeno-free long-term self-renewal and differentiation of human pluripotent stem cells. <i>Biomaterials</i> , 2014 , 35, 8496-502	15.6	34
107	Sequential pH-driven dimerization and stabilization of the N-terminal domain enables rapid spider silk formation. <i>Nature Communications</i> , 2014 , 5, 3254	17.4	96
106	Evaluation of Functionalized Spider Silk Matrices: Choice of Cell Types and Controls are Important for Detecting Specific Effects. <i>Frontiers in Bioengineering and Biotechnology</i> , 2014 , 2, 50	5.8	6

105	C/EBPlexpression is an independent predictor of overall survival in breast cancer patients by MHCII/CD4-dependent mechanism of metastasis formation. <i>Oncogenesis</i> , 2014 , 3, e125	6.6	12
104	Carbonic anhydrase generates CO2 and H+ that drive spider silk formation via opposite effects on the terminal domains. <i>PLoS Biology</i> , 2014 , 12, e1001921	9.7	109
103	The chaperone domain BRICHOS prevents CNS toxicity of amyloid-[peptide in Drosophila melanogaster. <i>DMM Disease Models and Mechanisms</i> , 2014 , 7, 659-65	4.1	36
102	The BRICHOS domain, amyloid fibril formation, and their relationship. <i>Biochemistry</i> , 2013 , 52, 7523-31	3.2	51
101	MiR-155-mediated loss of C/EBPIshifts the TGF-Iresponse from growth inhibition to epithelial-mesenchymal transition, invasion and metastasis in breast cancer. <i>Oncogene</i> , 2013 , 32, 5614-	2 ² -2	123
100	Morphology and composition of the spider major ampullate gland and dragline silk. <i>Biomacromolecules</i> , 2013 , 14, 2945-52	6.9	37
99	Harnessing the Self-Assembling Properties of Proteins in Spider Silk and Lung Surfactant 2013 , 455-470	0	1
98	Invited review current progress and limitations of spider silk for biomedical applications. <i>Biopolymers</i> , 2012 , 97, 468-78	2.2	64
97	BRICHOS domains efficiently delay fibrillation of amyloid Epeptide. <i>Journal of Biological Chemistry</i> , 2012 , 287, 31608-17	5.4	103
96	Control of amyloid assembly by autoregulation. <i>Biochemical Journal</i> , 2012 , 447, 185-92	3.8	13
95	pH-dependent dimerization of spider silk N-terminal domain requires relocation of a wedged tryptophan side chain. <i>Journal of Molecular Biology</i> , 2012 , 422, 477-87	6.5	61
94	Alterations of the C-terminal end do not affect in vitro or in vivo activity of surfactant protein C analogs. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2012 , 1818, 27-32	3.8	6
93	High-resolution structure of a BRICHOS domain and its implications for anti-amyloid chaperone activity on lung surfactant protein C. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 2325-9	11.5	87
92	Full-length minor ampullate spidroin gene sequence. <i>PLoS ONE</i> , 2012 , 7, e52293	3.7	58
91	BRICHOS domain associated with lung fibrosis, dementia and cancera chaperone that prevents amyloid fibril formation?. <i>FEBS Journal</i> , 2011 , 278, 3893-904	5.7	51
90	Spider silk proteins: recent advances in recombinant production, structure-function relationships and biomedical applications. <i>Cellular and Molecular Life Sciences</i> , 2011 , 68, 169-84	10.3	144
89	Functionalisation of recombinant spider silk with conjugated polyelectrolytes. <i>Journal of Materials Chemistry</i> , 2011 , 21, 2909		18
88	Self-assembly of spider silk proteins is controlled by a pH-sensitive relay. <i>Nature</i> , 2010 , 465, 236-8	50.4	328

(2008-2010)

87	Surfactant protein B propeptide contains a saposin-like protein domain with antimicrobial activity at low pH. <i>Journal of Immunology</i> , 2010 , 184, 975-83	5.3	51
86	Different effects of surfactant proteins B and C - implications for development of synthetic surfactants. <i>Neonatology</i> , 2010 , 97, 367-72	4	27
85	A pH-dependent dimer lock in spider silk protein. <i>Journal of Molecular Biology</i> , 2010 , 404, 328-36	6.5	47
84	The extracellular domain of Bri2 (ITM2B) binds the ABri peptide (1-23) and amyloid beta-peptide (Abeta1-40): Implications for Bri2 effects on processing of amyloid precursor protein and Abeta aggregation. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 393, 356-61	3.4	40
83	Amino acid sequence determinants and molecular chaperones in amyloid fibril formation. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 396, 2-6	3.4	24
82	Conformational preferences of non-polar amino acid residues: an additional factor in amyloid formation. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 402, 515-8	3.4	21
81	Synthetic surfactant based on analogues of SP-B and SP-C is superior to single-peptide surfactants in ventilated premature rabbits. <i>Neonatology</i> , 2010 , 98, 91-9	4	44
80	Sterilized recombinant spider silk fibers of low pyrogenicity. <i>Biomacromolecules</i> , 2010 , 11, 953-9	6.9	43
79	Recombinant spider silk as matrices for cell culture. <i>Biomaterials</i> , 2010 , 31, 9575-85	15.6	89
78	Tissue Response to Subcutaneously Implanted Recombinant Spider Silk: An in Vivo Study. <i>Materials</i> , 2009 , 2, 1908-1922	3.5	65
77	BRICHOS - a superfamily of multidomain proteins with diverse functions. <i>BMC Research Notes</i> , 2009 , 2, 180	2.3	64
76	Peptide-binding specificity of the prosurfactant protein C Brichos domain analyzed by electrospray ionization mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2009 , 23, 3591-8	2.2	15
75	Engineered disulfides improve mechanical properties of recombinant spider silk. <i>Protein Science</i> , 2009 , 18, 1012-22	6.3	34
74	The Brichos domain of prosurfactant protein C can hold and fold a transmembrane segment. <i>Protein Science</i> , 2009 , 18, 1175-82	6.3	31
73	Preventing amyloid formation by catching unfolded transmembrane segments. <i>Journal of Molecular Biology</i> , 2009 , 389, 227-9	6.5	24
72	Anti-amyloid activity of the C-terminal domain of proSP-C against amyloid beta-peptide and medin. <i>Biochemistry</i> , 2009 , 48, 3778-86	3.2	34
71	Biomimicry of surfactant protein C. Accounts of Chemical Research, 2008, 41, 1409-17	24.3	30
70	Structural properties of recombinant nonrepetitive and repetitive parts of major ampullate spidroin 1 from Euprosthenops australis: implications for fiber formation. <i>Biochemistry</i> , 2008 , 47, 3407-	1 3 ²	113

69	Surfactant proteins B and C are both necessary for alveolar stability at end expiration in premature rabbits with respiratory distress syndrome. <i>Journal of Applied Physiology</i> , 2008 , 104, 1101-8	3.7	47
68	Mutations linked to interstitial lung disease can abrogate anti-amyloid function of prosurfactant protein C. <i>Biochemical Journal</i> , 2008 , 416, 201-9	3.8	45
67	C-terminal, endoplasmic reticulum-lumenal domain of prosurfactant protein C - structural features and membrane interactions. <i>FEBS Journal</i> , 2008 , 275, 536-47	5.7	19
66	Macroscopic fibers self-assembled from recombinant miniature spider silk proteins. <i>Biomacromolecules</i> , 2007 , 8, 1695-701	6.9	144
65	Major ampullate spidroins from Euprosthenops australis: multiplicity at protein, mRNA and gene levels. <i>Insect Molecular Biology</i> , 2007 , 16, 551-61	3.4	39
64	N-terminal nonrepetitive domain common to dragline, flagelliform, and cylindriform spider silk proteins. <i>Biomacromolecules</i> , 2006 , 7, 3120-4	6.9	122
63	The Brichos domain-containing C-terminal part of pro-surfactant protein C binds to an unfolded poly-val transmembrane segment. <i>Journal of Biological Chemistry</i> , 2006 , 281, 21032-21039	5.4	54
62	Structure and influence on stability and activity of the N-terminal propeptide part of lung surfactant protein C. <i>FEBS Journal</i> , 2006 , 273, 926-35	5.7	7
61	Amyloid fibrils. FEBS Journal, 2005, 272, 5941	5.7	10
60	Mapping and analysis of the lytic and fusogenic domains of surfactant protein B. <i>Biochemistry</i> , 2005 , 44, 861-72	3.2	69
59	Processing of pulmonary surfactant protein B by napsin and cathepsin H. <i>Journal of Biological Chemistry</i> , 2004 , 279, 16178-84	5.4	105
58	N-terminally extended surfactant protein (SP) C isolated from SP-B-deficient children has reduced surface activity and inhibited lipopolysaccharide binding. <i>Biochemistry</i> , 2004 , 43, 3891-8	3.2	31
57	The N-terminal propeptide of lung surfactant protein C is necessary for biosynthesis and prevents unfolding of a metastable alpha-helix. <i>Journal of Molecular Biology</i> , 2004 , 338, 857-62	6.5	15
56	Cellular antiendotoxin activities of lung surfactant protein C in lipid vesicles. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2003 , 168, 335-41	10.2	28
55	A synthetic surfactant based on a poly-Leu SP-C analog and phospholipids: effects on tidal volumes and lung gas volumes in ventilated immature newborn rabbits. <i>Journal of Applied Physiology</i> , 2003 , 95, 2055-63	3.7	51
54	Interaction of pulmonary surfactant protein C with CD14 and lipopolysaccharide. <i>Infection and Immunity</i> , 2003 , 71, 61-7	3.7	56
53	Structural basis for interactions between lung surfactant protein C and bacterial lipopolysaccharide. <i>Journal of Biological Chemistry</i> , 2002 , 277, 23484-92	5.4	49
52	Hydrogen/deuterium exchange and aggregation of a polyvaline and a polyleucine alpha-helix investigated by matrix-assisted laser desorption ionization mass spectrometry. <i>Molecular and Cellular Proteomics</i> , 2002 , 1, 592-7	7.6	24

(1998-2002)

51	Molecular effects of proinsulin C-peptide. <i>Biochemical and Biophysical Research Communications</i> , 2002 , 295, 1035-40	3.4	54
50	Membrane properties and amyloid fibril formation of lung surfactant protein C. <i>Biochemical Society Transactions</i> , 2001 , 29, 601-6	5.1	24
49	Membrane properties and amyloid fibril formation of lung surfactant protein C. <i>Biochemical Society Transactions</i> , 2001 , 29, A56-A56	5.1	
48	Free energy barrier estimation of unfolding the alpha-helical surfactant-associated polypeptide C. <i>Proteins: Structure, Function and Bioinformatics</i> , 2001 , 43, 395-402	4.2	9
47	Prediction of amyloid fibril-forming proteins. <i>Journal of Biological Chemistry</i> , 2001 , 276, 12945-50	5.4	240
46	Artificial surfactants based on analogues of SP-B and SP-C. Fetal and Pediatric Pathology, 2001, 20, 501-	18	10
45	The palmitoyl groups of lung surfactant protein C reduce unfolding into a fibrillogenic intermediate. <i>Journal of Molecular Biology</i> , 2001 , 310, 937-50	6.5	42
44	Characterisation of variant forms of prophenin: mechanistic aspects of the fragmentation of proline-rich peptides. <i>Rapid Communications in Mass Spectrometry</i> , 2000 , 14, 2182-202	2.2	25
43	Palmitoylation of a pulmonary surfactant protein C analogue affects the surface associated lipid reservoir and film stability. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2000 , 1466, 169-78	3.8	35
42	Pulmonary surfactant protein B: a structural model and a functional analogue. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2000 , 1466, 179-86	3.8	57
41	A method for S- and O-palmitoylation of peptides: synthesis of pulmonary surfactant protein-C models. <i>Biochemical Journal</i> , 1999 , 343, 557-562	3.8	40
40	Isolation and characterization of hydrophobic polypeptides in human bile. <i>FEBS Journal</i> , 1999 , 266, 209	-14	25
39	Pulmonary surfactant: emerging protein analogues. <i>BioDrugs</i> , 1999 , 11, 71-7	7.9	2
38	Porcine pulmonary surfactant preparations contain the antibacterial peptide prophenin and a C-terminal 18-residue fragment thereof. <i>FEBS Letters</i> , 1999 , 460, 257-62	3.8	19
37	Amyloid fibril formation by pulmonary surfactant protein C. FEBS Letters, 1999, 464, 138-42	3.8	86
36	Biophysical activity of an artificial surfactant containing an analogue of surfactant protein (SP)-C and native SP-B. <i>Biochemical Journal</i> , 1999 , 339, 381-386	3.8	28
35	Synthetic peptide-containing surfactantsevaluation of transmembrane versus amphipathic helices and surfactant protein C poly-valyl to poly-leucyl substitution. <i>FEBS Journal</i> , 1998 , 255, 116-24		77
34	Determination of proteins, phosphatidylethanolamine, and phosphatidylserine in organic solvent extracts of tissue material by analysis of phenylthiocarbamyl derivatives. <i>Analytical Biochemistry</i> , 1998, 265, 97-102	3.1	28

33	Analysis of variant forms of porcine surfactant polypeptide-C by nano-electrospray mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 1998 , 12, 1104-14	2.2	11
32	Pulmonary surfactant-associated polypeptide C in a mixed organic solvent transforms from a monomeric alpha-helical state into insoluble beta-sheet aggregates. <i>Protein Science</i> , 1998 , 7, 2533-40	6.3	76
31	Structure and properties of surfactant protein C. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 1998 , 1408, 161-72	6.9	132
30	Secondary structure and limited proteolysis give experimental evidence that the precursor of pulmonary surfactant protein B contains three saposin-like domains. <i>FEBS Letters</i> , 1998 , 423, 1-4	3.8	19
29	Reverse-phase HPLC of the hydrophobic pulmonary surfactant proteins: detection of a surfactant protein C isoform containing Nepsilon-palmitoyl-lysine. <i>Biochemical Journal</i> , 1997 , 326 (Pt 3), 799-806	3.8	54
28	Molecular structures and interactions of pulmonary surfactant components. <i>FEBS Journal</i> , 1997 , 244, 675-93		233
27	The 21-residue surfactant peptide (LysLeu4)4Lys(KL4) is a transmembrane alpha-helix with a mixed nonpolar/polar surface. <i>FEBS Letters</i> , 1996 , 384, 185-8	3.8	55
26	Synthetic protein analogues in artificial surfactants. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 1996 , 85, 642-6	3.1	13
25	Pulmonary surfactant-associated polypeptide SP-C in lipid micelles: CD studies of intact SP-C and NMR secondary structure determination of depalmitoyl-SP-C(1-17). <i>FEBS Letters</i> , 1995 , 362, 261-5	3.8	52
24	An amphipathic helical motif common to tumourolytic polypeptide NK-lysin and pulmonary surfactant polypeptide SP-B. <i>FEBS Letters</i> , 1995 , 362, 328-32	3.8	123
23	The effect of environment on the stability of an integral membrane helix: molecular dynamics simulations of surfactant protein C in chloroform, methanol and water. <i>Journal of Molecular Biology</i> , 1995 , 247, 808-22	6.5	34
22	Secondary structure and biophysical activity of synthetic analogues of the pulmonary surfactant polypeptide SP-C. <i>Biochemical Journal</i> , 1995 , 307 (Pt 2), 535-41	3.8	85
21	Structural and Functional Importance of the C-Terminal Part of the Pulmonary Surfactant Polypeptide SP-C. <i>FEBS Journal</i> , 1995 , 229, 465-472		24
20	Secondary structure and membrane interaction of PR-39, a Pro+Arg-rich antibacterial peptide. <i>FEBS Journal</i> , 1994 , 224, 1019-27		87
19	The NMR structure of the pulmonary surfactant-associated polypeptide SP-C in an apolar solvent contains a valyl-rich alpha-helix. <i>Biochemistry</i> , 1994 , 33, 6015-23	3.2	184
18	The proteins of the surfactant system. European Respiratory Journal, 1994, 7, 372-91	13.6	152
17	Secondary structure and orientation of the surfactant protein SP-B in a lipid environment. A Fourier transform infrared spectroscopy study. <i>Biochemistry</i> , 1992 , 31, 9169-76	3.2	136
16	Identification of hydrophobic fragments of alpha 1-antitrypsin and C1 protease inhibitor in human bile, plasma and spleen. <i>FEBS Letters</i> , 1992 , 299, 146-8	3.8	32

LIST OF PUBLICATIONS

15	Cysteine reactivity in sorbitol and aldehyde dehydrogenases. Differences towards the pattern in alcohol dehydrogenase. <i>FEBS Letters</i> , 1992 , 303, 1-3	3.8	6
14	Human surfactant polypeptide SP-B. Disulfide bridges, C-terminal end, and peptide analysis of the airway form. <i>FEBS Letters</i> , 1992 , 301, 165-7	3.8	27
13	Structure and orientation of the surfactant-associated protein C in a lipid bilayer. <i>FEBS Journal</i> , 1992 , 203, 201-9		149
12	Surfactant protein B: disulfide bridges, structural properties, and kringle similarities. <i>Biochemistry</i> , 1991 , 30, 6917-21	3.2	112
11	Closely related isozymes of alcohol dehydrogenase. Carboxymethylation: gamma 1 gamma 1 differs widely from both beta 1 beta 1 and its equine equivalence EE. <i>FEBS Letters</i> , 1991 , 279, 119-22	3.8	5
10	Canine hydrophobic surfactant polypeptide SP-C. A lipopeptide with one thioester-linked palmitoyl group. <i>FEBS Letters</i> , 1991 , 281, 119-22	3.8	42
9	Hydrophobic surfactant-associated polypeptides: SP-C is a lipopeptide with two palmitoylated cysteine residues, whereas SP-B lacks covalently linked fatty acyl groups. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990 , 87, 2985-9	11.5	205
8	Low-molecular-mass surfactant protein type 1. The primary structure of a hydrophobic 8-kDa polypeptide with eight half-cystine residues. <i>FEBS Journal</i> , 1988 , 172, 521-5		86
7	Mitochondrial aldehyde dehydrogenase from horse liver. Correlations of the same species variants for both the cytosolic and the mitochondrial forms of an enzyme. <i>FEBS Journal</i> , 1988 , 172, 527-33		53
6	Size and structure of the hydrophobic low molecular weight surfactant-associated polypeptide. <i>Biochemistry</i> , 1988 , 27, 3544-7	3.2	74
5	Hydrophobic 3.7 kDa surfactant polypeptide: structural characterization of the human and bovine forms. <i>FEBS Letters</i> , 1988 , 232, 61-4	3.8	50
4	Alcohol dehydrogenases and aldehyde dehydrogenases. <i>Biochemical Society Transactions</i> , 1988 , 16, 223	-₹ .1	11
3	Superoxide dismutase in human testis preparations. <i>Bioscience Reports</i> , 1986 , 6, 535-41	4.1	2
2	Peptides of postulated inhibin activity. Lack of in vitro inhibin activity of a 94-residue peptide isolated from human seminal plasma, and of a synthetic replicate of its C-terminal 28-residue segment. <i>FEBS Letters</i> , 1986 , 199, 242-8	3.8	37
1	Engineered Spider Silk Proteins for Biomimetic Spinning of Fibers with Toughness Equal to Dragline Silks. <i>Advanced Functional Materials</i> ,2200986	15.6	О