

Qingshan Wang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6570512/publications.pdf>

Version: 2024-02-01

47
papers

1,684
citations

257450

24
h-index

302126

39
g-index

50
all docs

50
docs citations

50
times ranked

2030
citing authors

#	ARTICLE	IF	CITATIONS
1	Rotenone impairs learning and memory in mice through microglia-mediated blood brain barrier disruption and neuronal apoptosis. <i>Chemosphere</i> , 2022, 291, 132982.	8.2	24
2	Microglial Activation Damages Dopaminergic Neurons through MMP-2/-9-Mediated Increase of Blood-Brain Barrier Permeability in a Parkinson's Disease Mouse Model. <i>International Journal of Molecular Sciences</i> , 2022, 23, 2793.	4.1	24
3	A novel synthetic peptide SVHRSP attenuates dopaminergic neurodegeneration by inhibiting NADPH oxidase-mediated neuroinflammation in experimental models of Parkinson's disease. <i>Free Radical Biology and Medicine</i> , 2022, 188, 363-374.	2.9	9
4	2,5-hexanedione induces NLRP3 inflammasome activation and neurotoxicity through NADPH oxidase-dependent pathway. <i>Free Radical Biology and Medicine</i> , 2021, 162, 561-570.	2.9	9
5	Microglial activation contributes to cognitive impairments in rotenone-induced mouse Parkinson's disease model. <i>Journal of Neuroinflammation</i> , 2021, 18, 4.	7.2	70
6	Microglial Activation Mediates Noradrenergic Locus Coeruleus Neurodegeneration via Complement Receptor 3 in a Rotenone-Induced Parkinson's Disease Mouse Model. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 1341-1356.	3.5	17
7	Fertility intentions among the working population of Dalian City born between 1980 and 1989. <i>Journal of Biosocial Science</i> , 2021, , 1-12.	1.2	4
8	Inhibition of NLRP3 inflammasome by glibenclamide attenuated dopaminergic neurodegeneration and motor deficits in paraquat and maneb-induced mouse Parkinson's disease model. <i>Toxicology Letters</i> , 2021, 349, 1-11.	0.8	20
9	Clinical Efficacy and Safety of Lacosamide as an Adjunctive Treatment in Adults With Refractory Epilepsy. <i>Frontiers in Neurology</i> , 2021, 12, 712717.	2.4	5
10	Locus coeruleus neurons are most sensitive to chronic neuroinflammation-induced neurodegeneration. <i>Brain, Behavior, and Immunity</i> , 2020, 87, 359-368.	4.1	33
11	NLRP3 inflammasome mediates 2,5-hexanedione-induced neurotoxicity through regulation of macrophage infiltration in rats. <i>Chemico-Biological Interactions</i> , 2020, 330, 109232.	4.0	7
12	Integrin CD11b mediates locus coeruleus noradrenergic neurodegeneration in a mouse Parkinson's disease model. <i>Journal of Neuroinflammation</i> , 2020, 17, 148.	7.2	23
13	Glibenclamide attenuates 2,5-hexanedione-induced neurotoxicity in the spinal cord of rats through mitigation of NLRP3 inflammasome activation, neuroinflammation and oxidative stress. <i>Toxicology Letters</i> , 2020, 331, 152-158.	0.8	18
14	Nicotinamide Adenine Dinucleotide Phosphate Oxidase and Neurodegenerative Diseases: Mechanisms and Therapy. <i>Antioxidants and Redox Signaling</i> , 2020, 33, 374-393.	5.4	22
15	Loss of Brain Norepinephrine Elicits Neuroinflammation-Mediated Oxidative Injury and Selective Caudo-Rostral Neurodegeneration. <i>Molecular Neurobiology</i> , 2019, 56, 2653-2669.	4.0	50
16	Noradrenergic dysfunction accelerates LPS-elicited inflammation-related ascending sequential neurodegeneration and deficits in non-motor/motor functions. <i>Brain, Behavior, and Immunity</i> , 2019, 81, 374-387.	4.1	36
17	Lesion of the Locus Coeruleus Damages Learning and Memory Performance in Paraquat and Maneb-induced Mouse Parkinson's Disease Model. <i>Neuroscience</i> , 2019, 419, 129-140.	2.3	33
18	NADPH oxidase regulates paraquat and maneb-induced dopaminergic neurodegeneration through ferroptosis. <i>Toxicology</i> , 2019, 417, 64-73.	4.2	80

#	ARTICLE	IF	CITATIONS
19	Inhibition of NADPH oxidase by apocynin prevents learning and memory deficits in a mouse Parkinson's disease model. <i>Redox Biology</i> , 2019, 22, 101134.	9.0	85
20	NMDA Receptor Antagonist MK801 Protects Against 1-Bromopropane-Induced Cognitive Dysfunction. <i>Neuroscience Bulletin</i> , 2019, 35, 347-361.	2.9	11
21	Allyl Sulfide Counteracts 1-Bromopropane-Induced Neurotoxicity by Inhibiting Neuroinflammation and Oxidative Stress. <i>Toxicological Sciences</i> , 2019, 167, 397-407.	3.1	9
22	Taurine protects noradrenergic locus coeruleus neurons in a mouse Parkinson's disease model by inhibiting microglial M1 polarization. <i>Amino Acids</i> , 2018, 50, 547-556.	2.7	43
23	Physiological Concentration of Prostaglandin E2 Exerts Anti-inflammatory Effects by Inhibiting Microglial Production of Superoxide Through a Novel Pathway. <i>Molecular Neurobiology</i> , 2018, 55, 8001-8013.	4.0	10
24	Taurine protects dopaminergic neurons in a mouse Parkinson's disease model through inhibition of microglial M1 polarization. <i>Cell Death and Disease</i> , 2018, 9, 435.	6.3	100
25	2,5-Hexanedione induces dopaminergic neurodegeneration through integrin α 2 β 2/NADPH oxidase axis-mediated microglial activation. <i>Cell Death and Disease</i> , 2018, 9, 60.	6.3	23
26	Integrin CD11b mediates α -synuclein-induced activation of NADPH oxidase through a Rho-dependent pathway. <i>Redox Biology</i> , 2018, 14, 600-608.	9.0	69
27	Bone marrow mesenchymal stem cells protect against n-hexane-induced neuropathy through beclin 1-independent inhibition of autophagy. <i>Scientific Reports</i> , 2018, 8, 4516.	3.3	8
28	Complement receptor 3 mediates NADPH oxidase activation and dopaminergic neurodegeneration through a Src-Erk-dependent pathway. <i>Redox Biology</i> , 2018, 14, 250-260.	9.0	72
29	Hypertension and Diagnosis of Parkinson's Disease: A Meta-Analysis of Cohort Studies. <i>Frontiers in Neurology</i> , 2018, 9, 162.	2.4	48
30	The mechanism of neuroinflammation in Parkinson's disease: the involvement of integrin. <i>Proceedings for Annual Meeting of the Japanese Pharmacological Society</i> , 2018, WCP2018, PO4-1-125.	0.0	0
31	Paraquat and maneb co-exposure induces noradrenergic locus coeruleus neurodegeneration through NADPH oxidase-mediated microglial activation. <i>Toxicology</i> , 2017, 380, 1-10.	4.2	40
32	NADPH oxidase-derived H ₂ O ₂ mediates the regulatory effects of microglia on astrogliosis in experimental models of Parkinson's disease. <i>Redox Biology</i> , 2017, 12, 162-170.	9.0	54
33	Clozapine metabolites protect dopaminergic neurons through inhibition of microglial NADPH oxidase. <i>Journal of Neuroinflammation</i> , 2016, 13, 110.	7.2	42
34	Bone marrow mesenchymal stem cells attenuate 2,5-hexanedione-induced neuronal apoptosis through a NGF/AKT-dependent pathway. <i>Scientific Reports</i> , 2016, 6, 34715.	3.3	21
35	Neurons and astroglia govern microglial endotoxin tolerance through macrophage colony-stimulating factor receptor-mediated ERK1/2 signals. <i>Brain, Behavior, and Immunity</i> , 2016, 55, 260-272.	4.1	31
36	Substance P enhances microglial density in the substantia nigra through neurokinin-1 receptor/NADPH oxidase-mediated chemotaxis in mice. <i>Clinical Science</i> , 2015, 129, 757-767.	4.3	21

#	ARTICLE	IF	CITATIONS
37	Post-treatment with an ultra-low dose of NADPH oxidase inhibitor diphenyleneiodonium attenuates disease progression in multiple Parkinson's disease models. <i>Brain</i> , 2015, 138, 1247-1262.	7.6	86
38	A novel role of microglial NADPH oxidase in mediating extra-synaptic function of norepinephrine in regulating brain immune homeostasis. <i>Glia</i> , 2015, 63, 1057-1072.	4.9	53
39	PGE2 Inhibits IL-10 Production via EP2-Mediated β -Arrestin Signaling in Neuroinflammatory Condition. <i>Molecular Neurobiology</i> , 2015, 52, 587-600.	4.0	24
40	Microglial regulation of immunological and neuroprotective functions of astroglia. <i>Glia</i> , 2015, 63, 118-131.	4.9	84
41	Substance P Exacerbates Dopaminergic Neurodegeneration through Neurokinin-1 Receptor-Independent Activation of Microglial NADPH Oxidase. <i>Journal of Neuroscience</i> , 2014, 34, 12490-12503.	3.6	70
42	Subpicomolar diphenyleneiodonium inhibits microglial NADPH oxidase with high specificity and shows great potential as a therapeutic agent for neurodegenerative diseases. <i>Glia</i> , 2014, 62, 2034-2043.	4.9	46
43	Endogenous dynorphin protects against neurotoxin-elicited nigrostriatal dopaminergic neuron damage and motor deficits in mice. <i>Journal of Neuroinflammation</i> , 2012, 9, 124.	7.2	65
44	Naloxone inhibits immune cell function by suppressing superoxide production through a direct interaction with gp91 phox subunit of NADPH oxidase. <i>Journal of Neuroinflammation</i> , 2012, 9, 32.	7.2	56
45	Carboxyl-terminus of Hsc70 interacting protein mediates 2,5-hexanedione-induced neurofilament medium chain degradation. <i>Biochemical Pharmacology</i> , 2011, 81, 793-799.	4.4	14
46	Alterations in neurofilaments content and calpains activity of sciatic nerve of carbon disulfide-treated rats. <i>Archives of Toxicology</i> , 2009, 83, 587-594.	4.2	7
47	The reversibility of neurofilaments decline induced by 2,5-hexanedione in rat nerve tissues. <i>Biochemical Pharmacology</i> , 2008, 75, 737-744.	4.4	8