

# Piotr Donizy

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6572832/publications.pdf>

Version: 2024-02-01

50  
papers

567  
citations

840119

11  
h-index

713013

21  
g-index

50  
all docs

50  
docs citations

50  
times ranked

1133  
citing authors

#	ARTICLE	IF	CITATIONS
1	Cancer stem cells – the current status of an old concept: literature review and clinical approaches. <i>Biological Research</i> , 2014, 47, 66.	1.5	60
2	Clinical, Histopathological and Cytogenetic Prognosticators in Uveal Melanoma – A Comprehensive Review. <i>Anticancer Research</i> , 2017, 37, 6541-6549.	0.5	38
3	Prognostic significance of ALCAM (CD166/MEMD) expression in cutaneous melanoma patients. <i>Diagnostic Pathology</i> , 2015, 10, 86.	0.9	31
4	Golgi-Related Proteins GOLPH2 (GP73/GOLM1) and GOLPH3 (GOPP1/MIDAS) in Cutaneous Melanoma: Patterns of Expression and Prognostic Significance. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1619.	1.8	28
5	Augmented expression of Polo-like kinase 1 is a strong predictor of shorter cancer-specific overall survival in early stage breast cancer at 15-year follow-up. <i>Oncology Letters</i> , 2016, 12, 1667-1674.	0.8	28
6	Macrophages in skin melanoma-the key element in melanomagenesis (Review). <i>Oncology Letters</i> , 2018, 15, 5399-5404.	0.8	26
7	Paucity of tumor-infiltrating lymphocytes is an unfavorable prognosticator and predicts lymph node metastases in cutaneous melanoma patients. <i>Anticancer Research</i> , 2015, 35, 351-8.	0.5	26
8	Nuclear-cytoplasmic PARP-1 expression as an unfavorable prognostic marker in lymph node-negative early breast cancer: 15-year follow-up. <i>Oncology Reports</i> , 2014, 31, 1777-1787.	1.2	23
9	Mitotic rate is a more reliable unfavorable prognosticator than ulceration for early cutaneous melanoma: A 5-year survival analysis. <i>Oncology Reports</i> , 2014, 32, 2735-2743.	1.2	23
10	Up-Regulation of PARP1 Expression Significantly Correlated with Poor Survival in Mucosal Melanomas. <i>Cells</i> , 2020, 9, 1135.	1.8	22
11	Nucleoli cytomorphology in cutaneous melanoma cells – a new prognostic approach to an old concept. <i>Diagnostic Pathology</i> , 2017, 12, 88.	0.9	15
12	Angiotensin II Type 1 Receptor Expression in Renal Transplant Biopsies and Anti-AT1R Antibodies in Serum Indicates the Risk of Transplant Loss. <i>Transplantation Proceedings</i> , 2020, 52, 2299-2304.	0.3	13
13	Morphologic Diversity of Merkel Cell Carcinoma. <i>American Journal of Dermatopathology</i> , 2020, 42, 629-640.	0.3	13
14	TdT Expression Is a Marker of Better Survival in Merkel Cell Carcinoma, and Expression of B-Cell Markers Is Associated With Merkel Cell Polyomavirus. <i>American Journal of Clinical Pathology</i> , 2020, 154, 38-47.	0.4	13
15	The variety of clinical presentations in IgG4-related disease in Rheumatology. <i>Rheumatology International</i> , 2018, 38, 303-309.	1.5	11
16	SMAD7 is a novel independent predictor of survival in patients with cutaneous melanoma. <i>Translational Research</i> , 2019, 204, 72-81.	2.2	10
17	Ephrin Receptors (Eph): EphA1, EphA5, and EphA7 Expression in Uveal Melanoma – Associations with Clinical Parameters and Patient Survival. <i>Life</i> , 2020, 10, 225.	1.1	10
18	Merkel Cell Carcinoma of Unknown Primary: Immunohistochemical and Molecular Analyses Reveal Distinct UV-Signature/MCPyV-Negative and High Immunogenicity/MCPyV-Positive Profiles. <i>Cancers</i> , 2021, 13, 1621.	1.7	10

#	ARTICLE	IF	CITATIONS
19	Prognostic significance of immunohistochemical epithelial-mesenchymal transition markers in skin melanoma patients. <i>Biomarkers in Medicine</i> , 2016, 10, 975-985.	0.6	9
20	Epithelial-mesenchymal transition inducer Snail1 and invasive potential of intraductal breast cancer. <i>Journal of Surgical Oncology</i> , 2017, 116, 696-705.	0.8	9
21	The effect of YAP expression in tumor cells and tumor stroma on the prognosis of patients with squamous cell carcinoma of the oral cavity floor and oral surface of the tongue. <i>Oncology Letters</i> , 2019, 18, 3561-3570.	0.8	9
22	Histone Deacetylase (HDAC)-1, -2, -4, and -6 in Uveal Melanomas: Associations with Clinicopathological Parameters and Patients' Survival. <i>Cancers</i> , 2021, 13, 4763.	1.7	9
23	PARP1 as a Marker of an Aggressive Clinical Phenotype in Cutaneous Melanoma: A Clinical and an In Vitro Study. <i>Cells</i> , 2021, 10, 286.	1.8	9
24	Ductal carcinoma in situ on stereotactic biopsy of suspicious breast microcalcifications: Expression of SPARC (Secreted Protein, Acidic and Rich in Cysteine) can predict postoperative invasion. <i>Journal of Surgical Oncology</i> , 2016, 114, 548-556.	0.8	8
25	Upregulation of FOXP1 is a new independent unfavorable prognosticator and a specific predictor of lymphatic dissemination in cutaneous melanoma patients. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 1413-1422.	1.0	8
26	ROCK1 and ROCK2 Are Down-regulated in Aggressive and Advanced Skin Melanomas: A Clinicopathological Perspective. <i>Anticancer Research</i> , 2020, 40, 1931-1942.	0.5	8
27	High Percentage of ADAM-10 Positive Melanoma Cells Correlates with Paucity of Tumor-Infiltrating Lymphocytes but Does Not Predict Prognosis in Cutaneous Melanoma Patients. <i>Analytical Cellular Pathology</i> , 2015, 2015, 1-7.	0.7	7
28	Prognostic Role of Tumoral PD-L1 and IDO1 Expression, and Intratumoral CD8+ and FoxP3+ Lymphocyte Infiltrates in 132 Primary Cutaneous Merkel Cell Carcinomas. <i>International Journal of Molecular Sciences</i> , 2021, 22, 5489.	1.8	7
29	Selected Golgi-Localized Proteins and Carcinogenesis: What Do We Know?. <i>Results and Problems in Cell Differentiation</i> , 2019, 67, 487-529.	0.2	7
30	Biomarkers in Primary Focal Segmental Glomerulosclerosis in Optimal Diagnostic-Therapeutic Strategy. <i>Journal of Clinical Medicine</i> , 2022, 11, 3292.	1.0	7
31	Large nuclear size correlated with better overall survival, Merkel cell polyomavirus positivity, and terminal deoxynucleotidyl transferase expression in Merkel cell carcinoma. <i>Journal of the American Academy of Dermatology</i> , 2021, 84, 550-552.	0.6	6
32	Endothelin A Receptors Expressed in Glomeruli of Renal Transplant Patients May Be Associated with Antibody-Mediated Rejection. <i>Journal of Clinical Medicine</i> , 2021, 10, 422.	1.0	6
33	Clinical and Histopathological Factors Influencing IgA Nephropathy Outcome. <i>Diagnostics</i> , 2021, 11, 1764.	1.3	6
34	Amelanotic Uveal Melanomas Evaluated by Indirect Ophthalmoscopy Reveal Better Long-Term Prognosis Than Pigmented Primary Tumours: A Single Centre Experience. <i>Cancers</i> , 2022, 14, 2753.	1.7	6
35	Polo-like kinase-1 immunoreactivity is associated with metastases in cutaneous melanoma. <i>Journal of Cutaneous Pathology</i> , 2017, 44, 819-826.	0.7	5
36	Biological Aggressiveness of Subclinical No-Mass Ductal Carcinoma In Situ (DCIS) Can Be Reflected by the Expression Profiles of Epithelial-Mesenchymal Transition Triggers. <i>International Journal of Molecular Sciences</i> , 2018, 19, 3941.	1.8	5

#	ARTICLE	IF	CITATIONS
37	The Prognostic Values of PARP-1 Expression in Uveal Melanoma. <i>Cells</i> , 2021, 10, 285.	1.8	5
38	Downregulation of Polo-like kinase-1 (PLK-1) expression is associated with poor clinical outcome in uveal melanoma patients. <i>Folia Histochemica Et Cytobiologica</i> , 2020, 58, 108-116.	0.6	5
39	Intratumoral but not peritumoral lymphatic vessel density measured by D2-40 expression predicts poor outcome in gastric cancer-ROC curve analysis to find cut-off point. <i>Anticancer Research</i> , 2014, 34, 3113-8.	0.5	5
40	The Summarized Assessment of Endothelin A Receptor Expression in Renal Transplant Compartments Associated with Antibody-Mediated Rejection. <i>Diagnostics</i> , 2021, 11, 2366.	1.3	4
41	Nuclear pseudoinclusions in melanoma cells: prognostic fact or artifact? The possible role of Golgi phosphoprotein 3 overexpression in nuclear pseudoinclusions generation. <i>Pathology International</i> , 2018, 68, 117-122.	0.6	3
42	Clinical Significance of Nucleoli Cytomorphology Assessment in Patients With Uveal Melanoma. <i>Anticancer Research</i> , 2020, 40, 3505-3512.	0.5	3
43	Lymphangioinvasion in routine H&E staining is strongly associated with poor clinical outcome in lymph node-negative cutaneous melanoma patients. <i>Folia Histochemica Et Cytobiologica</i> , 2016, 54, 126-133.	0.6	3
44	Alpha-Enolase (ENO1) Correlates with Invasiveness of Cutaneous Melanoma—An In Vitro and a Clinical Study. <i>Diagnostics</i> , 2022, 12, 254.	1.3	3
45	Schwannoma in the porta hepatis – laparoscopic excision under laparoscopic ultrasound guidance. <i>Wideochirurgia I Inne Techniki Maloinwazyjne</i> , 2017, 3, 301-305.	0.3	2
46	EPHA2, EPHA4, and EPHA6 Expression in Uveal Melanomas: Searching for the Culprits of Neoplasia. <i>Diagnostics</i> , 2022, 12, 1025.	1.3	2
47	The Length of Leukocyte and Femoral Artery Telomeres in Patients with Peripheral Atherosclerosis. <i>Genes</i> , 2022, 13, 704.	1.0	1
48	SP700THE PRESENCE OF ANTI-AT1R ANTIBODIES IN BLOOD AND AT1 RECEPTOR EXPRESSION IN BIOPSY FOR CAUSE OF RENAL TRANSPLANT RECIPIENTS MAY BE ASSOCIATED WITH HIGHER GRAFT LOSS AND MORE ANTIBODY MEDIATED REJECTION CASES. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i582-i582.	0.4	0
49	Nuclear Pseudoinclusions and Intranuclear Grooves Have an Important Impact on the Long-term Survival of Patients With Uveal Melanoma. <i>Anticancer Research</i> , 2021, 41, 517-526.	0.5	0
50	BILLCD8 – A Multivariable Survival Model as a Simple and Clinically Useful Prognostic Tool to Identify High-risk Cutaneous Melanoma Patients. <i>Anticancer Research</i> , 2016, 36, 4739-4748.	0.5	0