

Xing Chen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6573426/publications.pdf>

Version: 2024-02-01

147
papers

5,625
citations

66343

42
h-index

110387

64
g-index

147
all docs

147
docs citations

147
times ranked

2598
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Healthy benefits and edible delivery systems of resveratrol: a review. Food Reviews International, 2023, 39, 3879-3905. | 8.4 | 2 |
| 2 | Insight Into the Effect of Carnosine on the Dispersibility of Myosin Under a Low-salt Condition and its Mechanism. Food Biophysics, 2023, 18, 71-81. | 3.0 | 1 |
| 3 | Impact of Phytophenols on Myofibrillar Proteins: Revisit the Interaction Scenarios Inspired for Meat Products Innovation. Food Reviews International, 2023, 39, 5637-5665. | 8.4 | 2 |
| 4 | Stabilization of O/W emulsions via interfacial protein concentrating induced by thermodynamic incompatibility between sarcoplasmic proteins and xanthan gum. Food Hydrocolloids, 2022, 124, 107242. | 10.7 | 22 |
| 5 | Trace the difference driven by unfolding-refolding pathway of myofibrillar protein: Emphasizing the changes on structural and emulsion properties. Food Chemistry, 2022, 367, 130688. | 8.2 | 37 |
| 6 | Optimizing 3D printing of chicken meat by response surface methodology and genetic algorithm: Feasibility study of 3D printed chicken product. LWT - Food Science and Technology, 2022, 154, 112693. | 5.2 | 20 |
| 7 | Effect of high-pressure treatment on the heat-induced emulsion gelation of rabbit myosin. LWT - Food Science and Technology, 2022, 154, 112719. | 5.2 | 4 |
| 8 | Interfacial rheology of alkali pH-shifted myofibrillar protein at O/W interface and impact of Tween 20 displacement. Food Hydrocolloids, 2022, 124, 107275. | 10.7 | 19 |
| 9 | Characterization and bioactivity of phlorotannin loaded protein-polysaccharide nanocomplexes. LWT - Food Science and Technology, 2022, 155, 112998. | 5.2 | 14 |
| 10 | Structural basis for high-intensity ultrasound treatment in the rheology of myofibrillar protein extracted from White Croaker in relation to their solubility. LWT - Food Science and Technology, 2022, 156, 112979. | 5.2 | 18 |
| 11 | Chemical Stability of Ascorbic Acid Integrated into Commercial Products: A Review on Bioactivity and Delivery Technology. Antioxidants, 2022, 11, 153. | 5.1 | 73 |
| 12 | Sequential changes in antioxidant activity and structure of curcumin-myofibrillar protein nanocomplex during in vitro digestion. Food Chemistry, 2022, 382, 132331. | 8.2 | 9 |
| 13 | Comparison of the interfacial properties of native and refolded myofibrillar proteins subjected to pH-shifting. Food Chemistry, 2022, 380, 131734. | 8.2 | 24 |
| 14 | Improving physicochemical properties of myofibrillar proteins from wooden breast of broiler by diverse glycation strategies. Food Chemistry, 2022, 382, 132328. | 8.2 | 23 |
| 15 | Soluble Aggregates of Myofibrillar Proteins Engineered by Gallic Acid: Colloidal Structure and Resistance to <i>In Vitro</i> Gastric Digestion. Journal of Agricultural and Food Chemistry, 2022, 70, 4066-4075. | 5.2 | 26 |
| 16 | New insights into the ultrasound impact on covalent reactions of myofibrillar protein. Ultrasonics Sonochemistry, 2022, 84, 105973. | 8.2 | 26 |
| 17 | Freeze-Thawing Treatment as a Simple Way to Tune the Gel Property and Digestibility of Minced Meat from Red Swamp Crayfish (<i>Procambarus clarkii</i>). Foods, 2022, 11, 837. | 4.3 | 2 |
| 18 | Real meat and plant-based meat analogues have different in vitro protein digestibility properties. Food Chemistry, 2022, 387, 132917. | 8.2 | 45 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 19 | Tailoring protein intrinsic charge by enzymatic deamidation for solubilizing chicken breast myofibrillar protein in water. <i>Food Chemistry</i> , 2022, 385, 132512. | 8.2 | 21 |
| 20 | Phenolic modification of myofibrillar protein enhanced by ultrasound: The structure of phenol matters. <i>Food Chemistry</i> , 2022, 386, 132662. | 8.2 | 34 |
| 21 | Comparative study on the in vitro digestibility of chicken protein after different modifications. <i>Food Chemistry</i> , 2022, 385, 132652. | 8.2 | 10 |
| 22 | Recovery of emulsifying and gelling protein from waste chicken exudate by using a sustainable pH-shifting treatment. <i>Food Chemistry</i> , 2022, 387, 132886. | 8.2 | 4 |
| 23 | Interactions between the protein-epigallocatechin gallate complex and nanocrystalline cellulose: A systematic study. <i>Food Chemistry</i> , 2022, 387, 132791. | 8.2 | 8 |
| 24 | Continuous cyclic wet heating glycation to prepare myofibrillar protein-glucose conjugates: A study on the structures, solubility and emulsifying properties. <i>Food Chemistry</i> , 2022, 388, 133035. | 8.2 | 23 |
| 25 | Interactions of water-soluble myofibrillar protein with chitosan: Phase behavior, microstructure and rheological properties. <i>Innovative Food Science and Emerging Technologies</i> , 2022, 78, 103013. | 5.6 | 18 |
| 26 | Synergistic effect of preheating and different power output high-intensity ultrasound on the physicochemical, structural, and gelling properties of myofibrillar protein from chicken wooden breast. <i>Ultrasonics Sonochemistry</i> , 2022, 86, 106030. | 8.2 | 18 |
| 27 | Effects of pulsed electric fields on the conformation and gelation properties of myofibrillar proteins isolated from pale, soft, exudative (PSE)-like chicken breast meat: A molecular dynamics study. <i>Food Chemistry</i> , 2021, 342, 128306. | 8.2 | 32 |
| 28 | Synergistic effects of UVA irradiation and phlorotannin extracts of <i>Laminaria japonica</i> on properties of grass carp myofibrillar protein gel. <i>Journal of the Science of Food and Agriculture</i> , 2021, 101, 2659-2667. | 3.5 | 8 |
| 29 | Changes of myofibrillar protein structure improved the stability and distribution of baicalein in emulsion. <i>LWT - Food Science and Technology</i> , 2021, 137, 110404. | 5.2 | 9 |
| 30 | Modification of myofibrillar protein functional properties prepared by various strategies: A comprehensive review. <i>Comprehensive Reviews in Food Science and Food Safety</i> , 2021, 20, 458-500. | 11.7 | 52 |
| 31 | Effects of oxidation on the structure of collagen fibers of sea cucumber (<i>Apostichopus japonicus</i>) body wall during thermal processing. <i>LWT - Food Science and Technology</i> , 2021, 138, 110528. | 5.2 | 14 |
| 32 | Resistance of detached-cells of biofilm formed by <i>Staphylococcus aureus</i> to ultra high pressure homogenization. <i>Food Research International</i> , 2021, 139, 109954. | 6.2 | 5 |
| 33 | Temperature-dependent in vitro digestion properties of isoelectric solubilization/precipitation (ISP)-isolated PSE-like chicken protein. <i>Food Chemistry</i> , 2021, 343, 128501. | 8.2 | 13 |
| 34 | Covalent chemical modification of myofibrillar proteins to improve their gelation properties: A systematic review. <i>Comprehensive Reviews in Food Science and Food Safety</i> , 2021, 20, 924-959. | 11.7 | 34 |
| 35 | Antioxidant activity and stability of α -tocopherol, resveratrol and epigallocatechin gallate in mixture and complexation with bovine serum albumin. <i>International Journal of Food Science and Technology</i> , 2021, 56, 1788-1800. | 2.7 | 13 |
| 36 | Quality and microbial community of high pressure shucked crab (<i>Eriocheir sinensis</i>) meat stored at 4°C. <i>Journal of Food Processing and Preservation</i> , 2021, 45, e15330. | 2.0 | 12 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 37 | Robustness of protein: Using pH shifting and low speed shearing to partially recover conformation and dispersibility of myosin from pale, soft, exudative (PSE)-like chicken breast. <i>LWT - Food Science and Technology</i> , 2021, 138, 110786. | 5.2 | 8 |
| 38 | Quality characteristics of shucked crab meat (<i>Eriocheir sinensis</i>) processed by high pressure during superchilled storage. <i>Journal of Food Biochemistry</i> , 2021, 45, e13708. | 2.9 | 8 |
| 39 | Protein deamidation to produce processable ingredients and engineered colloids for emerging food applications. <i>Comprehensive Reviews in Food Science and Food Safety</i> , 2021, 20, 3788-3817. | 11.7 | 44 |
| 40 | Effect of high intensity ultrasound on the gelation properties of wooden breast meat with different NaCl contents. <i>Food Chemistry</i> , 2021, 347, 129031. | 8.2 | 28 |
| 41 | Stability improvement of reduced-fat reduced-salt meat batter through modulation of secondary and tertiary protein structures by means of high pressure processing. <i>Meat Science</i> , 2021, 176, 108439. | 5.5 | 19 |
| 42 | Loss of immobilized water and intense protein aggregation responsible for quality deterioration of ready to eat firm tofu. <i>Journal of Texture Studies</i> , 2021, 52, 492-500. | 2.5 | 3 |
| 43 | Effects of high hydrostatic pressure treatment on the emulsifying behavior of myosin and its underlying mechanism. <i>LWT - Food Science and Technology</i> , 2021, 146, 111397. | 5.2 | 24 |
| 44 | Insight into the effect of charge regulation on the binding mechanism of curcumin to myofibrillar protein. <i>Food Chemistry</i> , 2021, 352, 129395. | 8.2 | 11 |
| 45 | Dual role (promotion and inhibition) of transglutaminase in mediating myofibrillar protein gelation under malondialdehyde-induced oxidative stress. <i>Food Chemistry</i> , 2021, 353, 129453. | 8.2 | 17 |
| 46 | Ultrasound-assisted covalent reaction of myofibrillar protein: The improvement of functional properties and its potential mechanism. <i>Ultrasonics Sonochemistry</i> , 2021, 76, 105652. | 8.2 | 45 |
| 47 | Enhanced heat stability and antioxidant activity of myofibrillar protein-dextran conjugate by the covalent adduction of polyphenols. <i>Food Chemistry</i> , 2021, 352, 129376. | 8.2 | 78 |
| 48 | Characterization of whey protein-based nanocomplex to load fucoxanthin and the mechanism of action on glial cells PC12. <i>LWT - Food Science and Technology</i> , 2021, 151, 112208. | 5.2 | 13 |
| 49 | Effect of high-pressure homogenization on structural changes and emulsifying properties of chicken liver proteins isolated by isoelectric solubilization/precipitation. <i>LWT - Food Science and Technology</i> , 2021, 151, 112092. | 5.2 | 10 |
| 50 | Structural and functional modification of food proteins by high power ultrasound and its application in meat processing. <i>Critical Reviews in Food Science and Nutrition</i> , 2021, 61, 1914-1933. | 10.3 | 58 |
| 51 | Self-powered, ultra-high detectivity and high-speed near-infrared photodetectors from stacked layered MoSe ₂ /Si heterojunction. <i>Nanotechnology</i> , 2021, 32, 075201. | 2.6 | 20 |
| 52 | Water-soluble myofibrillar protein-pectin complex for enhanced physical stability near the isoelectric point: Fabrication, rheology and thermal property. <i>International Journal of Biological Macromolecules</i> , 2020, 142, 615-623. | 7.5 | 52 |
| 53 | Glycation-induced structural modification of myofibrillar protein and its relation to emulsifying properties. <i>LWT - Food Science and Technology</i> , 2020, 117, 108664. | 5.2 | 62 |
| 54 | Conformational and rheological changes of high-pressure processing treated rabbit myosin subfragments during heating. <i>LWT - Food Science and Technology</i> , 2020, 122, 108994. | 5.2 | 4 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 55 | Physicochemical and structural properties of myofibrillar proteins isolated from pale, soft, exudative (PSE)-like chicken breast meat: Effects of pulsed electric field (PEF). <i>Innovative Food Science and Emerging Technologies</i> , 2020, 59, 102277. | 5.6 | 60 |
| 56 | Advances in converting of meat protein into functional ingredient via engineering modification of high pressure homogenization. <i>Trends in Food Science and Technology</i> , 2020, 106, 12-29. | 15.1 | 32 |
| 57 | High intake of chicken and pork proteins aggravates high-fat-diet-induced inflammation and disorder of hippocampal glutamatergic system. <i>Journal of Nutritional Biochemistry</i> , 2020, 85, 108487. | 4.2 | 7 |
| 58 | Effects of different ultrasound frequencies on the structure, rheological and functional properties of myosin: Significance of quorum sensing. <i>Ultrasonics Sonochemistry</i> , 2020, 69, 105268. | 8.2 | 35 |
| 59 | Effects of ultrasound frequency mode on myofibrillar protein structure and emulsifying properties. <i>International Journal of Biological Macromolecules</i> , 2020, 163, 1768-1779. | 7.5 | 55 |
| 60 | Gallic Acid-Aided Cross-Linking of Myofibrillar Protein Fabricated Soluble Aggregates for Enhanced Thermal Stability and a Tunable Colloidal State. <i>Journal of Agricultural and Food Chemistry</i> , 2020, 68, 11535-11544. | 5.2 | 62 |
| 61 | Fucoxanthin activities motivate its nano/micro-encapsulation for food or nutraceutical application: a review. <i>Food and Function</i> , 2020, 11, 9338-9358. | 4.6 | 39 |
| 62 | Effect of wooden breast myopathy on water-holding capacity and rheological and gelling properties of chicken broiler breast batters. <i>Poultry Science</i> , 2020, 99, 3742-3751. | 3.4 | 18 |
| 63 | Processing Properties and Improvement of Pale, Soft, and Exudative-Like Chicken Meat: a Review. <i>Food and Bioprocess Technology</i> , 2020, 13, 1280-1291. | 4.7 | 15 |
| 64 | Fabrication and characterisation of whey protein isolate-“propolis”-alginate complex particles for stabilising α -tocopherol-contained emulsions. <i>International Dairy Journal</i> , 2020, 109, 104756. | 3.0 | 17 |
| 65 | Borate suppresses the scavenging activity of gallic acid and plant polyphenol extracts on DPPH radical: A potential interference to DPPH assay. <i>LWT - Food Science and Technology</i> , 2020, 131, 109769. | 5.2 | 40 |
| 66 | Isolation of novel ACE-inhibitory peptide from naked oat globulin hydrolysates <i>in silico</i> approach: Molecular docking, <i>in vivo</i> antihypertension and effects on renin and intracellular endothelin-1. <i>Journal of Food Science</i> , 2020, 85, 1328-1337. | 3.1 | 32 |
| 67 | Influence of extreme alkaline pH induced unfolding and aggregation on PSE-like chicken protein edible film formation. <i>Food Chemistry</i> , 2020, 319, 126574. | 8.2 | 37 |
| 68 | Modification of myofibrillar protein via glycation: Physicochemical characterization, rheological behavior and solubility property. <i>Food Hydrocolloids</i> , 2020, 105, 105852. | 10.7 | 77 |
| 69 | Preparation, characterization, physicochemical property and potential application of porous starch: A review. <i>International Journal of Biological Macromolecules</i> , 2020, 148, 1169-1181. | 7.5 | 101 |
| 70 | A staining method for detection of <i>Enterocytozoon hepatopenaei</i> (EHP) spores with calcofluor white. <i>Journal of Invertebrate Pathology</i> , 2020, 172, 107347. | 3.2 | 15 |
| 71 | Physicochemical and microstructural attributes of marinated chicken breast influenced by breathing ultrasonic tumbling. <i>Ultrasonics Sonochemistry</i> , 2020, 64, 105022. | 8.2 | 28 |
| 72 | A TaqMan probe based real-time PCR for the detection of Decapod iridescent virus 1. <i>Journal of Invertebrate Pathology</i> , 2020, 173, 107367. | 3.2 | 31 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 73 | Overheating induced structural changes of type I collagen and impaired the protein digestibility. <i>Food Research International</i> , 2020, 134, 109225. | 6.2 | 47 |
| 74 | Effects of inulin on the gel properties and molecular structure of porcine myosin: A underlying mechanisms study. <i>Food Hydrocolloids</i> , 2020, 108, 105974. | 10.7 | 38 |
| 75 | Effects of ultrafine comminution treatment on gelling properties of myofibrillar proteins from chicken breast. <i>Food Hydrocolloids</i> , 2019, 97, 105199. | 10.7 | 43 |
| 76 | High-pressure homogenization combined with sulfhydryl blockage by hydrogen peroxide enhance the thermal stability of chicken breast myofibrillar protein aqueous solution. <i>Food Chemistry</i> , 2019, 285, 31-38. | 8.2 | 58 |
| 77 | Antioxidant activity of peptides in postmortem aged duck meat as affected by cooking and <i>in vitro</i> digestion. <i>International Journal of Food Properties</i> , 2019, 22, 727-736. | 3.0 | 14 |
| 78 | Impact of gum Arabic on the partition and stability of resveratrol in sunflower oil emulsions stabilized by whey protein isolate. <i>Colloids and Surfaces B: Biointerfaces</i> , 2019, 181, 749-755. | 5.0 | 27 |
| 79 | Description of a Natural Infection with Decapod Iridescent Virus 1 in Farmed Giant Freshwater Prawn, <i>Macrobrachium rosenbergii</i> . <i>Viruses</i> , 2019, 11, 354. | 3.3 | 74 |
| 80 | Susceptibility of <i>Exopalaemon carinicauda</i> to the Infection with Shrimp Hemocyte Iridescent Virus (SHIV 20141215), a Strain of Decapod Iridescent Virus 1 (DIV1). <i>Viruses</i> , 2019, 11, 387. | 3.3 | 52 |
| 81 | Isoelectric solubilization/precipitation processing modified sarcoplasmic protein from pale, soft, exudative-like chicken meat. <i>Food Chemistry</i> , 2019, 287, 1-10. | 8.2 | 15 |
| 82 | Effect of the disruption chamber geometry on the physicochemical and structural properties of water-soluble myofibrillar proteins prepared by high pressure homogenization (HPH). <i>LWT - Food Science and Technology</i> , 2019, 105, 215-223. | 5.2 | 17 |
| 83 | Myofibrillar protein-curcumin nanocomplexes prepared at different ionic strengths to improve oxidative stability of marinated chicken meat products. <i>LWT - Food Science and Technology</i> , 2019, 99, 69-76. | 5.2 | 29 |
| 84 | Structural changes and emulsion properties of goose liver proteins obtained by isoelectric solubilisation/precipitation processes. <i>LWT - Food Science and Technology</i> , 2019, 102, 190-196. | 5.2 | 28 |
| 85 | The case for thyroid disruption in early life stage exposures to thiram in zebrafish (<i>Danio rerio</i>). <i>General and Comparative Endocrinology</i> , 2019, 271, 73-81. | 1.8 | 24 |
| 86 | Oxidative stability of isoelectric solubilization/precipitation-isolated PSE-like chicken protein. <i>Food Chemistry</i> , 2019, 283, 646-655. | 8.2 | 24 |
| 87 | Structural and solubility properties of pale, soft and exudative (PSE)-like chicken breast myofibrillar protein: Effect of glycosylation. <i>LWT - Food Science and Technology</i> , 2018, 95, 209-215. | 5.2 | 36 |
| 88 | Effects of high-intensity ultrasound, high-pressure processing, and high-pressure homogenization on the physicochemical and functional properties of myofibrillar proteins. <i>Innovative Food Science and Emerging Technologies</i> , 2018, 45, 354-360. | 5.6 | 73 |
| 89 | Structural modification of myofibrillar proteins by high-pressure processing for functionally improved, value-added, and healthy muscle gelled foods. <i>Critical Reviews in Food Science and Nutrition</i> , 2018, 58, 2981-3003. | 10.3 | 80 |
| 90 | Applications of high pressure to pre-rigor rabbit muscles affect the water characteristics of myosin gels. <i>Food Chemistry</i> , 2018, 240, 59-66. | 8.2 | 28 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 91 | Effect of sodium alginate with three molecular weight forms on the water holding capacity of chicken breast myosin gel. <i>Food Chemistry</i> , 2018, 239, 1134-1142. | 8.2 | 81 |
| 92 | Chicken breast quality "normal, pale, soft and exudative (PSE) and woody" influences the functional properties of meat batters. <i>International Journal of Food Science and Technology</i> , 2018, 53, 654-664. | 2.7 | 36 |
| 93 | Rheological behavior, conformational changes and interactions of water-soluble myofibrillar protein during heating. <i>Food Hydrocolloids</i> , 2018, 77, 524-533. | 10.7 | 101 |
| 94 | Dose-dependent effects of rosmarinic acid on formation of oxidatively stressed myofibrillar protein emulsion gel at different NaCl concentrations. <i>Food Chemistry</i> , 2018, 243, 50-57. | 8.2 | 88 |
| 95 | Gelation properties of goose liver protein recovered by isoelectric solubilisation/precipitation process. <i>International Journal of Food Science and Technology</i> , 2018, 53, 356-364. | 2.7 | 12 |
| 96 | Thermal gelling properties and mechanism of porcine myofibrillar protein containing flaxseed gum at different NaCl concentrations. <i>LWT - Food Science and Technology</i> , 2018, 87, 361-367. | 5.2 | 61 |
| 97 | Alkaline pH-dependent thermal aggregation of chicken breast myosin: formation of soluble aggregates. <i>CYTA - Journal of Food</i> , 2018, 16, 765-775. | 1.9 | 17 |
| 98 | Inhibition of Heat-Induced Flocculation of Myosin-Based Emulsions through Steric Repulsion by Conformational Adaptation-Enhanced Interfacial Protein with an Alkaline pH-Shifting-Driven Method. <i>Langmuir</i> , 2018, 34, 8848-8856. | 3.5 | 10 |
| 99 | Manipulating interfacial behavior and emulsifying properties of myosin through alkali-heat treatment. <i>Food Hydrocolloids</i> , 2018, 85, 69-74. | 10.7 | 39 |
| 100 | Inhibition of Epigallocatechin-3-gallate/Protein Interaction by Methyl- β -cyclodextrin in Myofibrillar Protein Emulsion Gels under Oxidative Stress. <i>Journal of Agricultural and Food Chemistry</i> , 2018, 66, 8094-8103. | 5.2 | 30 |
| 101 | High-pressure effects on myosin in relation to heat gelation: A micro-perspective study. <i>Food Hydrocolloids</i> , 2018, 84, 219-228. | 10.7 | 14 |
| 102 | Effects of chicken myofibrillar protein concentration on protein oxidation and water holding capacity of its heat-induced gels. <i>Journal of Food Measurement and Characterization</i> , 2018, 12, 2302-2312. | 3.2 | 17 |
| 103 | Solubilization of myofibrillar proteins in water or low ionic strength media: Classical techniques, basic principles, and novel functionalities. <i>Critical Reviews in Food Science and Nutrition</i> , 2017, 57, 3260-3280. | 10.3 | 96 |
| 104 | Effect of Sodium Chloride on the Properties of Ready-to-Eat Pressure-Induced Gel-Type Chicken Meat Products. <i>Journal of Food Process Engineering</i> , 2017, 40, e12299. | 2.9 | 7 |
| 105 | Emulsifying Properties of Oxidatively Stressed Myofibrillar Protein Emulsion Gels Prepared with (α)-Epigallocatechin-3-gallate and NaCl. <i>Journal of Agricultural and Food Chemistry</i> , 2017, 65, 2816-2826. | 5.2 | 86 |
| 106 | Applications of high pressure to pre-rigor rabbit muscles affect the functional properties associated with heat-induced gelation. <i>Meat Science</i> , 2017, 129, 176-184. | 5.5 | 26 |
| 107 | Yield, thermal denaturation, and microstructure of proteins isolated from pale, soft, exudative chicken breast meat by using isoelectric solubilization/precipitation. <i>Process Biochemistry</i> , 2017, 58, 167-173. | 3.7 | 15 |
| 108 | Generation of bioactive peptides from duck meat during post-mortem aging. <i>Food Chemistry</i> , 2017, 237, 408-415. | 8.2 | 39 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Effects of high-pressure treatments on water characteristics and juiciness of rabbit meat sausages: Role of microstructure and chemical interactions. <i>Innovative Food Science and Emerging Technologies</i> , 2017, 41, 150-159. | 5.6 | 50 |
| 110 | Changes of Molecular Forces During Thermo-Gelling of Protein Isolated from PSE-Like Chicken Breast by Various Isoelectric Solubilization/Precipitation Extraction Strategies. <i>Food and Bioprocess Technology</i> , 2017, 10, 1240-1247. | 4.7 | 16 |
| 111 | High-pressure processing-induced conformational changes during heating affect water holding capacity of myosin gel. <i>International Journal of Food Science and Technology</i> , 2017, 52, 724-732. | 2.7 | 30 |
| 112 | In vitro protein digestibility of pork products is affected by the method of processing. <i>Food Research International</i> , 2017, 92, 88-94. | 6.2 | 92 |
| 113 | Changes in <i>in vitro</i> protein digestion of retort-pouched pork belly during 120-day storage. <i>International Journal of Food Science and Technology</i> , 2017, 52, 2684-2694. | 2.7 | 7 |
| 114 | Water-soluble myofibrillar proteins prepared by high-pressure homogenisation: a comparison study on the composition and functionality. <i>International Journal of Food Science and Technology</i> , 2017, 52, 2334-2342. | 2.7 | 11 |
| 115 | Stability of antioxidant peptides from duck meat after post-mortem ageing. <i>International Journal of Food Science and Technology</i> , 2017, 52, 2513-2521. | 2.7 | 7 |
| 116 | Influence of biofilm surface layer protein A (BslA) on the gel structure of myofibril protein from chicken breast. <i>Journal of the Science of Food and Agriculture</i> , 2017, 97, 4712-4720. | 3.5 | 14 |
| 117 | Effect of salt content on gelation of normal and wooden breast myopathy chicken pectoralis major meat batters. <i>International Journal of Food Science and Technology</i> , 2017, 52, 2068-2077. | 2.7 | 27 |
| 118 | High-pressure effects on the molecular aggregation and physicochemical properties of myosin in relation to heat gelation. <i>Food Research International</i> , 2017, 99, 413-418. | 6.2 | 17 |
| 119 | A comparative study of functional properties of normal and wooden breast broiler chicken meat with NaCl addition. <i>Poultry Science</i> , 2017, 96, 3473-3481. | 3.4 | 37 |
| 120 | Structural modification by high-pressure homogenization for improved functional properties of freeze-dried myofibrillar proteins powder. <i>Food Research International</i> , 2017, 100, 193-200. | 6.2 | 124 |
| 121 | Comparison of the Acidic and Alkaline Treatment on Emulsion Composite Gel Properties of the Proteins Recovered from Chicken Breast by Isoelectric Solubilization/Precipitation Process. <i>Journal of Food Processing and Preservation</i> , 2017, 41, e12884. | 2.0 | 7 |
| 122 | Precipitation and ultimate pH effect on chemical and gelation properties of protein prepared by isoelectric solubilization/precipitation process from pale, soft, exudative (PSE)-like chicken breast meat. <i>Poultry Science</i> , 2017, 96, 1504-1512. | 3.4 | 7 |
| 123 | L-histidine improves water retention of heat-induced gel of chicken breast myofibrillar proteins in low ionic strength solution. <i>International Journal of Food Science and Technology</i> , 2016, 51, 1195-1203. | 2.7 | 41 |
| 124 | Thermal gelling properties and mechanism of porcine myofibrillar protein containing flaxseed gum at various pH values. <i>CYTA - Journal of Food</i> , 2016, 14, 547-554. | 1.9 | 13 |
| 125 | Potential of high pressure homogenization to solubilize chicken breast myofibrillar proteins in water. <i>Innovative Food Science and Emerging Technologies</i> , 2016, 33, 170-179. | 5.6 | 131 |
| 126 | Conformational changes induced by high-pressure homogenization inhibit myosin filament formation in low ionic strength solutions. <i>Food Research International</i> , 2016, 85, 1-9. | 6.2 | 110 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 127 | Application of isoelectric solubilization/precipitation processing to improve gelation properties of protein isolated from pale, soft, exudative (PSE)-like chicken breast meat. <i>LWT - Food Science and Technology</i> , 2016, 72, 141-148. | 5.2 | 40 |
| 128 | Effects of sodium tripolyphosphate on functional properties of low-salt single-step high-pressure processed chicken breast sausage. <i>International Journal of Food Science and Technology</i> , 2016, 51, 2106-2113. | 2.7 | 12 |
| 129 | Comparative study of extraction efficiency and composition of protein recovered from chicken liver by acid-alkaline treatment. <i>Process Biochemistry</i> , 2016, 51, 1629-1635. | 3.7 | 23 |
| 130 | Different physicochemical, structural and digestibility characteristics of myofibrillar protein from PSE and normal pork before and after oxidation. <i>Meat Science</i> , 2016, 121, 228-237. | 5.5 | 35 |
| 131 | Optimization of textural properties of reduced-fat and reduced-salt emulsion-type sausages treated with high pressure using a response surface methodology. <i>Innovative Food Science and Emerging Technologies</i> , 2016, 33, 162-169. | 5.6 | 13 |
| 132 | Solubilisation of myosin in a solution of low ionic strength l-histidine: Significance of the imidazole ring. <i>Food Chemistry</i> , 2016, 196, 42-49. | 8.2 | 100 |
| 133 | Effects of Heat-oxidized Soy Protein Isolate on Growth Performance and Digestive Function of Broiler Chickens at Early Age. <i>Asian-Australasian Journal of Animal Sciences</i> , 2015, 28, 544-550. | 2.4 | 14 |
| 134 | High pressure/thermal combinations on texture and water holding capacity of chicken batters. <i>Innovative Food Science and Emerging Technologies</i> , 2015, 30, 8-14. | 5.6 | 56 |
| 135 | Effect of high pressure on cooking losses and functional properties of reduced-fat and reduced-salt pork sausage emulsions. <i>Innovative Food Science and Emerging Technologies</i> , 2015, 29, 125-133. | 5.6 | 38 |
| 136 | Effect of Cooking on <i>in Vitro</i> Digestion of Pork Proteins: A Peptidomic Perspective. <i>Journal of Agricultural and Food Chemistry</i> , 2015, 63, 250-261. | 5.2 | 88 |
| 137 | High pressure processing alters water distribution enabling the production of reduced-fat and reduced-salt pork sausages. <i>Meat Science</i> , 2015, 102, 69-78. | 5.5 | 59 |
| 138 | Effects of High-Pressure Processing on the Cooking Loss and Gel Strength of Chicken Breast Actomyosin Containing Sodium Alginate. <i>Food and Bioprocess Technology</i> , 2014, 7, 3608-3617. | 4.7 | 41 |
| 139 | Contribution of Three Ionic Types of Polysaccharides to the Thermal Gelling Properties of Chicken Breast Myosin. <i>Journal of Agricultural and Food Chemistry</i> , 2014, 62, 2655-2662. | 5.2 | 50 |
| 140 | Effect of pH on heat-induced gelation of duck blood plasma protein. <i>Food Hydrocolloids</i> , 2014, 35, 324-331. | 10.7 | 26 |
| 141 | Effects of high pressure processing on the thermal gelling properties of chicken breast myosin containing I ^κ -carrageenan. <i>Food Hydrocolloids</i> , 2014, 40, 262-272. | 10.7 | 131 |
| 142 | Low-field NMR study of heat-induced gelation of pork myofibrillar proteins and its relationship with microstructural characteristics. <i>Food Research International</i> , 2014, 62, 1175-1182. | 6.2 | 298 |
| 143 | The mechanism of high pressure-induced gels of rabbit myosin. <i>Innovative Food Science and Emerging Technologies</i> , 2012, 16, 41-46. | 5.6 | 130 |
| 144 | Influence of Various Levels of Flaxseed Gum Addition on the Water Holding Capacities of Heat-Induced Porcine Myofibrillar Protein. <i>Journal of Food Science</i> , 2011, 76, C472-8. | 3.1 | 68 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | Effect of microbial transglutaminase on NMR relaxometry and microstructure of pork myofibrillar protein gel. <i>European Food Research and Technology</i> , 2009, 228, 665-670. | 3.3 | 157 |
| 146 | Influence of caspase3 selective inhibitor on proteolysis of chicken skeletal muscle proteins during post mortem aging. <i>Food Chemistry</i> , 2009, 115, 181-186. | 8.2 | 56 |
| 147 | Changes of intramuscular phospholipids and free fatty acids during the processing of Nanjing dry-cured duck. <i>Food Chemistry</i> , 2008, 110, 279-284. | 8.2 | 67 |