

# Tomoyuki Nagata

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6573983/publications.pdf>

Version: 2024-02-01

39  
papers

602  
citations

623734

14  
h-index

642732

23  
g-index

39  
all docs

39  
docs citations

39  
times ranked

975  
citing authors

#	ARTICLE	IF	CITATIONS
1	Association between DNA Methylation of the BDNF Promoter Region and Clinical Presentation in Alzheimer's Disease. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2015, 5, 64-73.	1.3	65
2	Increase in the IgG avidity index due to herpes simplex virus type 1 reactivation and its relationship with cognitive function in amnesic mild cognitive impairment and Alzheimer's disease. <i>Biochemical and Biophysical Research Communications</i> , 2013, 430, 907-911.	2.1	62
3	Plasma BDNF levels are correlated with aggressiveness in patients with amnesic mild cognitive impairment or Alzheimer disease. <i>Journal of Neural Transmission</i> , 2014, 121, 433-441.	2.8	36
4	Association between executive dysfunction and hippocampal volume in Alzheimer's disease. <i>International Psychogeriatrics</i> , 2011, 23, 764-771.	1.0	33
5	Association between BDNF Polymorphism (Val66Met) and Executive Function in Patients with Amnesic Mild Cognitive Impairment or Mild Alzheimer Disease. <i>Dementia and Geriatric Cognitive Disorders</i> , 2012, 33, 266-272.	1.5	33
6	Development of Biomarkers Based on DNA Methylation in the NCAPH2/LMF2 Promoter Region for Diagnosis of Alzheimer's Disease and Amnesic Mild Cognitive Impairment. <i>PLoS ONE</i> , 2016, 11, e0146449.	2.5	29
7	Psychosocial or clinico-demographic factors related to neuropsychiatric symptoms in patients with Alzheimer's disease needing interventional treatment: analysis of the CATIE-AD study. <i>International Journal of Geriatric Psychiatry</i> , 2017, 32, 1264-1271.	2.7	27
8	Relationship of frontal lobe dysfunction and aberrant motor behaviors in patients with Alzheimer's disease. <i>International Psychogeriatrics</i> , 2010, 22, 463-469.	1.0	26
9	Usefulness of DNA Methylation Levels in COASY and SPINT1 Gene Promoter Regions as Biomarkers in Diagnosis of Alzheimer's Disease and Amnesic Mild Cognitive Impairment. <i>PLoS ONE</i> , 2016, 11, e0168816.	2.5	26
10	Correlation between a reduction in Frontal Assessment Battery scores and delusional thoughts in patients with Alzheimer's disease. <i>Psychiatry and Clinical Neurosciences</i> , 2009, 63, 449-454.	1.8	25
11	A case in which mirtazapine reduced auditory hallucinations in a patient with Parkinson disease. <i>International Psychogeriatrics</i> , 2013, 25, 1199-1201.	1.0	21
12	Classification of Neuropsychiatric Symptoms Requiring Antipsychotic Treatment in Patients with Alzheimer's Disease: Analysis of the CATIE-AD Study. <i>Journal of Alzheimer's Disease</i> , 2016, 50, 839-845.	2.6	19
13	Differentiation between amnesic mild cognitive impairment and early-stage Alzheimer's disease using the Frontal Assessment Battery test. <i>Psychogeriatrics</i> , 2011, 11, 235-241.	1.2	17
14	Increased blood COASY DNA methylation levels a potential biomarker for early pathology of Alzheimer's disease. <i>Scientific Reports</i> , 2020, 10, 12217.	3.3	17
15	Association between brain-derived neurotrophic factor ( <i>BDNF</i> ) gene polymorphisms and executive function in Japanese patients with Alzheimer's disease. <i>Psychogeriatrics</i> , 2011, 11, 141-149.	1.2	15
16	Anosognosia in patients with Alzheimer's disease: current perspectives. <i>Psychogeriatrics</i> , 2020, 20, 345-352.	1.2	14
17	Association between Neuropsychiatric Improvement and Neurocognitive Change in Alzheimer's Disease: Analysis of the CATIE-AD Study. <i>Journal of Alzheimer's Disease</i> , 2018, 66, 139-148.	2.6	13
18	Effectiveness of carbamazepine for benzodiazepine-resistant impulsive aggression in a patient with frontal infarctions. <i>Psychiatry and Clinical Neurosciences</i> , 2007, 61, 695-697.	1.8	10

#	ARTICLE	IF	CITATIONS
19	Association between Nerve Growth Factor Gene Polymorphism and Executive Dysfunction in Japanese Patients with Early-Stage Alzheimer's Disease and Amnesic Mild Cognitive Impairment. <i>Dementia and Geriatric Cognitive Disorders</i> , 2011, 32, 379-386.	1.5	10
20	Anosognosia in mild Alzheimer's disease is correlated with not only neural dysfunction but also compensation. <i>Psychogeriatrics</i> , 2018, 18, 81-88.	1.2	10
21	Cognitive Dysfunction in Patients With Late-Life Somatic Symptom Disorder: A Comparison According to Disease Severity. <i>Psychosomatics</i> , 2015, 56, 486-494.	2.5	9
22	Sex differences in the severity of neuropsychiatric symptoms and their relationship with clinico-demographic and psychosocial factors in patients with amnesic mild cognitive impairment and mild Alzheimer's disease. <i>Aging and Mental Health</i> , 2020, 24, 431-438.	2.8	8
23	Effects of neuropsychiatric symptoms of dementia on reductions in activities of daily living in patients with Alzheimer's disease. <i>Geriatrics and Gerontology International</i> , 2020, 20, 584-588.	1.5	8
24	Genetic Association between Neurotrophin-3 Polymorphisms and Alzheimer's Disease in Japanese Patients. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2013, 3, 272-280.	1.3	7
25	Baseline Predictors of Antipsychotic Treatment Continuation and Response at Week 8 in Patients with Alzheimer's Disease with Psychosis or Aggressive Symptoms: An Analysis of the CATIE-AD Study. <i>Journal of Alzheimer's Disease</i> , 2017, 60, 263-272.	2.6	7
26	Pharmacological management of behavioral disturbances in patients with Alzheimer's disease. <i>Expert Opinion on Pharmacotherapy</i> , 2020, 21, 1093-1102.	1.8	7
27	Early Improvements of Individual Symptoms With Antipsychotics Predict Subsequent Treatment Response of Neuropsychiatric Symptoms in Alzheimer's Disease. <i>Journal of Clinical Psychiatry</i> , 2020, 81, .	2.2	6
28	Suppressors of Cytokine Signaling Are Decreased in Major Depressive Disorder Patients. <i>Journal of Personalized Medicine</i> , 2022, 12, 1040.	2.5	6
29	Genetic association between <i>RAGE</i> polymorphisms and Alzheimer's disease and Lewy body dementias in a Japanese cohort: a case-control study. <i>International Journal of Geriatric Psychiatry</i> , 2017, 32, 1241-1246.	2.7	5
30	A case of V180I genetic mutation Creutzfeldt Jakob disease (CJD) with delusional misidentification as an initial symptom. <i>Prion</i> , 2022, 16, 7-13.	1.8	5
31	Anorexia nervosa with chronic episodes for more than 30 years in a patient with a comorbid schizotypal personality disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2007, 61, 434-436.	1.8	4
32	Association between the catechol-O-methyltransferase polymorphism Val158Met and Alzheimer's disease in a Japanese population. <i>International Journal of Geriatric Psychiatry</i> , 2015, 30, 927-933.	2.7	4
33	Correlation between cognition and symptomatic severity in patients with late-life somatoform disorders. <i>Aging and Mental Health</i> , 2015, 19, 169-174.	2.8	4
34	Executive Dysfunction Correlated With 2-Year Treatment Response in Patients With Late-Life Undifferentiated Somatoform Disorders. <i>Psychosomatics</i> , 2016, 57, 378-389.	2.5	3
35	Blood DNA Methylation Levels in the WNT5A Gene Promoter Region: A Potential Biomarker for Agitation in Subjects with Dementia. <i>Journal of Alzheimer's Disease</i> , 2021, 81, 1601-1611.	2.6	3
36	Education level is associated with neuropsychiatric symptoms in patients with amnesic mild cognitive impairment. <i>Psychogeriatrics</i> , 2022, 22, 343-352.	1.2	3

#	ARTICLE	IF	CITATIONS
37	Age-Related Association between Apolipoprotein E e4 and Cognitive Function in Japanese Patients with Alzheimer's Disease. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2013, 3, 66-73.	1.3	2
38	Genetic Association between Presenilin 2 Polymorphisms and Alzheimer's Disease and Dementia of Lewy Body Type in a Japanese Population. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2016, 6, 90-97.	1.3	2
39	The time-dependent trajectory of neuropsychiatric symptoms in patients with Alzheimer's disease. <i>Psychogeriatrics</i> , 2020, 20, 542-543.	1.2	1