

# Tomoyuki Nagata

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6573983/publications.pdf>

Version: 2024-02-01

39  
papers

602  
citations

623734

14  
h-index

642732

23  
g-index

39  
all docs

39  
docs citations

39  
times ranked

975  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Education level is associated with neuropsychiatric symptoms in patients with amnesicâ€mild cognitive impairment. <i>Psychogeriatrics</i> , 2022, 22, 343-352.   | 1.2 | 3         |
| 2  | A case of V180I genetic mutation Creutzfeldt Jakob disease (CJD) with delusional misidentification as an initial symptom. <i>Prion</i> , 2022, 16, 7-13.   | 1.8 | 5         |
| 3  | Suppressors of Cytokine Signaling Are Decreased in Major Depressive Disorder Patients. <i>Journal of Personalized Medicine</i> , 2022, 12, 1040.   | 2.5 | 6         |
| 4  | Blood DNA Methylation Levels in the WNT5A Gene Promoter Region: A Potential Biomarker for Agitation in Subjects with Dementia. <i>Journal of Alzheimer's Disease</i> , 2021, 81, 1601-1611.  | 2.6 | 3         |
| 5  | Sex differences in the severity of neuropsychiatric symptoms and their relationship with clinico-demographic and psychosocial factors in patients with amnesic mild cognitive impairment and mild Alzheimerâ€™s disease. <i>Aging and Mental Health</i> , 2020, 24, 431-438. | 2.8 | 8         |
| 6  | Increased blood COASY DNA methylation levels a potential biomarker for early pathology of Alzheimerâ€™s disease. <i>Scientific Reports</i> , 2020, 10, 12217.  | 3.3 | 17        |
| 7  | The timeâ€dependent trajectory of neuropsychiatric symptoms in patients with Alzheimer's disease. <i>Psychogeriatrics</i> , 2020, 20, 542-543.   | 1.2 | 1         |
| 8  | Early Improvements of Individual Symptoms With Antipsychotics Predict Subsequent Treatment Response of Neuropsychiatric Symptoms in Alzheimerâ€™s Disease. <i>Journal of Clinical Psychiatry</i> , 2020, 81, .   | 2.2 | 6         |
| 9  | Anosognosia in patients with Alzheimer's disease: current perspectives. <i>Psychogeriatrics</i> , 2020, 20, 345-352.   | 1.2 | 14        |
| 10 | Effects of neuropsychiatric symptoms of dementia on reductions in activities of daily living in patients with Alzheimer's disease. <i>Geriatrics and Gerontology International</i> , 2020, 20, 584-588.  | 1.5 | 8         |
| 11 | Pharmacological management of behavioral disturbances in patients with Alzheimerâ€™s disease. <i>Expert Opinion on Pharmacotherapy</i> , 2020, 21, 1093-1102.  | 1.8 | 7         |
| 12 | Anosognosia in mild <scp>A</scp>lzheimer's disease is correlated with not only neural dysfunction but also compensation. <i>Psychogeriatrics</i> , 2018, 18, 81-88.  | 1.2 | 10        |
| 13 | Association between Neuropsychiatric Improvement and Neurocognitive Change in Alzheimerâ€™s Disease: Analysis of the CATIE-AD Study. <i>Journal of Alzheimer's Disease</i> , 2018, 66, 139-148.  | 2.6 | 13        |
| 14 | Genetic association between <i>RAGE</i> polymorphisms and Alzheimer's disease and Lewy body dementias in a Japanese cohort: a caseâ€control study. <i>International Journal of Geriatric Psychiatry</i> , 2017, 32, 1241-1246.   | 2.7 | 5         |
| 15 | Psychosocial or clinicoâ€demographic factors related to neuropsychiatric symptoms in patients with Alzheimer's disease needing interventional treatment: analysis of the CATIEâ€AD study. <i>International Journal of Geriatric Psychiatry</i> , 2017, 32, 1264-1271.        | 2.7 | 27        |
| 16 | Baseline Predictors of Antipsychotic Treatment Continuation and Response at Week 8 in Patients with Alzheimerâ€™s Disease with Psychosis or Aggressive Symptoms: An Analysis of the CATIE-AD Study. <i>Journal of Alzheimer's Disease</i> , 2017, 60, 263-272.               | 2.6 | 7         |
| 17 | Genetic Association between Presenilin 2 Polymorphisms and Alzheimer's Disease and Dementia of Lewy Body Type in a Japanese Population. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2016, 6, 90-97.  | 1.3 | 2         |
| 18 | Development of Biomarkers Based on DNA Methylation in the NCAPH2/LMF2 Promoter Region for Diagnosis of Alzheimerâ€™s Disease and Amnesic Mild Cognitive Impairment. <i>PLoS ONE</i> , 2016, 11, e0146449.  | 2.5 | 29        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Usefulness of DNA Methylation Levels in COASY and SPINT1 Gene Promoter Regions as Biomarkers in Diagnosis of Alzheimer's Disease and Amnestic Mild Cognitive Impairment. <i>PLoS ONE</i> , 2016, 11, e0168816.   | 2.5 | 26        |
| 20 | Classification of Neuropsychiatric Symptoms Requiring Antipsychotic Treatment in Patients with Alzheimer's Disease: Analysis of the CATIE-AD Study. <i>Journal of Alzheimer's Disease</i> , 2016, 50, 839-845.   | 2.6 | 19        |
| 21 | Executive Dysfunction Correlated With 2-Year Treatment Response in Patients With Late-Life Undifferentiated Somatoform Disorders. <i>Psychosomatics</i> , 2016, 57, 378-389.   | 2.5 | 3         |
| 22 | Cognitive Dysfunction in Patients With Late-Life Somatic Symptom Disorder: A Comparison According to Disease Severity. <i>Psychosomatics</i> , 2015, 56, 486-494.  | 2.5 | 9         |
| 23 | Association between DNA Methylation of the BDNF Promoter Region and Clinical Presentation in Alzheimer's Disease. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2015, 5, 64-73.  | 1.3 | 65        |
| 24 | Association between the catechol-O-methyltransferase polymorphism Val158Met and Alzheimer's disease in a Japanese population. <i>International Journal of Geriatric Psychiatry</i> , 2015, 30, 927-933.  | 2.7 | 4         |
| 25 | Correlation between cognition and symptomatic severity in patients with late-life somatoform disorders. <i>Aging and Mental Health</i> , 2015, 19, 169-174.  | 2.8 | 4         |
| 26 | Plasma BDNF levels are correlated with aggressiveness in patients with amnestic mild cognitive impairment or Alzheimer disease. <i>Journal of Neural Transmission</i> , 2014, 121, 433-441.  | 2.8 | 36        |
| 27 | Increase in the IgG avidity index due to herpes simplex virus type 1 reactivation and its relationship with cognitive function in amnestic mild cognitive impairment and Alzheimer's disease. <i>Biochemical and Biophysical Research Communications</i> , 2013, 430, 907-911. | 2.1 | 62        |
| 28 | Age-Related Association between Apolipoprotein E e4 and Cognitive Function in Japanese Patients with Alzheimer's Disease. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2013, 3, 66-73.  | 1.3 | 2         |
| 29 | A case in which mirtazapine reduced auditory hallucinations in a patient with Parkinson disease. <i>International Psychogeriatrics</i> , 2013, 25, 1199-1201.  | 1.0 | 21        |
| 30 | Genetic Association between Neurotrophin-3 Polymorphisms and Alzheimer's Disease in Japanese Patients. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2013, 3, 272-280.   | 1.3 | 7         |
| 31 | Association between BDNF Polymorphism (Val66Met) and Executive Function in Patients with Amnestic Mild Cognitive Impairment or Mild Alzheimer Disease. <i>Dementia and Geriatric Cognitive Disorders</i> , 2012, 33, 266-272.  | 1.5 | 33        |
| 32 | Association between Nerve Growth Factor Gene Polymorphism and Executive Dysfunction in Japanese Patients with Early-Stage Alzheimer's Disease and Amnestic Mild Cognitive Impairment. <i>Dementia and Geriatric Cognitive Disorders</i> , 2011, 32, 379-386.                   | 1.5 | 10        |
| 33 | Association between executive dysfunction and hippocampal volume in Alzheimer's disease. <i>International Psychogeriatrics</i> , 2011, 23, 764-771.  | 1.0 | 33        |
| 34 | Association between brain-derived neurotrophic factor (BDNF) gene polymorphisms and executive function in Japanese patients with Alzheimer's disease. <i>Psychogeriatrics</i> , 2011, 11, 141-149.   | 1.2 | 15        |
| 35 | Differentiation between amnestic mild cognitive impairment and early-stage Alzheimer's disease using the Frontal Assessment Battery test. <i>Psychogeriatrics</i> , 2011, 11, 235-241.   | 1.2 | 17        |
| 36 | Relationship of frontal lobe dysfunction and aberrant motor behaviors in patients with Alzheimer's disease. <i>International Psychogeriatrics</i> , 2010, 22, 463-469.   | 1.0 | 26        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Correlation between a reduction in Frontal Assessment Battery scores and delusional thoughts in patients with Alzheimer's disease. <i>Psychiatry and Clinical Neurosciences</i> , 2009, 63, 449-454. | 1.8 | 25        |
| 38 | Anorexia nervosa with chronic episodes for more than 30 years in a patient with a comorbid schizotypal personality disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2007, 61, 434-436.       | 1.8 | 4         |
| 39 | Effectiveness of carbamazepine for benzodiazepine-resistant impulsive aggression in a patient with frontal infarctions. <i>Psychiatry and Clinical Neurosciences</i> , 2007, 61, 695-697.            | 1.8 | 10        |