## Asif Ahmed

## List of Publications by Citations

Source: https://exaly.com/author-pdf/6585448/asif-ahmed-publications-by-citations.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

143 papers

8,157 citations

49 h-index

87 g-index

165 ext. papers

8,965 ext. citations

6.7 avg, IF

5.76 L-index

#	Paper	IF	Citations
143	Elevated placental soluble vascular endothelial growth factor receptor-1 inhibits angiogenesis in preeclampsia. <i>Circulation Research</i> , <b>2004</b> , 95, 884-91	15.7	400
142	Negative regulation of soluble Flt-1 and soluble endoglin release by heme oxygenase-1. <i>Circulation</i> , <b>2007</b> , 115, 1789-97	16.7	338
141	Fetal growth restriction results in remodeled and less efficient hearts in children. <i>Circulation</i> , <b>2010</b> , 121, 2427-36	16.7	289
140	Regulation of placental vascular endothelial growth factor (VEGF) and placenta growth factor (PIGF) and soluble Flt-1 by oxygena review. <i>Placenta</i> , <b>2000</b> , 21 Suppl A, S16-24	3.4	246
139	Vascular endothelial growth factor receptor-1 modulates vascular endothelial growth factor-mediated angiogenesis via nitric oxide. <i>American Journal of Pathology</i> , <b>2001</b> , 159, 993-1008	5.8	242
138	Vascular endothelial growth factor receptor localization and activation in human trophoblast and choriocarcinoma cells. <i>Biology of Reproduction</i> , <b>1994</b> , 51, 524-30	3.9	214
137	Colocalisation of vascular endothelial growth factor and its Flt-1 receptor in human placenta. <i>Growth Factors</i> , <b>1995</b> , 12, 235-43	1.6	213
136	Carbon monoxide expedites metabolic exhaustion to inhibit tumor growth. <i>Cancer Research</i> , <b>2013</b> , 73, 7009-21	10.1	210
135	Dysregulation of hydrogen sulfide producing enzyme cystathionine Eyase contributes to maternal hypertension and placental abnormalities in preeclampsia. <i>Circulation</i> , <b>2013</b> , 127, 2514-22	16.7	195
134	Cardiac dysfunction and cell damage across clinical stages of severity in growth-restricted fetuses. <i>American Journal of Obstetrics and Gynecology</i> , <b>2008</b> , 199, 254.e1-8	6.4	191
133	Release and complex formation of soluble VEGFR-1 from endothelial cells and biological fluids. <i>Laboratory Investigation</i> , <b>2000</b> , 80, 443-54	5.9	182
132	Bifunctional role for VEGF-induced heme oxygenase-1 in vivo: induction of angiogenesis and inhibition of leukocytic infiltration. <i>Blood</i> , <b>2004</b> , 103, 761-6	2.2	166
131	Localisation of placenta growth factor (PIGF) in human term placenta. <i>Growth Factors</i> , <b>1996</b> , 13, 243-50,color plates I-II,pre.bk cov	1.6	158
130	Angiopoietin-1 and angiopoietin-2 activate trophoblast Tie-2 to promote growth and migration during placental development. <i>American Journal of Pathology</i> , <b>2000</b> , 156, 2185-99	5.8	150
129	Autoantibody from women with preeclampsia induces soluble Fms-like tyrosine kinase-1 production via angiotensin type 1 receptor and calcineurin/nuclear factor of activated T-cells signaling. <i>Hypertension</i> , <b>2008</b> , 51, 1010-9	8.5	142
128	Reduction of circulating soluble Flt-1 alleviates preeclampsia-like symptoms in a mouse model. <i>Journal of Cellular and Molecular Medicine</i> , <b>2010</b> , 14, 1857-67	5.6	134
127	Induction of Placental Heme Oxygenase-1 Is Protective Against TNF⊞nduced Cytotoxicity and Promotes Vessel Relaxation. <i>Molecular Medicine</i> , <b>2000</b> , 6, 391-409	6.2	130

126	Establishment of porcine and human expanded potential stem cells. Nature Cell Biology, 2019, 21, 687-6	5 <b>2</b> 9.4	127
125	VEGF induces Tie2 shedding via a phosphoinositide 3-kinase/Akt dependent pathway to modulate Tie2 signaling. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , <b>2007</b> , 27, 2619-26	9.4	127
124	Angiotensin II induces soluble fms-Like tyrosine kinase-1 release via calcineurin signaling pathway in pregnancy. <i>Circulation Research</i> , <b>2007</b> , 100, 88-95	15.7	124
123	Direct evidence for endothelial vascular endothelial growth factor receptor-1 function in nitric oxide-mediated angiogenesis. <i>Circulation Research</i> , <b>2006</b> , 99, 715-22	15.7	119
122	Flt1 acts as a negative regulator of tip cell formation and branching morphogenesis in the zebrafish embryo. <i>Development (Cambridge)</i> , <b>2011</b> , 138, 2111-20	6.6	110
121	Activation of vascular endothelial growth factor receptor-1 sustains angiogenesis and Bcl-2 expression via the phosphatidylinositol 3-kinase pathway in endothelial cells. <i>Diabetes</i> , <b>2003</b> , 52, 2959-6	58 <sup>.9</sup>	108
120	The role of H2S bioavailability in endothelial dysfunction. <i>Trends in Pharmacological Sciences</i> , <b>2015</b> , 36, 568-78	13.2	106
119	Immunolocalisation of vascular endothelial growth factor in human endometrium. <i>Growth Factors</i> , <b>1994</b> , 11, 277-82	1.6	103
118	New insights into the etiology of preeclampsia: identification of key elusive factors for the vascular complications. <i>Thrombosis Research</i> , <b>2011</b> , 127 Suppl 3, S72-5	8.2	100
117	Hypoxia induces dilated cardiomyopathy in the chick embryo: mechanism, intervention, and		
,	long-term consequences. <i>PLoS ONE</i> , <b>2009</b> , 4, e5155	3.7	95
116	Hemeoxygenase-1 inhibits human myometrial contractility via carbon monoxide and is upregulated by progesterone during pregnancy. <i>Journal of Clinical Investigation</i> , <b>1998</b> , 101, 949-55	3·7 15.9	95
	Hemeoxygenase-1 inhibits human myometrial contractility via carbon monoxide and is upregulated		
116	Hemeoxygenase-1 inhibits human myometrial contractility via carbon monoxide and is upregulated by progesterone during pregnancy. <i>Journal of Clinical Investigation</i> , <b>1998</b> , 101, 949-55  The role of heterodimerization between VEGFR-1 and VEGFR-2 in the regulation of endothelial cell	15.9	94
116	Hemeoxygenase-1 inhibits human myometrial contractility via carbon monoxide and is upregulated by progesterone during pregnancy. <i>Journal of Clinical Investigation</i> , <b>1998</b> , 101, 949-55  The role of heterodimerization between VEGFR-1 and VEGFR-2 in the regulation of endothelial cell homeostasis. <i>Nature Communications</i> , <b>2012</b> , 3, 972  Nitric oxide-dependent bone marrow progenitor mobilization by carbon monoxide enhances	15.9 17.4	94
116 115 114	Hemeoxygenase-1 inhibits human myometrial contractility via carbon monoxide and is upregulated by progesterone during pregnancy. <i>Journal of Clinical Investigation</i> , <b>1998</b> , 101, 949-55  The role of heterodimerization between VEGFR-1 and VEGFR-2 in the regulation of endothelial cell homeostasis. <i>Nature Communications</i> , <b>2012</b> , 3, 972  Nitric oxide-dependent bone marrow progenitor mobilization by carbon monoxide enhances endothelial repair after vascular injury. <i>Circulation</i> , <b>2010</b> , 121, 537-48  Cellular localization of AT1 receptor mRNA and protein in normal placenta and its reduced expression in intrauterine growth restriction. Angiotensin II stimulates the release of	15.9 17.4 16.7	94 92 91
116 115 114	Hemeoxygenase-1 inhibits human myometrial contractility via carbon monoxide and is upregulated by progesterone during pregnancy. <i>Journal of Clinical Investigation</i> , <b>1998</b> , 101, 949-55  The role of heterodimerization between VEGFR-1 and VEGFR-2 in the regulation of endothelial cell homeostasis. <i>Nature Communications</i> , <b>2012</b> , 3, 972  Nitric oxide-dependent bone marrow progenitor mobilization by carbon monoxide enhances endothelial repair after vascular injury. <i>Circulation</i> , <b>2010</b> , 121, 537-48  Cellular localization of AT1 receptor mRNA and protein in normal placenta and its reduced expression in intrauterine growth restriction. Angiotensin II stimulates the release of vasorelaxants. <i>Journal of Clinical Investigation</i> , <b>1998</b> , 101, 442-54	15.9 17.4 16.7	94 92 91 91 90
116 115 114 113	Hemeoxygenase-1 inhibits human myometrial contractility via carbon monoxide and is upregulated by progesterone during pregnancy. <i>Journal of Clinical Investigation</i> , <b>1998</b> , 101, 949-55  The role of heterodimerization between VEGFR-1 and VEGFR-2 in the regulation of endothelial cell homeostasis. <i>Nature Communications</i> , <b>2012</b> , 3, 972  Nitric oxide-dependent bone marrow progenitor mobilization by carbon monoxide enhances endothelial repair after vascular injury. <i>Circulation</i> , <b>2010</b> , 121, 537-48  Cellular localization of AT1 receptor mRNA and protein in normal placenta and its reduced expression in intrauterine growth restriction. Angiotensin II stimulates the release of vasorelaxants. <i>Journal of Clinical Investigation</i> , <b>1998</b> , 101, 442-54  Dual role of angiotensin II in the human endometrium. <i>Human Reproduction</i> , <b>1996</b> , 11 Suppl 2, 95-108	15.9 17.4 16.7 15.9	94 92 91 91

108	Vascular endothelial growth factor-induced endothelial cell proliferation is regulated by interaction between VEGFR-2, SH-PTP1 and eNOS. <i>Microvascular Research</i> , <b>2006</b> , 71, 20-31	3.7	76
107	Immunohistochemical localization of vascular endothelial growth factor (VEGF) and its two specific receptors, Flt-1 and KDR, in the porcine placenta and non-pregnant uterus. <i>Placenta</i> , <b>1999</b> , 20, 35-43	3.4	69
106	Small Molecule, Big Prospects: MicroRNA in Pregnancy and Its Complications. <i>Journal of Pregnancy</i> , <b>2017</b> , 2017, 6972732	2.5	68
105	Antiproliferative effects of carbon monoxide on pancreatic cancer. <i>Digestive and Liver Disease</i> , <b>2014</b> , 46, 369-75	3.3	65
104	Hepatitis C virus infection reduces hepatocellular polarity in a vascular endothelial growth factor-dependent manner. <i>Gastroenterology</i> , <b>2010</b> , 138, 1134-42	13.3	62
103	Colocalization of acidic and basic fibroblast growth factor (FGF) in human placenta and the cellular effects of bFGF in trophoblast cell line JEG-3. <i>Growth Factors</i> , <b>1994</b> , 10, 259-68	1.6	60
102	Therapeutic potential of statins and the induction of heme oxygenase-1 in preeclampsia. <i>Journal of Reproductive Immunology</i> , <b>2014</b> , 101-102, 153-160	4.2	54
101	Ontogeny of hepatocyte growth factor (HGF) and its receptor (c-met) in human placenta: reduced HGF expression in intrauterine growth restriction. <i>American Journal of Pathology</i> , <b>1998</b> , 153, 1139-47	5.8	54
100	Evidence-Based Revised View of the Pathophysiology of Preeclampsia. <i>Advances in Experimental Medicine and Biology</i> , <b>2017</b> , 956, 355-374	3.6	53
99	Down-regulation of angiopoietin-1 expression in menorrhagia. <i>American Journal of Pathology</i> , <b>2002</b> , 160, 773-80	5.8	53
98	Induction of placental heme oxygenase-1 is protective against TNFalpha-induced cytotoxicity and promotes vessel relaxation. <i>Molecular Medicine</i> , <b>2000</b> , 6, 391-409	6.2	52
97	Localization of the angiotensin II and its receptor subtype expression in human endometrium and identification of a novel high-affinity angiotensin II binding site. <i>Journal of Clinical Investigation</i> , <b>1995</b> , 96, 848-57	15.9	51
96	Unravelling the theories of pre-eclampsia: are the protective pathways the new paradigm?. <i>British Journal of Pharmacology</i> , <b>2015</b> , 172, 1574-86	8.6	50
95	Vascular endothelial growth factor promotes physical wound repair and is anti-apoptotic in primary distal lung epithelial and A549 cells. <i>Critical Care Medicine</i> , <b>2007</b> , 35, 2164-70	1.4	50
94	Hypoxia or hyperoxia in placental insufficiency?. Lancet, The, 1997, 350, 826-7	40	48
93	Activation of functional oxytocin receptors stimulates cell proliferation in human trophoblast and choriocarcinoma cell lines. <i>Endocrinology</i> , <b>2001</b> , 142, 1130-6	4.8	48
92	Parathyroid hormone-related protein in human term placenta and membranes. <i>Journal of Endocrinology</i> , <b>1994</b> , 142, 217-24	4.7	48
91	Autocrine activity of soluble Flt-1 controls endothelial cell function and angiogenesis. <i>Vascular Cell</i> , <b>2011</b> , 3, 15	1	47

## (2001-2009)

90	Can the biology of VEGF and haem oxygenases help solve pre-eclampsia?. <i>Biochemical Society Transactions</i> , <b>2009</b> , 37, 1237-42	5.1	47
89	Localization of mRNA for basic fibroblast growth factor in human placenta. <i>Growth Factors</i> , <b>1994</b> , 11, 105-11	1.6	47
88	Loss of Akt activity increases circulating soluble endoglin release in preeclampsia: identification of inter-dependency between Akt-1 and heme oxygenase-1. <i>European Heart Journal</i> , <b>2012</b> , 33, 1150-8	9.5	46
87	Resveratrol inhibits the release of soluble fms-like tyrosine kinase (sFlt-1) from human placenta. <i>American Journal of Obstetrics and Gynecology</i> , <b>2012</b> , 206, 253.e10-5	6.4	46
86	Physical and biomechanical characteristics of rat cervical ripening are not consistent with increased collagenase activity. <i>American Journal of Obstetrics and Gynecology</i> , <b>2004</b> , 191, 1695-704	6.4	46
85	Pravastatin for early-onset pre-eclampsia: a randomised, blinded, placebo-controlled trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , <b>2020</b> , 127, 478-488	3.7	44
84	Basic FGF activates phospholipase D in endothelial cells in the absence of inositol-lipid hydrolysis. <i>American Journal of Physiology - Cell Physiology</i> , <b>1994</b> , 266, C206-12	5.4	42
83	Angiopoietin-2 confers Atheroprotection in apoE-/- mice by inhibiting LDL oxidation via nitric oxide. <i>Circulation Research</i> , <b>2009</b> , 104, 1333-6	15.7	40
82	Heparin elevates circulating soluble fms-like tyrosine kinase-1 immunoreactivity in pregnant women receiving anticoagulation therapy. <i>Circulation</i> , <b>2011</b> , 124, 2543-53	16.7	40
81	Localization, quantification, and activation of platelet-activating factor receptor in human endometrium during the menstrual cycle: PAF stimulates NO, VEGF, and FAKpp125. <i>FASEB Journal</i> , <b>1998</b> , 12, 831-43	0.9	40
80	Role of VEGF receptor-1 (Flt-1) in mediating calcium-dependent nitric oxide release and limiting DNA synthesis in human trophoblast cells. <i>Laboratory Investigation</i> , <b>1997</b> , 76, 779-91	5.9	40
79	Increased expression of placenta growth factor in proliferative diabetic retinopathy. <i>Laboratory Investigation</i> , <b>1998</b> , 78, 109-16	5.9	40
78	Hypoxia down-regulates placenta growth factor, whereas fetal growth restriction up-regulates placenta growth factor expression: molecular evidence for "placental hyperoxia" in intrauterine growth restriction. <i>Laboratory Investigation</i> , <b>1999</b> , 79, 151-70	5.9	40
77	Prediction of pre-eclampsia: review of reviews. <i>Ultrasound in Obstetrics and Gynecology</i> , <b>2019</b> , 54, 16-27	5.8	39
76	Carbon monoxide inhibits sprouting angiogenesis and vascular endothelial growth factor receptor-2 phosphorylation. <i>Thrombosis and Haemostasis</i> , <b>2015</b> , 113, 329-37	7	37
75	Role of epithelial Na+ channels in endothelial function. <i>Journal of Cell Science</i> , <b>2016</b> , 129, 290-7	5.3	36
74	The elevation in circulating anti-angiogenic factors is independent of markers of neutrophil activation in preeclampsia. <i>Angiogenesis</i> , <b>2012</b> , 15, 333-40	10.6	36
73	Tie-2 is expressed on thyroid follicular cells, is increased in goiter, and is regulated by thyrotropin through cyclic adenosine 3RSRmonophosphate. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2001</b> , 86, 2709-16	5.6	36

72	Localisation of hepatocyte growth factor and its receptor (c-met) protein and mRNA in human term placenta. <i>Growth Factors</i> , <b>1996</b> , 13, 133-9	1.6	34
71	Vascular endothelial growth factor receptor-2-mediated mitogenesis is negatively regulated by vascular endothelial growth factor receptor-1 in tumor epithelial cells. <i>American Journal of Pathology</i> , <b>2001</b> , 158, 265-73	5.8	34
70	Haemostatometer: a new in vitro technique for assessing haemostatic activity of blood. <i>Thrombosis Research</i> , <b>1984</b> , 34, 341-57	8.2	33
69	Placenta growth factor-1 exerts time-dependent stabilization of adherens junctions following VEGF-induced vascular permeability. <i>PLoS ONE</i> , <b>2011</b> , 6, e18076	3.7	32
68	Activation of proteinase-activated receptor 2 stimulates soluble vascular endothelial growth factor receptor 1 release via epidermal growth factor receptor transactivation in endothelial cells. <i>Hypertension</i> , <b>2010</b> , 55, 689-97	8.5	31
67	Expression of VEGF-C and activation of its receptors VEGFR-2 and VEGFR-3 in trophoblast. Histology and Histopathology, <b>2001</b> , 16, 359-75	1.4	30
66	Activation of protein kinase C is required for oxytocin-induced contractility in human pregnant myometrium. <i>Human Reproduction</i> , <b>1996</b> , 11, 2285-90	5.7	29
65	Inhibition of pyruvate kinase M2 by reactive oxygen species contributes to the development of pulmonary arterial hypertension. <i>Journal of Molecular and Cellular Cardiology</i> , <b>2016</b> , 91, 179-87	5.8	28
64	Compartmentalization and cyclic variation of immunoreactivity of renin and angiotensin converting enzyme in human endometrium throughout the menstrual cycle. <i>Human Reproduction</i> , <b>1997</b> , 12, 2804-9	5.7	28
63	VEGF-E activates endothelial nitric oxide synthase to induce angiogenesis via cGMP and PKG-independent pathways. <i>Biochemical and Biophysical Research Communications</i> , <b>2006</b> , 345, 1275-82	3.4	28
62	Drug Repurposing Screen Identifies Foxo1-Dependent Angiopoietin-2 Regulation in Sepsis. <i>Critical Care Medicine</i> , <b>2015</b> , 43, e230-40	1.4	27
61	Modulation of Trophoblast Cell Death by Oxygen and EGF. <i>Molecular Medicine</i> , <b>2002</b> , 8, 847-856	6.2	26
60	Vascular endothelial growth factor signaling requires glycine to promote angiogenesis. <i>Scientific Reports</i> , <b>2017</b> , 7, 14749	4.9	24
59	Eicosanoid synthesis and Helicobacter pylori associated gastritis: increase in leukotriene C4 generation associated with H. pylori colonization. <i>Prostaglandins</i> , <b>1992</b> , 44, 75-86		23
58	Soluble Fms-Like Tyrosine Kinase-1 Alters Cellular Metabolism and Mitochondrial Bioenergetics in Preeclampsia. <i>Frontiers in Physiology</i> , <b>2018</b> , 9, 83	4.6	22
57	Immunohistochemical localization of parathyroid hormone-related protein (PTHrP) in human term placenta and membranes. <i>Placenta</i> , <b>1994</b> , 15, 653-60	3.4	22
56	Platelet-activating factor stimulates phospholipase C activity in human endometrium. <i>Journal of Cellular Physiology</i> , <b>1992</b> , 152, 207-14	7	22
55	Antiangiogenic effect of soluble vascular endothelial growth factor receptor-1 in placental angiogenesis. <i>Endothelium: Journal of Endothelial Cell Research</i> , <b>2005</b> , 12, 89-95		21

54	The release of nitric oxide from S-nitrosothiols promotes angiogenesis. <i>PLoS ONE</i> , <b>2006</b> , 1, e25	3.7	21
53	Vesicle miR-195 derived from Endothelial Cells Inhibits Expression of Serotonin Transporter in Vessel Smooth Muscle Cells. <i>Scientific Reports</i> , <b>2017</b> , 7, 43546	4.9	20
52	Selective inhibition of the human tie-1 promoter with triplex-forming oligonucleotides targeted to Ets binding sites. <i>Molecular Medicine</i> , <b>2006</b> , 12, 8-16	6.2	20
51	Functional platelet-activating factor receptors linked to inositol lipid hydrolysis, calcium mobilization and tyrosine kinase activity in the human endometrial HEC-1B cell line. <i>Reproduction</i> , <b>1994</b> , 101, 459-66	3.8	20
50	Upregulation of urotensin II receptor in preeclampsia causes in vitro placental release of soluble vascular endothelial growth factor receptor 1 in hypoxia. <i>Hypertension</i> , <b>2010</b> , 56, 172-8	8.5	19
49	Angiopoietin-1 induces migration of monocytes in a tie-2 and integrin-independent manner. <i>Hypertension</i> , <b>2010</b> , 56, 477-83	8.5	18
48	Development and validation of Prediction models for Risks of complications in Early-onset Pre-eclampsia (PREP): a prospective cohort study. <i>Health Technology Assessment</i> , <b>2017</b> , 21, 1-100	4.4	18
47	Survival after cardiac resynchronization therapy: results from 50\(\textit{D}\)84 implantations. <i>Europace</i> , <b>2019</b> , 21, 754-762	3.9	18
46	Molecular mechanisms and therapeutic implications of the carbon monoxide/hmox1 and the hydrogen sulfide/CSE pathways in the prevention of pre-eclampsia and fetal growth restriction. <i>Pregnancy Hypertension</i> , <b>2014</b> , 4, 243-4	2.6	17
45	Antenatal corticosteroids impact the inflammatory rather than the antiangiogenic profile of women with preeclampsia. <i>Hypertension</i> , <b>2014</b> , 63, 1285-92	8.5	17
44	Platelet-activating factor mediates phosphatidylcholine hydrolysis by phospholipase D in human endometrium. <i>Biology of Reproduction</i> , <b>1992</b> , 47, 59-65	3.9	17
43	Activation of Functional Oxytocin Receptors Stimulates Cell Proliferation in Human Trophoblast and Choriocarcinoma Cell Lines		17
42	Hormone Replacement Therapy: Would it be Possible to Replicate a Functional Ovary?. <i>International Journal of Molecular Sciences</i> , <b>2018</b> , 19,	6.3	16
41	Hemin Conditioning Targets the Vascular and Immunologic Compartments and Restrains Prostate Tumor Development. <i>Clinical Cancer Research</i> , <b>2017</b> , 23, 5135-5148	12.9	15
40	Hydrogen sulfide: A novel mechanism for the vascular protection by resveratrol under oxidative stress in mouse aorta. <i>Vascular Pharmacology</i> , <b>2016</b> , 87, 76-82	5.9	15
39	Multiple roles of angiopoietins in atherogenesis. Current Opinion in Lipidology, <b>2011</b> , 22, 380-5	4.4	15
38	Activation of phospholipase A2 and phospholipase C by endothelin-1 in human endometrium. <i>Journal of Endocrinology</i> , <b>1992</b> , 135, 383-90	4.7	15
37	Angiopoietin-1 promotes atherosclerosis by increasing the proportion of circulating Gr1+ monocytes. <i>Cardiovascular Research</i> , <b>2017</b> , 113, 81-89	9.9	14

36	Resveratrol Stimulates Hydrogen Sulfide (H2 S) Formation to Relax Murine Corpus Cavernosum. Journal of Sexual Medicine, <b>2015</b> , 12, 2004-12	1.1	13
35	Carbon monoxide inhibits inward rectifier potassium channels in cardiomyocytes. <i>Nature Communications</i> , <b>2014</b> , 5, 4676	17.4	13
34	Fibroblast growth factor receptor-1 is a critical component for endometrial remodeling: localization and expression of basic fibroblast growth factor and FGF-R1 in human endometrium during the menstrual cycle and decreased FGF-R1 expression in menorrhagia. <i>Laboratory</i>	5.9	13
33	Investigation, 1997, 77, 389-402 Bioenergetic effects of hydrogen sulfide suppress soluble Flt-1 and soluble endoglin in cystathionine gamma-lyase compromised endothelial cells. <i>Scientific Reports</i> , 2020, 10, 15810	4.9	11
32	Heparin-binding angiogenic growth factors in pregnancy. <i>Placenta</i> , <b>1997</b> , 18, 215-258	3.4	10
31	Hydrogen sulfide releasing molecule MZe786 inhibits soluble Flt-1 and prevents preeclampsia in a refined RUPP mouse model. <i>Redox Biology</i> , <b>2021</b> , 38, 101814	11.3	10
30	Effect of ethnicity on live birth rates after in vitro fertilisation or intracytoplasmic sperm injection treatment: possible explanations and further observations. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , <b>2014</b> , 121, 1576-7	3.7	8
29	MZe786 Rescues Cardiac Mitochondrial Activity in High sFlt-1 and Low HO-1 Environment. <i>Antioxidants</i> , <b>2020</b> , 9,	7.1	8
28	MZe786, a hydrogen sulfide-releasing aspirin prevents preeclampsia in heme oxygenase-1 haplodeficient pregnancy under high soluble flt-1 environment. <i>Redox Biology</i> , <b>2021</b> , 38, 101768	11.3	8
27	Heterodimerisation between VEGFR-1 and VEGFR-2 and not the homodimers of VEGFR-1 inhibit VEGFR-2 activity. <i>Vascular Pharmacology</i> , <b>2017</b> , 88, 11-20	5.9	7
26	External validation, update and development of prediction models for pre-eclampsia using an Individual Participant Data (IPD) meta-analysis: the International Prediction of Pregnancy Complication Network (IPPIC pre-eclampsia) protocol. <i>Diagnostic and Prognostic Research</i> , <b>2017</b> , 1, 16	5.5	7
25	Oestradiol-17 beta modulates PAF-evoked phospholipase D activity but not inositide-lipid hydrolysis in human endometrial cell line, HEC-1B. <i>Cellular Signalling</i> , <b>1995</b> , 7, 403-9	4.9	7
24	Validation and development of models using clinical, biochemical and ultrasound markers for predicting pre-eclampsia: an individual participant data meta-analysis. <i>Health Technology Assessment</i> , <b>2020</b> , 24, 1-252	4.4	6
23	Activation of human endometrial phospholipase D by bradykinin. <i>Cellular Signalling</i> , <b>1995</b> , 7, 599-609	4.9	5
22	Eicosanoid synthesis in duodenal ulcer disease: decrease in leukotriene C4 by colloidal bismuth subcitrate. <i>Gut</i> , <b>1992</b> , 33, 159-63	19.2	5
21	External validation of prognostic models predicting pre-eclampsia: individual participant data meta-analysis. <i>BMC Medicine</i> , <b>2020</b> , 18, 302	11.4	4
20	Response to letter regarding article, "dysregulation of hydrogen sulfide (H2S) producing enzyme cystathionine Eyase (CSE) contributes to maternal hypertension and placental abnormalities in preeclampsia". <i>Circulation</i> , <b>2014</b> , 129, e517-8	16.7	4
19	Localization of bradykinin type II receptor mRNA in human endometrium. <i>Molecular Human Reproduction</i> , <b>1996</b> , 2, 35-9	4.4	4

18	The role of platelet-activating factor and its receptor in endometrial receptivity. <i>Advances in Experimental Medicine and Biology</i> , <b>1996</b> , 416, 277-90	3.6	4
17	Hyperglycaemia up-regulates placental growth factor (PlGF) expression and secretion in endothelial cells via suppression of PI3 kinase-Akt signalling and activation of FOXO1. <i>Scientific Reports</i> , <b>2021</b> , 11, 16344	4.9	3
16	Development and validation of prediction models for risk of adverse outcomes in women with early-onset pre-eclampsia: protocol of the prospective cohort PREP study. <i>Diagnostic and Prognostic Research</i> , <b>2017</b> , 1, 6	5.5	2
15	Placenta growth factor (PIGF) expression and function in human placenta. <i>Placenta</i> , <b>1996</b> , 17, A43	3.4	2
14	Localisation of bradykinin-like immunoreactivity and modulation of bradykinin-evoked phospholipase D activity by 17 beta-oestradiol in human endometrium. <i>Growth Factors</i> , <b>1995</b> , 12, 203-9	1.6	2
13	Growth Factor Regulators of Placental Angiogenesis <b>2000</b> , 149-162		2
12	The localisation and expression of hepatocyte growth factor (HGF) and its receptor, c-met, in human placenta from the 1st, 2nd, and 3rd trimesters and in the presence of severe IUGR. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , <b>1998</b> , 105, 1231-1232	3.7	1
11	The role of hepatocyte growth factor and its receptor, C-met, in placental development and fetal growth restriction. <i>Placenta</i> , <b>1998</b> , 19, 159-176	3.4	1
10	Abdominal tuberculosis: the great mimic. British Journal of Hospital Medicine, 2001, 62, 368-9		1
9	The impact of COVID-19 on pregnancy and therapeutic drug development. <i>British Journal of Pharmacology</i> , <b>2021</b> ,	8.6	1
8	Re: Oral and maxillofacial surgery simulation: efficacy for medical students. <i>British Journal of Oral and Maxillofacial Surgery</i> , <b>2020</b> , 58, 1357-1358	1.4	
7	OC22.04: Cardiovascular remodeling in children who suffered growth restriction in utero. <i>Ultrasound in Obstetrics and Gynecology</i> , <b>2010</b> , 36, 41-41	5.8	
6	Nitric Oxide Released VIA Vegfr-1 Suppresses Vegfr-2 Mediated Endothelial Cell Growth and Regulates Angiogenesis. <i>Clinical Science</i> , <b>2001</b> , 100, 19P-19P		
5	Angiopoietin-1 Promotes Trophoblast Migration and Angiopoietin-2 Stimulates Trophoblast Proliferation and Release of Nitric Oxide during Placental Development. <i>Clinical Science</i> , <b>2001</b> , 100, 19P	-20P	
4	The Role of Heme Oxygenase in Pregnancy <b>2002</b> , 67-82		
3	A New Paradigm for Vascular Endothelial Cell Growth Factor Receptor Signaling in Adult Endothelium <b>2003</b> , 153-166		
2	Flt1 acts as a negative regulator of tip cell formation and branching morphogenesis in the zebrafish embryo. <i>Journal of Cell Science</i> , <b>2011</b> , 124, e1-e1	5.3	
1	Angiogenesis in Cervical Neoplasia289-300		