

Asif Ahmed

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6585448/asif-ahmed-publications-by-year.pdf>

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

143 papers	8,157 citations	49 h-index	87 g-index
165 ext. papers	8,965 ext. citations	6.7 avg, IF	5.76 L-index

#	Paper	IF	Citations
143	The impact of COVID-19 on pregnancy and therapeutic drug development. <i>British Journal of Pharmacology</i> , 2021 ,	8.6	1
142	MZe786, a hydrogen sulfide-releasing aspirin prevents preeclampsia in heme oxygenase-1 haplodeficient pregnancy under high soluble Flt-1 environment. <i>Redox Biology</i> , 2021 , 38, 101768	11.3	8
141	Hydrogen sulfide releasing molecule MZe786 inhibits soluble Flt-1 and prevents preeclampsia in a refined RUPP mouse model. <i>Redox Biology</i> , 2021 , 38, 101814	11.3	10
140	Hyperglycaemia up-regulates placental growth factor (PlGF) expression and secretion in endothelial cells via suppression of PI3 kinase-Akt signalling and activation of FOXO1. <i>Scientific Reports</i> , 2021 , 11, 16344	4.9	3
139	External validation of prognostic models predicting pre-eclampsia: individual participant data meta-analysis. <i>BMC Medicine</i> , 2020 , 18, 302	11.4	4
138	Re: Oral and maxillofacial surgery simulation: efficacy for medical students. <i>British Journal of Oral and Maxillofacial Surgery</i> , 2020 , 58, 1357-1358	1.4	
137	Validation and development of models using clinical, biochemical and ultrasound markers for predicting pre-eclampsia: an individual participant data meta-analysis. <i>Health Technology Assessment</i> , 2020 , 24, 1-252	4.4	6
136	Pravastatin for early-onset pre-eclampsia: a randomised, blinded, placebo-controlled trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2020 , 127, 478-488	3.7	44
135	Bioenergetic effects of hydrogen sulfide suppress soluble Flt-1 and soluble endoglin in cystathionine gamma-lyase compromised endothelial cells. <i>Scientific Reports</i> , 2020 , 10, 15810	4.9	11
134	MZe786 Rescues Cardiac Mitochondrial Activity in High sFlt-1 and Low HO-1 Environment. <i>Antioxidants</i> , 2020 , 9,	7.1	8
133	Establishment of porcine and human expanded potential stem cells. <i>Nature Cell Biology</i> , 2019 , 21, 687-699	9.4	127
132	Survival after cardiac resynchronization therapy: results from 50084 implantations. <i>Europace</i> , 2019 , 21, 754-762	3.9	18
131	Prediction of pre-eclampsia: review of reviews. <i>Ultrasound in Obstetrics and Gynecology</i> , 2019 , 54, 16-27	5.8	39
130	Soluble Fms-Like Tyrosine Kinase-1 Alters Cellular Metabolism and Mitochondrial Bioenergetics in Preeclampsia. <i>Frontiers in Physiology</i> , 2018 , 9, 83	4.6	22
129	Hormone Replacement Therapy: Would it be Possible to Replicate a Functional Ovary?. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	16
128	Angiopoietin-1 promotes atherosclerosis by increasing the proportion of circulating Gr1+ monocytes. <i>Cardiovascular Research</i> , 2017 , 113, 81-89	9.9	14
127	Development and validation of prediction models for risk of adverse outcomes in women with early-onset pre-eclampsia: protocol of the prospective cohort PREP study. <i>Diagnostic and Prognostic Research</i> , 2017 , 1, 6	5.5	2

126	Vesicle miR-195 derived from Endothelial Cells Inhibits Expression of Serotonin Transporter in Vessel Smooth Muscle Cells. <i>Scientific Reports</i> , 2017 , 7, 43546	4.9	20
125	Hemin Conditioning Targets the Vascular and Immunologic Compartments and Restrains Prostate Tumor Development. <i>Clinical Cancer Research</i> , 2017 , 23, 5135-5148	12.9	15
124	Heterodimerisation between VEGFR-1 and VEGFR-2 and not the homodimers of VEGFR-1 inhibit VEGFR-2 activity. <i>Vascular Pharmacology</i> , 2017 , 88, 11-20	5.9	7
123	Small Molecule, Big Prospects: MicroRNA in Pregnancy and Its Complications. <i>Journal of Pregnancy</i> , 2017 , 2017, 6972732	2.5	68
122	External validation, update and development of prediction models for pre-eclampsia using an Individual Participant Data (IPD) meta-analysis: the International Prediction of Pregnancy Complication Network (IPPIC pre-eclampsia) protocol. <i>Diagnostic and Prognostic Research</i> , 2017 , 1, 16	5.5	7
121	Evidence-Based Revised View of the Pathophysiology of Preeclampsia. <i>Advances in Experimental Medicine and Biology</i> , 2017 , 956, 355-374	3.6	53
120	Vascular endothelial growth factor signaling requires glycine to promote angiogenesis. <i>Scientific Reports</i> , 2017 , 7, 14749	4.9	24
119	Development and validation of Prediction models for Risks of complications in Early-onset Pre-eclampsia (PREP): a prospective cohort study. <i>Health Technology Assessment</i> , 2017 , 21, 1-100	4.4	18
118	Hydrogen sulfide: A novel mechanism for the vascular protection by resveratrol under oxidative stress in mouse aorta. <i>Vascular Pharmacology</i> , 2016 , 87, 76-82	5.9	15
117	Inhibition of pyruvate kinase M2 by reactive oxygen species contributes to the development of pulmonary arterial hypertension. <i>Journal of Molecular and Cellular Cardiology</i> , 2016 , 91, 179-87	5.8	28
116	Role of epithelial Na ⁺ channels in endothelial function. <i>Journal of Cell Science</i> , 2016 , 129, 290-7	5.3	36
115	The role of H ₂ S bioavailability in endothelial dysfunction. <i>Trends in Pharmacological Sciences</i> , 2015 , 36, 568-78	13.2	106
114	Unravelling the theories of pre-eclampsia: are the protective pathways the new paradigm?. <i>British Journal of Pharmacology</i> , 2015 , 172, 1574-86	8.6	50
113	Drug Repurposing Screen Identifies Foxo1-Dependent Angiopoietin-2 Regulation in Sepsis. <i>Critical Care Medicine</i> , 2015 , 43, e230-40	1.4	27
112	Carbon monoxide inhibits sprouting angiogenesis and vascular endothelial growth factor receptor-2 phosphorylation. <i>Thrombosis and Haemostasis</i> , 2015 , 113, 329-37	7	37
111	Resveratrol Stimulates Hydrogen Sulfide (H ₂ S) Formation to Relax Murine Corpus Cavernosum. <i>Journal of Sexual Medicine</i> , 2015 , 12, 2004-12	1.1	13
110	Molecular mechanisms and therapeutic implications of the carbon monoxide/hmox1 and the hydrogen sulfide/CSE pathways in the prevention of pre-eclampsia and fetal growth restriction. <i>Pregnancy Hypertension</i> , 2014 , 4, 243-4	2.6	17
109	Response to letter regarding article, "dysregulation of hydrogen sulfide (H ₂ S) producing enzyme cystathionine γ -lyase (CSE) contributes to maternal hypertension and placental abnormalities in preeclampsia". <i>Circulation</i> , 2014 , 129, e517-8	16.7	4

108	Carbon monoxide inhibits inward rectifier potassium channels in cardiomyocytes. <i>Nature Communications</i> , 2014 , 5, 4676	17.4	13
107	Effect of ethnicity on live birth rates after in vitro fertilisation or intracytoplasmic sperm injection treatment: possible explanations and further observations. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2014 , 121, 1576-7	3.7	8
106	Antenatal corticosteroids impact the inflammatory rather than the antiangiogenic profile of women with preeclampsia. <i>Hypertension</i> , 2014 , 63, 1285-92	8.5	17
105	Antiproliferative effects of carbon monoxide on pancreatic cancer. <i>Digestive and Liver Disease</i> , 2014 , 46, 369-75	3.3	65
104	Therapeutic potential of statins and the induction of heme oxygenase-1 in preeclampsia. <i>Journal of Reproductive Immunology</i> , 2014 , 101-102, 153-160	4.2	54
103	Carbon monoxide expedites metabolic exhaustion to inhibit tumor growth. <i>Cancer Research</i> , 2013 , 73, 7009-21	10.1	210
102	Dysregulation of hydrogen sulfide producing enzyme cystathionine β -lyase contributes to maternal hypertension and placental abnormalities in preeclampsia. <i>Circulation</i> , 2013 , 127, 2514-22	16.7	195
101	The role of heterodimerization between VEGFR-1 and VEGFR-2 in the regulation of endothelial cell homeostasis. <i>Nature Communications</i> , 2012 , 3, 972	17.4	92
100	Loss of Akt activity increases circulating soluble endoglin release in preeclampsia: identification of inter-dependency between Akt-1 and heme oxygenase-1. <i>European Heart Journal</i> , 2012 , 33, 1150-8	9.5	46
99	The elevation in circulating anti-angiogenic factors is independent of markers of neutrophil activation in preeclampsia. <i>Angiogenesis</i> , 2012 , 15, 333-40	10.6	36
98	Resveratrol inhibits the release of soluble fms-like tyrosine kinase (sFlt-1) from human placenta. <i>American Journal of Obstetrics and Gynecology</i> , 2012 , 206, 253.e10-5	6.4	46
97	New insights into the etiology of preeclampsia: identification of key elusive factors for the vascular complications. <i>Thrombosis Research</i> , 2011 , 127 Suppl 3, S72-5	8.2	100
96	Multiple roles of angiopoietins in atherogenesis. <i>Current Opinion in Lipidology</i> , 2011 , 22, 380-5	4.4	15
95	Is inflammation the cause of pre-eclampsia?. <i>Biochemical Society Transactions</i> , 2011 , 39, 1619-27	5.1	81
94	Placenta growth factor-1 exerts time-dependent stabilization of adherens junctions following VEGF-induced vascular permeability. <i>PLoS ONE</i> , 2011 , 6, e18076	3.7	32
93	Autocrine activity of soluble Flt-1 controls endothelial cell function and angiogenesis. <i>Vascular Cell</i> , 2011 , 3, 15	1	47
92	Flt1 acts as a negative regulator of tip cell formation and branching morphogenesis in the zebrafish embryo. <i>Development (Cambridge)</i> , 2011 , 138, 2111-20	6.6	110
91	Heparin elevates circulating soluble fms-like tyrosine kinase-1 immunoreactivity in pregnant women receiving anticoagulation therapy. <i>Circulation</i> , 2011 , 124, 2543-53	16.7	40

90	Flt1 acts as a negative regulator of tip cell formation and branching morphogenesis in the zebrafish embryo. <i>Journal of Cell Science</i> , 2011 , 124, e1-e1	5.3	
89	Angiopoietin-1 induces migration of monocytes in a tie-2 and integrin-independent manner. <i>Hypertension</i> , 2010 , 56, 477-83	8.5	18
88	Fetal growth restriction results in remodeled and less efficient hearts in children. <i>Circulation</i> , 2010 , 121, 2427-36	16.7	289
87	Activation of proteinase-activated receptor 2 stimulates soluble vascular endothelial growth factor receptor 1 release via epidermal growth factor receptor transactivation in endothelial cells. <i>Hypertension</i> , 2010 , 55, 689-97	8.5	31
86	Upregulation of urotensin II receptor in preeclampsia causes in vitro placental release of soluble vascular endothelial growth factor receptor 1 in hypoxia. <i>Hypertension</i> , 2010 , 56, 172-8	8.5	19
85	Hepatitis C virus infection reduces hepatocellular polarity in a vascular endothelial growth factor-dependent manner. <i>Gastroenterology</i> , 2010 , 138, 1134-42	13.3	62
84	Nitric oxide-dependent bone marrow progenitor mobilization by carbon monoxide enhances endothelial repair after vascular injury. <i>Circulation</i> , 2010 , 121, 537-48	16.7	91
83	Reduction of circulating soluble Flt-1 alleviates preeclampsia-like symptoms in a mouse model. <i>Journal of Cellular and Molecular Medicine</i> , 2010 , 14, 1857-67	5.6	134
82	OC22.04: Cardiovascular remodeling in children who suffered growth restriction in utero. <i>Ultrasound in Obstetrics and Gynecology</i> , 2010 , 36, 41-41	5.8	
81	Angiopoietin-2 confers Atheroprotection in apoE ^{-/-} mice by inhibiting LDL oxidation via nitric oxide. <i>Circulation Research</i> , 2009 , 104, 1333-6	15.7	40
80	Can the biology of VEGF and haem oxygenases help solve pre-eclampsia?. <i>Biochemical Society Transactions</i> , 2009 , 37, 1237-42	5.1	47
79	Hypoxia induces dilated cardiomyopathy in the chick embryo: mechanism, intervention, and long-term consequences. <i>PLoS ONE</i> , 2009 , 4, e5155	3.7	95
78	Autoantibody from women with preeclampsia induces soluble Fms-like tyrosine kinase-1 production via angiotensin type 1 receptor and calcineurin/nuclear factor of activated T-cells signaling. <i>Hypertension</i> , 2008 , 51, 1010-9	8.5	142
77	Cardiac dysfunction and cell damage across clinical stages of severity in growth-restricted fetuses. <i>American Journal of Obstetrics and Gynecology</i> , 2008 , 199, 254.e1-8	6.4	191
76	Angiotensin II induces soluble fms-Like tyrosine kinase-1 release via calcineurin signaling pathway in pregnancy. <i>Circulation Research</i> , 2007 , 100, 88-95	15.7	124
75	Negative regulation of soluble Flt-1 and soluble endoglin release by heme oxygenase-1. <i>Circulation</i> , 2007 , 115, 1789-97	16.7	338
74	VEGF induces Tie2 shedding via a phosphoinositide 3-kinase/Akt dependent pathway to modulate Tie2 signaling. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 2619-26	9.4	127
73	Vascular endothelial growth factor promotes physical wound repair and is anti-apoptotic in primary distal lung epithelial and A549 cells. <i>Critical Care Medicine</i> , 2007 , 35, 2164-70	1.4	50

72	Direct evidence for endothelial vascular endothelial growth factor receptor-1 function in nitric oxide-mediated angiogenesis. <i>Circulation Research</i> , 2006 , 99, 715-22	15.7	119
71	VEGF-E activates endothelial nitric oxide synthase to induce angiogenesis via cGMP and PKG-independent pathways. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 345, 1275-82	3.4	28
70	Vascular endothelial growth factor-induced endothelial cell proliferation is regulated by interaction between VEGFR-2, SH-PTP1 and eNOS. <i>Microvascular Research</i> , 2006 , 71, 20-31	3.7	76
69	Selective inhibition of the human tie-1 promoter with triplex-forming oligonucleotides targeted to Ets binding sites. <i>Molecular Medicine</i> , 2006 , 12, 8-16	6.2	20
68	The release of nitric oxide from S-nitrosothiols promotes angiogenesis. <i>PLoS ONE</i> , 2006 , 1, e25	3.7	21
67	Antiangiogenic effect of soluble vascular endothelial growth factor receptor-1 in placental angiogenesis. <i>Endothelium: Journal of Endothelial Cell Research</i> , 2005 , 12, 89-95		21
66	Physical and biomechanical characteristics of rat cervical ripening are not consistent with increased collagenase activity. <i>American Journal of Obstetrics and Gynecology</i> , 2004 , 191, 1695-704	6.4	46
65	Elevated placental soluble vascular endothelial growth factor receptor-1 inhibits angiogenesis in preeclampsia. <i>Circulation Research</i> , 2004 , 95, 884-91	15.7	400
64	Bifunctional role for VEGF-induced heme oxygenase-1 in vivo: induction of angiogenesis and inhibition of leukocytic infiltration. <i>Blood</i> , 2004 , 103, 761-6	2.2	166
63	Activation of vascular endothelial growth factor receptor-1 sustains angiogenesis and Bcl-2 expression via the phosphatidylinositol 3-kinase pathway in endothelial cells. <i>Diabetes</i> , 2003 , 52, 2959-68	6.9	108
62	A New Paradigm for Vascular Endothelial Cell Growth Factor Receptor Signaling in Adult Endothelium 2003 , 153-166		
61	Modulation of Trophoblast Cell Death by Oxygen and EGF. <i>Molecular Medicine</i> , 2002 , 8, 847-856	6.2	26
60	Down-regulation of angiopoietin-1 expression in menorrhagia. <i>American Journal of Pathology</i> , 2002 , 160, 773-80	5.8	53
59	The Role of Heme Oxygenase in Pregnancy 2002 , 67-82		
58	Nitric Oxide Released VIA Vegfr-1 Suppresses Vegfr-2 Mediated Endothelial Cell Growth and Regulates Angiogenesis. <i>Clinical Science</i> , 2001 , 100, 19P-19P		
57	Angiopoietin-1 Promotes Trophoblast Migration and Angiopoietin-2 Stimulates Trophoblast Proliferation and Release of Nitric Oxide during Placental Development. <i>Clinical Science</i> , 2001 , 100, 19P-20P		
56	Abdominal tuberculosis: the great mimic. <i>British Journal of Hospital Medicine</i> , 2001 , 62, 368-9		1
55	Tie-2 is expressed on thyroid follicular cells, is increased in goiter, and is regulated by thyrotropin through cyclic adenosine 3',5'-monophosphate. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 2709-16	5.6	36

54	Vascular endothelial growth factor receptor-1 modulates vascular endothelial growth factor-mediated angiogenesis via nitric oxide. <i>American Journal of Pathology</i> , 2001 , 159, 993-1008	5.8	242
53	Vascular endothelial growth factor receptor-2-mediated mitogenesis is negatively regulated by vascular endothelial growth factor receptor-1 in tumor epithelial cells. <i>American Journal of Pathology</i> , 2001 , 158, 265-73	5.8	34
52	Activation of functional oxytocin receptors stimulates cell proliferation in human trophoblast and choriocarcinoma cell lines. <i>Endocrinology</i> , 2001 , 142, 1130-6	4.8	48
51	Expression of VEGF-C and activation of its receptors VEGFR-2 and VEGFR-3 in trophoblast. <i>Histology and Histopathology</i> , 2001 , 16, 359-75	1.4	30
50	Release and complex formation of soluble VEGFR-1 from endothelial cells and biological fluids. <i>Laboratory Investigation</i> , 2000 , 80, 443-54	5.9	182
49	Angiogenesis and intrauterine growth restriction. <i>Best Practice and Research in Clinical Obstetrics and Gynaecology</i> , 2000 , 14, 981-98	4.6	84
48	Regulation of placental vascular endothelial growth factor (VEGF) and placenta growth factor (PIGF) and soluble Flt-1 by oxygen--a review. <i>Placenta</i> , 2000 , 21 Suppl A, S16-24	3.4	246
47	Induction of Placental Heme Oxygenase-1 Is Protective Against TNF α -Induced Cytotoxicity and Promotes Vessel Relaxation. <i>Molecular Medicine</i> , 2000 , 6, 391-409	6.2	130
46	Angiopoietin-1 and angiopoietin-2 activate trophoblast Tie-2 to promote growth and migration during placental development. <i>American Journal of Pathology</i> , 2000 , 156, 2185-99	5.8	150
45	Induction of placental heme oxygenase-1 is protective against TNF α -induced cytotoxicity and promotes vessel relaxation. <i>Molecular Medicine</i> , 2000 , 6, 391-409	6.2	52
44	Growth Factor Regulators of Placental Angiogenesis 2000 , 149-162		2
43	Immunohistochemical localization of vascular endothelial growth factor (VEGF) and its two specific receptors, Flt-1 and KDR, in the porcine placenta and non-pregnant uterus. <i>Placenta</i> , 1999 , 20, 35-43	3.4	69
42	A molecular target for viral killer toxin: TOK1 potassium channels. <i>Cell</i> , 1999 , 99, 283-91	56.2	89
41	Hypoxia down-regulates placenta growth factor, whereas fetal growth restriction up-regulates placenta growth factor expression: molecular evidence for "placental hyperoxia" in intrauterine growth restriction. <i>Laboratory Investigation</i> , 1999 , 79, 151-70	5.9	40
40	The localisation and expression of hepatocyte growth factor (HGF) and its receptor, c-met, in human placenta from the 1st, 2nd, and 3rd trimesters and in the presence of severe IUGR. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 1998 , 105, 1231-1232	3.7	1
39	The role of hepatocyte growth factor and its receptor, C-met, in placental development and fetal growth restriction. <i>Placenta</i> , 1998 , 19, 159-176	3.4	1
38	Ontogeny of hepatocyte growth factor (HGF) and its receptor (c-met) in human placenta: reduced HGF expression in intrauterine growth restriction. <i>American Journal of Pathology</i> , 1998 , 153, 1139-47	5.8	54
37	Localization, quantification, and activation of platelet-activating factor receptor in human endometrium during the menstrual cycle: PAF stimulates NO, VEGF, and FAKp125. <i>FASEB Journal</i> , 1998 , 12, 831-43	0.9	40

36	Cellular localization of AT1 receptor mRNA and protein in normal placenta and its reduced expression in intrauterine growth restriction. Angiotensin II stimulates the release of vasorelaxants. <i>Journal of Clinical Investigation</i> , 1998 , 101, 442-54	15.9	91
35	Hemeoxygenase-1 inhibits human myometrial contractility via carbon monoxide and is upregulated by progesterone during pregnancy. <i>Journal of Clinical Investigation</i> , 1998 , 101, 949-55	15.9	94
34	Increased expression of placenta growth factor in proliferative diabetic retinopathy. <i>Laboratory Investigation</i> , 1998 , 78, 109-16	5.9	40
33	Compartmentalization and cyclic variation of immunoreactivity of renin and angiotensin converting enzyme in human endometrium throughout the menstrual cycle. <i>Human Reproduction</i> , 1997 , 12, 2804-9	5.7	28
32	Hypoxia or hyperoxia in placental insufficiency?. <i>Lancet, The</i> , 1997 , 350, 826-7	40	48
31	Heparin-binding angiogenic growth factors in pregnancy. <i>Placenta</i> , 1997 , 18, 215-258	3.4	10
30	Role of VEGF receptor-1 (Flt-1) in mediating calcium-dependent nitric oxide release and limiting DNA synthesis in human trophoblast cells. <i>Laboratory Investigation</i> , 1997 , 76, 779-91	5.9	40
29	Fibroblast growth factor receptor-1 is a critical component for endometrial remodeling: localization and expression of basic fibroblast growth factor and FGF-R1 in human endometrium during the menstrual cycle and decreased FGF-R1 expression in menorrhagia. <i>Laboratory Investigation</i> , 1997 , 77, 389-402	5.9	13
28	Localisation of hepatocyte growth factor and its receptor (c-met) protein and mRNA in human term placenta. <i>Growth Factors</i> , 1996 , 13, 133-9	1.6	34
27	Dual role of angiotensin II in the human endometrium. <i>Human Reproduction</i> , 1996 , 11 Suppl 2, 95-108	5.7	90
26	Localisation of placenta growth factor (PIGF) in human term placenta. <i>Growth Factors</i> , 1996 , 13, 243-50,color plates I-II,pre.bk cov	1.6	158
25	Activation of protein kinase C is required for oxytocin-induced contractility in human pregnant myometrium. <i>Human Reproduction</i> , 1996 , 11, 2285-90	5.7	29
24	Placenta growth factor (PIGF) expression and function in human placenta. <i>Placenta</i> , 1996 , 17, A43	3.4	2
23	Localization of bradykinin type II receptor mRNA in human endometrium. <i>Molecular Human Reproduction</i> , 1996 , 2, 35-9	4.4	4
22	The role of platelet-activating factor and its receptor in endometrial receptivity. <i>Advances in Experimental Medicine and Biology</i> , 1996 , 416, 277-90	3.6	4
21	Oestradiol-17 beta modulates PAF-evoked phospholipase D activity but not inositide-lipid hydrolysis in human endometrial cell line, HEC-1B. <i>Cellular Signalling</i> , 1995 , 7, 403-9	4.9	7
20	Activation of human endometrial phospholipase D by bradykinin. <i>Cellular Signalling</i> , 1995 , 7, 599-609	4.9	5
19	Colocalisation of vascular endothelial growth factor and its Flt-1 receptor in human placenta. <i>Growth Factors</i> , 1995 , 12, 235-43	1.6	213

18	Localisation of bradykinin-like immunoreactivity and modulation of bradykinin-evoked phospholipase D activity by 17 beta-oestradiol in human endometrium. <i>Growth Factors</i> , 1995 , 12, 203-9	1.6	2
17	Localization of the angiotensin II and its receptor subtype expression in human endometrium and identification of a novel high-affinity angiotensin II binding site. <i>Journal of Clinical Investigation</i> , 1995 , 96, 848-57	15.9	51
16	Basic FGF activates phospholipase D in endothelial cells in the absence of inositol-lipid hydrolysis. <i>American Journal of Physiology - Cell Physiology</i> , 1994 , 266, C206-12	5.4	42
15	Localization of mRNA for basic fibroblast growth factor in human placenta. <i>Growth Factors</i> , 1994 , 11, 105-11	1.6	47
14	Colocalization of acidic and basic fibroblast growth factor (FGF) in human placenta and the cellular effects of bFGF in trophoblast cell line JEG-3. <i>Growth Factors</i> , 1994 , 10, 259-68	1.6	60
13	Parathyroid hormone-related protein in human term placenta and membranes. <i>Journal of Endocrinology</i> , 1994 , 142, 217-24	4.7	48
12	Vascular endothelial growth factor receptor localization and activation in human trophoblast and choriocarcinoma cells. <i>Biology of Reproduction</i> , 1994 , 51, 524-30	3.9	214
11	Functional platelet-activating factor receptors linked to inositol lipid hydrolysis, calcium mobilization and tyrosine kinase activity in the human endometrial HEC-1B cell line. <i>Reproduction</i> , 1994 , 101, 459-66	3.8	20
10	Immunohistochemical localization of parathyroid hormone-related protein (PTHrP) in human term placenta and membranes. <i>Placenta</i> , 1994 , 15, 653-60	3.4	22
9	Immunolocalisation of vascular endothelial growth factor in human endometrium. <i>Growth Factors</i> , 1994 , 11, 277-82	1.6	103
8	Activation of phospholipase A2 and phospholipase C by endothelin-1 in human endometrium. <i>Journal of Endocrinology</i> , 1992 , 135, 383-90	4.7	15
7	Eicosanoid synthesis in duodenal ulcer disease: decrease in leukotriene C4 by colloidal bismuth subcitrate. <i>Gut</i> , 1992 , 33, 159-63	19.2	5
6	Platelet-activating factor mediates phosphatidylcholine hydrolysis by phospholipase D in human endometrium. <i>Biology of Reproduction</i> , 1992 , 47, 59-65	3.9	17
5	Eicosanoid synthesis and Helicobacter pylori associated gastritis: increase in leukotriene C4 generation associated with H. pylori colonization. <i>Prostaglandins</i> , 1992 , 44, 75-86		23
4	Platelet-activating factor stimulates phospholipase C activity in human endometrium. <i>Journal of Cellular Physiology</i> , 1992 , 152, 207-14	7	22
3	Haemostatometer: a new in vitro technique for assessing haemostatic activity of blood. <i>Thrombosis Research</i> , 1984 , 34, 341-57	8.2	33
2	Activation of Functional Oxytocin Receptors Stimulates Cell Proliferation in Human Trophoblast and Choriocarcinoma Cell Lines		17
1	Angiogenesis in Cervical Neoplasia	289-300	

