Reeta Davis

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6585580/publications.pdf

Version: 2024-02-01

752698
7 20
dex g-index
0 1512
0 1312
ranked citing authors

#	Article	IF	CITATIONS
1	Melamine Detection in Infant Formula Powder Using Near- and Mid-Infrared Spectroscopy. Journal of Agricultural and Food Chemistry, 2009, 57, 3974-3980.	5.2	256
2	Conversion of grass biomass into fermentable sugars and its utilization for medium chain length polyhydroxyalkanoate (mcl-PHA) production by Pseudomonas strains. Bioresource Technology, 2013, 150, 202-209.	9.6	129
3	Medium chain length polyhydroxyalkanoate (mcl-PHA) production from volatile fatty acids derived from the anaerobic digestion of grass. Applied Microbiology and Biotechnology, 2014, 98, 611-620.	3.6	68
4	Production and characterization of bacterial polyhydroxyalkanoate copolymers and evaluation of their blends by fourier transform infrared spectroscopy and scanning electron microscopy. Indian Journal of Microbiology, 2009, 49, 251-258.	2.7	58
5	Use of a mannitol rich ensiled grass press juice (EGPJ) as a sole carbon source for polyhydroxyalkanoates (PHAs) production through high cell density cultivation. Bioresource Technology, 2015, 191, 45-52.	9.6	57
6	Detection ofâ€, <i>E. coli ⟨i⟩â€,O157:H7 from Ground Beef Using Fourier Transform Infrared (FTâ€IR) Spectroscopy and Chemometrics. Journal of Food Science, 2010, 75, M340-6.</i>	3.1	54
7	High cell density cultivation of <i>Pseudomonas putida</i> KT2440 using glucose without the need for oxygen enriched air supply. Biotechnology and Bioengineering, 2015, 112, 725-733.	3.3	53
8	Recent developments in biocatalysis beyond the laboratory. Biotechnology Letters, 2015, 37, 943-954.	2.2	48
9	Chemical, physical and biotechnological approaches to the production of the potent antioxidant hydroxytyrosol. Applied Microbiology and Biotechnology, 2019, 103, 5957-5974.	3.6	46
10	Subtyping of Listeria monocytogenes at the haplotype level by Fourier transform infrared (FT-IR) spectroscopy and multivariate statistical analysis. International Journal of Food Microbiology, 2011, 150, 140-149.	4.7	37
11	Surfactant-mediated hydrothermal pretreatment of Ryegrass followed by enzymatic saccharification for polyhydroxyalkanoate production. Industrial Crops and Products, 2018, 111, 625-632.	5.2	29
12	Detection and differentiation of live and heat-treated Salmonella enterica serovars inoculated onto chicken breast using Fourier transform infrared (FT-IR) spectroscopy. Journal of Applied Microbiology, 2010, 109, 2019-2031.	3.1	27
13	Biosynthesis of polyhydroxyalkanoates co-polymer in E. coli using genes from Pseudomonas and Bacillus. Antonie Van Leeuwenhoek, 2008, 94, 207-216.	1.7	25
14	Differentiation of live, dead and treated cells of Escherichia coli O157:H7 using FT-IR spectroscopy. Journal of Applied Microbiology, 2012, 112, 743-751.	3.1	21
15	Fed-batch strategies using butyrate for high cell density cultivation of Pseudomonas putida and its use as a biocatalyst. Applied Microbiology and Biotechnology, 2014, 98, 9217-9228.	3.6	21
16	Evaluation of Fourier transform infrared (FT-IR) spectroscopy and chemometrics as a rapid approach for sub-typing Escherichia coli O157:H7 isolates. Food Microbiology, 2012, 31, 181-190.	4.2	20
17	Potent anti-melanogenic activity and favorable toxicity profile of selected 4-phenyl hydroxycoumarins in the zebrafish model and the computational molecular modeling studies. Bioorganic and Medicinal Chemistry, 2017, 25, 6286-6296.	3.0	19
18	Role of (R)-specific enoyl coenzyme A hydratases of Pseudomonas sp in the production of polyhydroxyalkanoates. Antonie Van Leeuwenhoek, 2008, 93, 285-296.	1.7	16

#	Article	lF	CITATIONS
19	Biocatalytic versatility of engineered and wild-type tyrosinase from R. solanacearum for the synthesis of 4-halocatechols. Applied Microbiology and Biotechnology, 2018, 102, 5121-5131.	3.6	9