

Katarzyna Taranta-Janusz

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6588827/katarzyna-taranta-janusz-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

28

papers

414

citations

10

h-index

20

g-index

35

ext. papers

571

ext. citations

4.5

avg, IF

3.65

L-index

#	Paper	IF	Citations
28	KIM-1 and NGAL: new markers of obstructive nephropathy. <i>Pediatric Nephrology</i> , 2011 , 26, 579-86	3.2	99
27	Salivary Biomarkers of Oxidative Stress in Children with Chronic Kidney Disease. <i>Journal of Clinical Medicine</i> , 2018 , 7,	5.1	51
26	A Case-Control Study of Salivary Redox Homeostasis in Hypertensive Children. Can Salivary Uric Acid be a Marker of Hypertension?. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	30
25	Urinary cytokine profiles in unilateral congenital hydronephrosis. <i>Pediatric Nephrology</i> , 2012 , 27, 2107-2113	3.13	30
24	Risk Factors for Early Dialysis Dependency in Autosomal Recessive Polycystic Kidney Disease. <i>Journal of Pediatrics</i> , 2018 , 199, 22-28.e6	3.6	28
23	Salivary FRAP as A Marker of Chronic Kidney Disease Progression in Children. <i>Antioxidants</i> , 2019 , 8,	7.1	27
22	Dysfunction of Salivary Glands, Disturbances in Salivary Antioxidants and Increased Oxidative Damage in Saliva of Overweight and Obese Adolescents. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	23
21	Urinary angiotensinogen as a novel marker of obstructive nephropathy in children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2013 , 102, e429-33	3.1	20
20	Salivary Gland Dysfunction, Protein Glycooxidation and Nitrosative Stress in Children with Chronic Kidney Disease. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	18
19	Rare Variants in BNC2 Are Implicated in Autosomal-Dominant Congenital Lower Urinary-Tract Obstruction. <i>American Journal of Human Genetics</i> , 2019 , 104, 994-1006	11	11
18	New tubular injury markers in children with a solitary functioning kidney. <i>Pediatric Nephrology</i> , 2014 , 29, 1599-605	3.2	10
17	Refining genotype-phenotype correlations in 304 patients with autosomal recessive polycystic kidney disease and PKHD1 gene variants. <i>Kidney International</i> , 2021 , 100, 650-659	9.9	9
16	Determining normal values of urinary phosphorus excretion in 3913 healthy children aged 2-18 to aid early diagnosis and treatment for urolithiasis. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2017 , 106, 1170-1175	3.1	8
15	Severe neurological outcomes after very early bilateral nephrectomies in patients with autosomal recessive polycystic kidney disease (ARPKD). <i>Scientific Reports</i> , 2020 , 10, 16025	4.9	8
14	Osteopontin and symmetric dimethylarginine plasma levels in solitary functioning kidney in children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2012 , 101, e369-72	3.1	7
13	Serum Renalase Levels in Adolescents with Primary Hypertension. <i>Pediatric Cardiology</i> , 2018 , 39, 1258-1264	12.64	6
12	Urine exoglycosidases are potential markers of renal tubular injury in children with ureteropelvic junction obstruction. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2015 , 104, e518-23	3.1	6

11	Mild X-linked Alport syndrome due to the COL4A5 G624D variant originating in the Middle Ages is predominant in Central/East Europe and causes kidney failure in midlife. <i>Kidney International</i> , 2021 , 99, 1451-1458	9.9	5
10	Pediatric reference data on activity of urinary N-acetyl-D-hexosaminidase and its isoenzymes. <i>Advances in Medical Sciences</i> , 2018 , 63, 94-99	2.8	4
9	Early childhood height-adjusted total kidney volume as a risk marker of kidney survival in ARPKD. <i>Scientific Reports</i> , 2021 , 11, 21677	4.9	3
8	Urine NGAL and KIM-1: tubular injury markers in acute lymphoblastic leukemia survivors. <i>Cancer Chemotherapy and Pharmacology</i> , 2020 , 86, 741-749	3.5	2
7	Urinary exoglycosidases, reference values in healthy children. <i>Advances in Medical Sciences</i> , 2018 , 63, 224-229	2.8	2
6	Low-Dose Antibiotic Prophylaxis Induces Rapid Modifications of the Gut Microbiota in Infants With Vesicoureteral Reflux. <i>Frontiers in Pediatrics</i> , 2021 , 9, 674716	3.4	2
5	Is urine intercellular adhesion molecule-1 a marker of renal disorder in children with ureteropelvic junction obstruction?. <i>Biomarkers</i> , 2016 , 21, 123-8	2.6	2
4	Untargeted Metabolomics Analysis of the Serum Metabolic Signature of Childhood Obesity.. <i>Nutrients</i> , 2022 , 14,	6.7	1
3	Tumor Necrosis Factor-Like Weak Inducer of Apoptosis and Selected Cytokines-Potential Biomarkers in Children with Solitary Functioning Kidney. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	1
2	Urinary procollagen III aminoterminal propeptide and Ecatenin - New diagnostic biomarkers in solitary functioning kidney?. <i>Advances in Medical Sciences</i> , 2019 , 64, 189-194	2.8	0
1	Activity of lysosomal exoglycosidases in the urine of healthy normotensive and pre-hypertensive children. <i>Advances in Medical Sciences</i> , 2019 , 64, 24-31	2.8	