

Yoshitaka Hashimoto

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6603755/yoshitaka-hashimoto-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

147
papers

4,157
citations

26
h-index

62
g-index

178
ext. papers

5,391
ext. citations

4
avg. IF

5.21
L-index

| # | Paper | IF | Citations |
|-----|--|-----|-----------|
| 147 | Relationship between eosinophils counts and muscle mass decline in older people with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort.. <i>Experimental Gerontology</i> , 2022 , 159, 1116-1121 | 4.5 | 1 |
| 146 | Characterization of Peripheral Blood TCR in Patients with Type 1 Diabetes Mellitus by BD Rhapsody™ VDJ CDR3 Assay. <i>Cells</i> , 2022 , 11, 1623 | 7.9 | 0 |
| 145 | Asymptomatic postprandial hypotension in patients with diabetes: The KAMOGAWA-HBP study. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 837-844 | 3.9 | 0 |
| 144 | Sarcopenia Is Associated With a Risk of Mortality in People With Type 2 Diabetes Mellitus. <i>Frontiers in Endocrinology</i> , 2021 , 12, 783363 | 5.7 | 0 |
| 143 | Association between Geriatric Nutrition Risk Index and The Presence of Sarcopenia in People with Type 2 Diabetes Mellitus: A Cross-Sectional Study. <i>Nutrients</i> , 2021 , 13, | 6.7 | 1 |
| 142 | Impact of extracellular-to-intracellular fluid volume ratio on albuminuria in patients with type 2 diabetes: A cross-sectional and longitudinal cohort study. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 1202-1211 | 3.9 | 1 |
| 141 | Association of mean corpuscular volume with sarcopenia and visceral obesity in individuals without anemia. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 1287-1292 | 3.9 | 1 |
| 140 | Group 3 Innate Lymphoid Cells Protect Steatohepatitis From High-Fat Diet Induced Toxicity. <i>Frontiers in Immunology</i> , 2021 , 12, 648754 | 8.4 | 9 |
| 139 | Trigger finger is associated with risk of incident cardiovascular disease in individuals with type 2 diabetes: a retrospective cohort study. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9, | 4.5 | 1 |
| 138 | Changes in the Size of a Ruptured Pheochromocytoma after Transcatheter Arterial Embolization. <i>Case Reports in Medicine</i> , 2021 , 2021, 5568978 | 0.7 | 0 |
| 137 | Short energy intake is associated with muscle mass loss in older patients with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort. <i>Clinical Nutrition</i> , 2021 , 40, 1613-1620 | 5.9 | 7 |
| 136 | Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with Glycemic Variability in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2021 , 13, | 6.7 | 1 |
| 135 | Trans Fatty Acid Intake Induces Intestinal Inflammation and Impaired Glucose Tolerance. <i>Frontiers in Immunology</i> , 2021 , 12, 669672 | 8.4 | 5 |
| 134 | The effect of COVID-19 pandemic on the lifestyle and glycemic control in patients with type 1 diabetes: a retrospective cohort study. <i>Diabetology International</i> , 2021 , 1-6 | 2.3 | 7 |
| 133 | U-shaped association between the triglyceride-glucose index and the risk of incident diabetes in people with normal glycemic level: A population-base longitudinal cohort study. <i>Clinical Nutrition</i> , 2021 , 40, 1555-1561 | 5.9 | 8 |
| 132 | Clinical characteristics and longitudinal changes of patients with non-alcoholic fatty liver disease in 2 decades: the NAGALA study. <i>BMC Gastroenterology</i> , 2021 , 21, 223 | 3 | 0 |
| 131 | Erythritol Ameliorates Small Intestinal Inflammation Induced by High-Fat Diets and Improves Glucose Tolerance. <i>International Journal of Molecular Sciences</i> , 2021 , 22, | 6.3 | 5 |

| | | | |
|-----|---|-----|----|
| 130 | 1182-P: Trans Fatty Acid Intake Induces Impaired Glucose Tolerance via Dysbiosis. <i>Diabetes</i> , 2021 , 70, 1182-P | 0.9 | |
| 129 | Habitual Dietary Intake Affects the Altered Pattern of Gut Microbiome by Acarbose in Patients with Type 2 Diabetes. <i>Nutrients</i> , 2021 , 13, | 6.7 | 3 |
| 128 | 801-P: SGLT2 Inhibitor, Luseogliflozin, Prevent Sarcopenia by Improving Extracellular Lipidome. <i>Diabetes</i> , 2021 , 70, 801-P | 0.9 | |
| 127 | ILC2s Improve Glucose Metabolism Through the Control of Saturated Fatty Acid Absorption Within Visceral Fat. <i>Frontiers in Immunology</i> , 2021 , 12, 669629 | 8.4 | 4 |
| 126 | Microbe-associated metabolites as targets for incident type2 diabetes. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 476-478 | 3.9 | 1 |
| 125 | Distinct associations of intraperitoneal and retroperitoneal visceral adipose tissues with metabolic syndrome and its components. <i>Clinical Nutrition</i> , 2021 , 40, 3479-3484 | 5.9 | 1 |
| 124 | Effect of coronavirus disease 2019 pandemic on the lifestyle and glycemic control in patients with type 2 diabetes: a cross-section and retrospective cohort study. <i>Endocrine Journal</i> , 2021 , 68, 201-210 | 2.9 | 18 |
| 123 | Association between the frequency of toothbrushing and lifestyle in people with type 2 diabetes mellitus: at the baseline date of the Kamogawa-DM cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2021 , 69, 294-298 | 3.1 | |
| 122 | Status of online diet management program users in Japan during the 2020 Coronavirus disease 2019 pandemic. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2021 , 69, 305-310 | 3.1 | 0 |
| 121 | A survey on consciousness towards the proper use of metformin and medical cost in Japanese patients with type 2 diabetes. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2021 , 69, 286-293 | 3.1 | 0 |
| 120 | Sarcopenic obesity is associated with macroalbuminuria in patients with type 2 diabetes: a cross-sectional study. <i>Endocrine Journal</i> , 2021 , 68, 781-789 | 2.9 | 2 |
| 119 | Low circulating arachidonic acid is associated with macroalbuminuria in diabetic patients: a cross-sectional examination of the KAMOGAWA-DM cohort study. <i>BMC Nephrology</i> , 2021 , 22, 68 | 2.7 | 0 |
| 118 | Vitamin Intake and Loss of Muscle Mass in Older People with Type 2 Diabetes: A Prospective Study of the KAMOGAWA-DM Cohort. <i>Nutrients</i> , 2021 , 13, | 6.7 | 2 |
| 117 | Obesity and metabolic abnormalities as risks of alcoholic fatty liver in men: NAGALA study. <i>BMC Gastroenterology</i> , 2021 , 21, 321 | 3 | |
| 116 | Metabolic associated fatty liver disease is a risk factor for chronic kidney disease. <i>Journal of Diabetes Investigation</i> , 2021 , | 3.9 | 4 |
| 115 | Low circulating dihomo-gamma-linolenic acid is associated with diabetic retinopathy: a cross sectional study of KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2021 , 68, 421-428 | 2.9 | 6 |
| 114 | Let-7e-5p Regulates IGF2BP2, and Induces Muscle Atrophy.. <i>Frontiers in Endocrinology</i> , 2021 , 12, 791363 | 5.7 | |
| 113 | Intraperitoneal, but not retroperitoneal, visceral adipose tissue is associated with diabetes mellitus: a cross-sectional, retrospective pilot analysis. <i>Diabetology and Metabolic Syndrome</i> , 2020 , 12, 103 | 5.6 | 2 |

| | | | |
|-----|---|-----|----|
| 112 | Usefulness of Exercise for Home Blood Pressure Control in People with Diabetes: A Study Protocol for a Crossover Randomized Controlled Trial. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2020 , 13, 4747-4753 | 3.4 | 1 |
| 111 | Immune modulating effects of additional supplementation of estradiol combined with testosterone in murine testosterone-deficient NAFLD model. <i>American Journal of Physiology - Renal Physiology</i> , 2020 , 318, G989-G999 | 5.1 | 10 |
| 110 | Japanese radio calisthenics prevents the reduction of skeletal muscle mass volume in people with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8, | 4.5 | 5 |
| 109 | Changes in metabolic complications in patients with alcoholic fatty liver disease monitored over two decades: NAGALA study. <i>BMJ Open Gastroenterology</i> , 2020 , 7, e000359 | 3.9 | 4 |
| 108 | Intake of sucrose affects gut dysbiosis in patients with type2 diabetes. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1623-1634 | 3.9 | 14 |
| 107 | Handgrip measurement as a useful benchmark for locomotive syndrome in patients with type2 diabetes mellitus: A KAMOGAWA-DM cohort study. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1602-1613 | 3.9 | 3 |
| 106 | Liver Stiffness Is Associated With Progression of Albuminuria in Adults With Type 2 Diabetes: Nonalcoholic Fatty Disease Cohort Study. <i>Canadian Journal of Diabetes</i> , 2020 , 44, 428-433 | 2.1 | 1 |
| 105 | Reduced dietary omega-3 fatty acids intake is associated with sarcopenia in elderly patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 66, 233-237 | 3.1 | 6 |
| 104 | Creatinine to Body Weight Ratio Is Associated with Incident Diabetes: Population-Based Cohort Study. <i>Journal of Clinical Medicine</i> , 2020 , 9, | 5.1 | 2 |
| 103 | Association between Sleep Duration and Incident Chronic Kidney Disease: A Population-Based Cohort Analysis of the NAGALA Study. <i>Kidney and Blood Pressure Research</i> , 2020 , 45, 339-349 | 3.1 | 2 |
| 102 | Association between sleep disorder and quality of life in patients with type 2 diabetes: a cross-sectional study. <i>BMC Endocrine Disorders</i> , 2020 , 20, 98 | 3.3 | 3 |
| 101 | Visceral Adiposity Index is a predictor of incident colorectal cancer: a population-based longitudinal study. <i>BMJ Open Gastroenterology</i> , 2020 , 7, | 3.9 | 4 |
| 100 | Low-attenuation muscle is a predictor of diabetes mellitus: A population-based cohort study. <i>Nutrition</i> , 2020 , 74, 110752 | 4.8 | 12 |
| 99 | Impact of metabolically healthy obesity on the risk of incident gastric cancer: a population-based cohort study. <i>BMC Endocrine Disorders</i> , 2020 , 20, 11 | 3.3 | 9 |
| 98 | Living alone is associated with visit-to-visit HbA1c variability in men but not in women in people with type 2 diabetes: KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2020 , 67, 419-426 | 2.9 | 3 |
| 97 | Limited joint mobility of the hand correlates incident hospitalisation with infection in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2020 , 161, 108049 | 7.4 | 0 |
| 96 | miR-23b-3p acts as a counter-response against skeletal muscle atrophy. <i>Journal of Endocrinology</i> , 2020 , 244, 535-547 | 4.7 | 3 |
| 95 | Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with a Low Prevalence of Sarcopenia in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2020 , 13, | 6.7 | 6 |

| | | | |
|----|---|-----|----|
| 94 | 510-P: Serum Mac-2 Binding Protein Is Associated with Diabetic Micro- and Macroangiopathy in People with Type 2 Diabetes. <i>Diabetes</i> , 2020 , 69, 510-P | 0.9 | |
| 93 | 698-P: Moderate-to-Vigorous Physical Activity Associated with Microalbuminuria in Type 2 Diabetes: A Cross-Sectional Study. <i>Diabetes</i> , 2020 , 69, 698-P | 0.9 | |
| 92 | Effect of probiotics, G9-1, on gastrointestinal symptoms in patients with type 2 diabetes mellitus: study protocol for open-label, single-arm, exploratory research trial (Big STAR study). <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 67, 223-227 | 3.1 | 0 |
| 91 | The visceral adiposity index is a predictor of incident nonalcoholic fatty liver disease: A population-based longitudinal study. <i>Clinics and Research in Hepatology and Gastroenterology</i> , 2020 , 44, 375-383 | 2.4 | 4 |
| 90 | Trunk muscle quality and quantity predict the development of metabolic syndrome and the increase in the number of its components in individuals without metabolic syndrome. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 1161-1168 | 4.5 | 4 |
| 89 | Skipping breakfast is associated with glycemic variability in patients with type 2 diabetes. <i>Nutrition</i> , 2020 , 71, 110639 | 4.8 | 10 |
| 88 | Creatinine-to-bodyweight ratio is a predictor of incident non-alcoholic fatty liver disease: A population-based longitudinal study. <i>Hepatology Research</i> , 2020 , 50, 57-66 | 5.1 | 9 |
| 87 | Relationship between nonalcoholic fatty liver disease and muscle quality as well as quantity evaluated by computed tomography. <i>Liver International</i> , 2020 , 40, 120-130 | 7.9 | 14 |
| 86 | Triglyceride-glucose index (TyG index) is a predictor of incident colorectal cancer: a population-based longitudinal study. <i>BMC Endocrine Disorders</i> , 2020 , 20, 113 | 3.3 | 7 |
| 85 | Eating Fast Is Associated with Nonalcoholic Fatty Liver Disease in Men But Not in Women with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2020 , 12, | 6.7 | 5 |
| 84 | Effect of Exercise Habit on Skeletal Muscle Mass Varies with Protein Intake in Elderly Patients with Type 2 Diabetes: A Retrospective Cohort Study. <i>Nutrients</i> , 2020 , 12, | 6.7 | 6 |
| 83 | Combined effect of hemoglobin and mean corpuscular volume levels on incident metabolic syndrome: A population-based cohort study. <i>Clinical Nutrition ESPEN</i> , 2020 , 40, 314-319 | 1.3 | 0 |
| 82 | Effect of alcohol consumption and the presence of fatty liver on the risk for incident type 2 diabetes: a population-based longitudinal study. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8, | 4.5 | 5 |
| 81 | Eating Fast Has a Significant Impact on Glycemic Excursion in Healthy Women: Randomized Controlled Cross-Over Trial. <i>Nutrients</i> , 2020 , 12, | 6.7 | 4 |
| 80 | The Effects of Metformin on the Gut Microbiota of Patients with Type 2 Diabetes: A Two-Center, Quasi-Experimental Study. <i>Life</i> , 2020 , 10, | 3 | 9 |
| 79 | Relationship between metabolic syndrome and trunk muscle quality as well as quantity evaluated by computed tomography. <i>Clinical Nutrition</i> , 2020 , 39, 1818-1825 | 5.9 | 12 |
| 78 | The Visceral Adiposity Index Is a Predictor of Incident Chronic Kidney Disease: A Population-Based Longitudinal Study. <i>Kidney and Blood Pressure Research</i> , 2020 , 45, 407-418 | 3.1 | 4 |
| 77 | Creatinine/(cystatin C [body weight]) ratio is associated with skeletal muscle mass index. <i>Endocrine Journal</i> , 2020 , 67, 733-740 | 2.9 | 9 |

| | | | |
|----|--|-----|----|
| 76 | Late-night-dinner deteriorates postprandial glucose and insulin whereas consuming dinner dividedly ameliorates them in patients with type 2 diabetes: A randomized crossover clinical trial. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2020 , 29, 68-76 | 1 | 1 |
| 75 | Tomato juice preload has a significant impact on postprandial glucose concentration in healthy women: A randomized cross-over trial. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2020 , 29, 491-497 | 1 | 0 |
| 74 | Platelet to lymphocyte ratio correlates with diabetic foot risk and foot ulcer in patients with type 2 diabetes. <i>Endocrine Journal</i> , 2019 , 66, 905-913 | 2.9 | 10 |
| 73 | Frequent Usage of Convenience Stores is Associated with Low Diet Quality. <i>Nutrients</i> , 2019 , 11, | 6.7 | 8 |
| 72 | High brain natriuretic peptide is associated with sarcopenia in patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2019 , 66, 369-377 | 2.9 | 10 |
| 71 | Protein intake is not associated with progression of diabetic kidney disease in patients without macroalbuminuria. <i>Diabetes/Metabolism Research and Reviews</i> , 2019 , 35, e3150 | 7.5 | 3 |
| 70 | Triglyceride-glucose index is a predictor of incident chronic kidney disease: a population-based longitudinal study. <i>Clinical and Experimental Nephrology</i> , 2019 , 23, 948-955 | 2.5 | 12 |
| 69 | Potential impact of the joint association of total bilirubin and gamma-glutamyltransferase with metabolic syndrome. <i>Diabetology and Metabolic Syndrome</i> , 2019 , 11, 12 | 5.6 | 7 |
| 68 | Understanding of antidiabetic medication is associated with blood glucose in patients with type 2 diabetes: At baseline date of the KAMOGAWA-DM cohort study. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 458-465 | 3.9 | 4 |
| 67 | The sodium-glucose cotransporter 2 inhibitor luseoglitazone can suppress muscle atrophy in Db/Db mice by suppressing the expression of. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2019 , 65, 23-28 | 3.1 | 10 |
| 66 | Malnutrition assessed by controlling nutritional status is correlated to carotid atherosclerosis in patients with type 2 diabetes. <i>Endocrine Journal</i> , 2019 , 66, 1073-1082 | 2.9 | 3 |
| 65 | Non-alcoholic fatty liver disease with obesity as an independent predictor for incident gastric and colorectal cancer: a population-based longitudinal study. <i>BMJ Open Gastroenterology</i> , 2019 , 6, e000295 | 3.9 | 19 |
| 64 | The Triglyceride and Glucose Index Is a Predictor of Incident Nonalcoholic Fatty Liver Disease: A Population-Based Cohort Study. <i>Canadian Journal of Gastroenterology and Hepatology</i> , 2019 , 2019, 5121574 | 2.8 | 26 |
| 63 | Effects of dietary salt restriction on home blood pressure in diabetic patients with excessive salt intake: a pilot study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2019 , 65, 252-257 | 3.1 | 2 |
| 62 | Short sleep duration is a risk of incident nonalcoholic fatty liver disease: a population-based longitudinal study. <i>Journal of Gastrointestinal and Liver Diseases</i> , 2019 , 28, 73-81 | 1.4 | 16 |
| 61 | 1937-P: miR-23b-3p Upregulates Glucose Uptake in C2C12 Myotube Cells by Regulating the Expression of PTEN. <i>Diabetes</i> , 2019 , 68, 1937-P | 0.9 | |
| 60 | 1943-P: The Sodium-Glucose Cotransporter 2 Inhibitor Luseoglitazone Can Suppress Muscle Atrophy in Db/Db Mice by Suppressing the Expression of Foxo1. <i>Diabetes</i> , 2019 , 68, 1943-P | 0.9 | |
| 59 | 752-P: Japanese Radio Calisthenics Prevents Reduction of Skeletal Muscle Volume in Patients with Type 2 Diabetes. <i>Diabetes</i> , 2019 , 68, 752-P | 0.9 | 1 |

| | | | |
|----|--|-----|----------------|
| 58 | Evaluation of the efficacy of simplified nutritional instructions from physicians on dietary salt restriction for patients with type 2 diabetes mellitus consuming excessive salt: protocol for a randomized controlled trial. <i>Trials</i> , 2019 , 20, 761 | 2.8 | 0 |
| 57 | Reduction of Fat to Muscle Mass Ratio Is Associated with Improvement of Liver Stiffness in Diabetic Patients with Non-Alcoholic Fatty Liver Disease. <i>Journal of Clinical Medicine</i> , 2019 , 8, | 5.1 | 8 |
| 56 | Shortage of energy intake rather than protein intake is associated with sarcopenia in elderly patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort. <i>Journal of Diabetes</i> , 2019 , 11, 477-483 | 3.8 | 29 |
| 55 | Impact of different timing of consuming sweet snack on postprandial glucose excursions in healthy women. <i>Diabetes and Metabolism</i> , 2019 , 45, 369-374 | 5.4 | 7 |
| 54 | Sarcopenia is associated with tongue pressure in older patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort study. <i>Geriatrics and Gerontology International</i> , 2019 , 19, 153-158 | 2.9 | 22 |
| 53 | Ectopic fat obesity presents the greatest risk for incident type 2 diabetes: a population-based longitudinal study. <i>International Journal of Obesity</i> , 2019 , 43, 139-148 | 5.5 | 77 |
| 52 | Decreased the creatinine to cystatin C ratio is a surrogate marker of sarcopenia in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018 , 139, 52-58 | 7.4 | 54 |
| 51 | Divided consumption of late-night-dinner improves glucose excursions in young healthy women: A randomized cross-over clinical trial. <i>Diabetes Research and Clinical Practice</i> , 2018 , 136, 78-84 | 7.4 | 6 |
| 50 | Metabolically healthy obesity without fatty liver and risk of incident type 2 diabetes: A meta-analysis of prospective cohort studies. <i>Obesity Research and Clinical Practice</i> , 2018 , 12, 4-15 | 5.4 | 22 |
| 49 | The Association Between Taste Impairment and Serum Zinc Concentration in Adult Patients With Type 2 Diabetes. <i>Canadian Journal of Diabetes</i> , 2018 , 42, 520-524 | 2.1 | 3 |
| 48 | Asia-Pacific Working Party on Non-alcoholic Fatty Liver Disease guidelines 2017-Part 1: Definition, risk factors and assessment. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2018 , 33, 70-85 | 4 | 211 |
| 47 | The Asia-Pacific Working Party on Non-alcoholic Fatty Liver Disease guidelines 2017-Part 2: Management and special groups. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2018 , 33, 86-98 ⁴ | | 7 ¹ |
| 46 | Impact of respiratory function on the progression from metabolically healthy non-overweight to metabolically abnormal phenotype. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018 , 28, 922-928 ^{4.5} | | 4 |
| 45 | Low urine pH is a risk for non-alcoholic fatty liver disease: A population-based longitudinal study. <i>Clinics and Research in Hepatology and Gastroenterology</i> , 2018 , 42, 570-576 | 2.4 | 6 |
| 44 | Consuming snacks mid-afternoon compared with just after lunch improves mean amplitude of glycaemic excursions in patients with type 2 diabetes: A randomized crossover clinical trial. <i>Diabetes and Metabolism</i> , 2018 , 44, 482-487 | 5.4 | 1 |
| 43 | Sarcopenia is associated with blood pressure variability in older patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort study. <i>Geriatrics and Gerontology International</i> , 2018 , 18, 1345-1349 | 2.9 | 23 |
| 42 | Neutrophil-lymphocyte ratio correlates with limited joint mobility of hand in patients with type 2 diabetes. <i>Endocrine Journal</i> , 2018 , 65, 1011-1017 | 2.9 | 2 |
| 41 | Metabolically healthy obesity and risk of leukoaraiosis; a population based cross-sectional study. <i>Endocrine Journal</i> , 2018 , 65, 669-675 | 2.9 | 6 |

| | | | |
|----|--|-----|----|
| 40 | Intake of Carbohydrate to Fiber Ratio Is a Useful Marker for Metabolic Syndrome in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Annals of Nutrition and Metabolism</i> , 2018 , 72, 329-335 | 4.5 | 9 |
| 39 | Serum N-terminal Pro-brain Natriuretic Peptide Level is Associated with the Development of Chronic Kidney Diseases in Patients with Type 2 Diabetes. <i>Endocrine, Metabolic and Immune Disorders - Drug Targets</i> , 2018 , 18, 590-595 | 2.2 | 3 |
| 38 | Late-night-dinner is associated with poor glycemic control in people with type 2 diabetes: The KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2018 , 65, 395-402 | 2.9 | 39 |
| 37 | Urinary pH is a predictor of diabetes in men; a population based large scale cohort study. <i>Diabetes Research and Clinical Practice</i> , 2017 , 130, 9-14 | 7.4 | 16 |
| 36 | Fatty liver as a risk factor for progression from metabolically healthy to metabolically abnormal in non-overweight individuals. <i>Endocrine</i> , 2017 , 57, 89-97 | 4 | 32 |
| 35 | Divided consumption of late-night-dinner improves glycemic excursions in patients with type 2 diabetes: A randomized cross-over clinical trial. <i>Diabetes Research and Clinical Practice</i> , 2017 , 129, 206-212 | 7.4 | 16 |
| 34 | Weight gain since age of 20 as risk of metabolic syndrome even in non-overweight individuals. <i>Endocrine</i> , 2017 , 58, 253-261 | 4 | 15 |
| 33 | Impact of fatty liver disease and metabolic syndrome on incident type 2 diabetes; a population based cohort study. <i>Endocrine Journal</i> , 2017 , 64, 1105-1114 | 2.9 | 6 |
| 32 | Urinary pH reflects dietary acid load in patients with type 2 diabetes. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2017 , 61, 74-77 | 3.1 | 11 |
| 31 | Combined effect of body mass index and waist-height ratio on incident diabetes; a population based cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2017 , 61, 118-122 | 3.1 | 15 |
| 30 | Relationship between limited joint mobility of hand and carotid atherosclerosis in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2017 , 132, 79-84 | 7.4 | 6 |
| 29 | Relationship between limited joint mobility of the hand and diabetic foot risk in patients with type 2 diabetes. <i>Journal of Diabetes</i> , 2017 , 9, 628-633 | 3.8 | 12 |
| 28 | Relationship between skeletal muscle mass and hepatic fibrosis in patients with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2017 , 43, 184-186 | 5.4 | 11 |
| 27 | Protein Intake, Especially Vegetable Protein Intake, Is Associated with Higher Skeletal Muscle Mass in Elderly Patients with Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2017 , 2017, 7985728 | 3.9 | 12 |
| 26 | Nonoverweight nonalcoholic fatty liver disease and incident cardiovascular disease: A post hoc analysis of a cohort study. <i>Medicine (United States)</i> , 2017 , 96, e6712 | 1.8 | 44 |
| 25 | The relationship between hepatic steatosis and skeletal muscle mass index in men with type 2 diabetes. <i>Endocrine Journal</i> , 2016 , 63, 877-884 | 2.9 | 40 |
| 24 | Caffeine intake enhances the benefits of sodium glucose transporter 2 inhibitor. <i>Diabetes/Metabolism Research and Reviews</i> , 2016 , 32, 694-699 | 7.5 | 3 |
| 23 | BMI history and risk of incident fatty liver: a population-based large-scale cohort study. <i>European Journal of Gastroenterology and Hepatology</i> , 2016 , 28, 1188-93 | 2.2 | 15 |

| | | | |
|----|---|------|-----|
| 22 | Transient remission of nonalcoholic fatty liver disease decreases the risk of incident type 2 diabetes mellitus in Japanese men. <i>European Journal of Gastroenterology and Hepatology</i> , 2016 , 28, 1443-1449 | 2.2 | 26 |
| 21 | Triglycerides to high-density lipoprotein cholesterol ratio is an independent predictor of incident fatty liver; a population-based cohort study. <i>Liver International</i> , 2016 , 36, 713-20 | 7.9 | 47 |
| 20 | Impact of low-carbohydrate diet on body composition: meta-analysis of randomized controlled studies. <i>Obesity Reviews</i> , 2016 , 17, 499-509 | 10.6 | 53 |
| 19 | Impact of low-carbohydrate diet on renal function: a meta-analysis of over 1000 individuals from nine randomised controlled trials. <i>British Journal of Nutrition</i> , 2016 , 116, 632-8 | 3.6 | 26 |
| 18 | The impact of non-alcoholic fatty liver disease on incident type 2 diabetes mellitus in non-overweight individuals. <i>Liver International</i> , 2016 , 36, 275-83 | 7.9 | 90 |
| 17 | Sodium-chloride Difference and Metabolic Syndrome: A Population-based Large-scale Cohort Study. <i>Internal Medicine</i> , 2016 , 55, 3085-3090 | 1.1 | 15 |
| 16 | Which Measurement of Blood Pressure Is More Associated With Albuminuria in Patients With Type 2 Diabetes: Central Blood Pressure or Peripheral Blood Pressure?. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 790-5 | 2.3 | 4 |
| 15 | Heart rate-corrected QT interval is a novel risk marker for the progression of albuminuria in people with Type2 diabetes. <i>Diabetic Medicine</i> , 2015 , 32, 1221-6 | 3.5 | 6 |
| 14 | Metabolically healthy obesity and risk of incident CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2015 , 10, 578-83 | 6.9 | 94 |
| 13 | Hemoglobin concentration and incident metabolic syndrome: a population-based large-scale cohort study. <i>Endocrine</i> , 2015 , 50, 390-6 | 4 | 26 |
| 12 | Modest alcohol consumption reduces the incidence of fatty liver in men: a population-based large-scale cohort study. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2015 , 30, 546-52 | 4 | 66 |
| 11 | Lower vegetable protein intake and higher dietary acid load associated with lower carbohydrate intake are risk factors for metabolic syndrome in patients with type2 diabetes: Post-hoc analysis of a cross-sectional study. <i>Journal of Diabetes Investigation</i> , 2015 , 6, 465-72 | 3.9 | 24 |
| 10 | High-sensitivity cardiac troponin T is associated with coronary artery calcification. <i>Journal of Cardiovascular Computed Tomography</i> , 2015 , 9, 209-14 | 2.8 | 5 |
| 9 | Association of non-alcoholic fatty liver disease with chronic kidney disease: a systematic review and meta-analysis. <i>PLoS Medicine</i> , 2014 , 11, e1001680 | 11.6 | 363 |
| 8 | A case of diabetic ketoacidosis complicated by fatal acute abdominal aortic thrombosis. <i>Diabetology International</i> , 2013 , 4, 201-204 | 2.3 | 2 |
| 7 | Aging is a risk factor of nonalcoholic fatty liver disease in premenopausal women. <i>World Journal of Gastroenterology</i> , 2012 , 18, 237-43 | 5.6 | 86 |
| 6 | Identification of individuals with non-alcoholic fatty liver disease by the diagnostic criteria for the metabolic syndrome. <i>World Journal of Gastroenterology</i> , 2012 , 18, 1508-16 | 5.6 | 67 |
| 5 | Nonalcoholic fatty liver disease is a novel predictor of cardiovascular disease. <i>World Journal of Gastroenterology</i> , 2007 , 13, 1579-84 | 5.6 | 415 |

| | | | |
|---|---|-----|-----|
| 4 | The severity of ultrasonographic findings in nonalcoholic fatty liver disease reflects the metabolic syndrome and visceral fat accumulation. <i>American Journal of Gastroenterology</i> , 2007 , 102, 2708-15 | 0.7 | 514 |
| 3 | The metabolic syndrome as a predictor of nonalcoholic fatty liver disease. <i>Annals of Internal Medicine</i> , 2005 , 143, 722-8 | 8 | 739 |
| 2 | Reduction by phytate-reduced soybean beta-conglycinin of plasma triglyceride level of young and adult rats. <i>Bioscience, Biotechnology and Biochemistry</i> , 2001 , 65, 1071-5 | 2.1 | 60 |
| 1 | Soy protein isolate and its hydrolysate reduce body fat of dietary obese rats and genetically obese mice (yellow KK). <i>Nutrition</i> , 2000 , 16, 349-54 | 4.8 | 149 |