

Athanassios K Boudalis

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6605229/athanassios-k-boudalis-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

85
papers

2,277
citations

26
h-index

44
g-index

95
ext. papers

2,380
ext. citations

3.8
avg. IF

4.51
L-index

#	Paper	IF	Citations
85	Room-temperature spin nutations in a magnetically condensed phase of [Y(pc)] \square <i>Chemical Communications</i> , 2021 , 57, 11505-11508	5.8	1
84	Room-Temperature Cu(II) Radical-Triggered Alkyne C-H Activation. <i>Jacs Au</i> , 2021 , 1, 1937-1948		0
83	Half-Integer Spin Triangles: Old Dogs, New Tricks. <i>Chemistry - A European Journal</i> , 2021 , 27, 7022-7042	4.8	5
82	Observation and deconvolution of a unique EPR signal from two cocrystallized spin triangles. <i>Physical Chemistry Chemical Physics</i> , 2021 , 23, 14415-14421	3.6	
81	Broadband electron paramagnetic resonance of a molecular spin triangle. <i>Physical Chemistry Chemical Physics</i> , 2021 , 23, 20268-20274	3.6	
80	Supramolecular Assemblies of Trinuclear Copper(II)-Pyrazolato Units: A Structural, Magnetic and EPR Study. <i>Chemistry</i> , 2020 , 2, 626-644	2.1	2
79	Origin of Ferromagnetism and Magnetic Anisotropy in a Family of Copper(II) Triangles. <i>Chemistry - A European Journal</i> , 2020 , 26, 12769-12784	4.8	5
78	Chromium(III)-pyrazole complexes. X-Ray crystal structures, 1H NMR investigation of ligand fluxional behavior and EPR studies. <i>Inorganica Chimica Acta</i> , 2020 , 502, 119299	2.7	2
77	Nitrite Reduction by Trinuclear Copper Pyrazolate Complexes: An Example of a Catalytic, Synthetic Polynuclear NO Releasing System. <i>Inorganic Chemistry</i> , 2019 , 58, 7537-7544	5.1	13
76	Relevance of Dzyaloshinskii-Moriya spectral broadenings in promoting spin decoherence: a comparative pulsed-EPR study of two structurally related iron(iii) and chromium(iii) spin-triangle molecular qubits. <i>Physical Chemistry Chemical Physics</i> , 2019 , 21, 19575-19584	3.6	3
75	Polyanisotropic Magnetoelectric Coupling in an Electrically Controlled Molecular Spin Qubit. <i>Journal of the American Chemical Society</i> , 2019 , 141, 19765-19775	16.4	9
74	The [Fe{(SePPh ₂) ₂ N} ₂] Complex Revisited: X-ray Crystallography, Magnetometry, High-Frequency EPR, and Mössbauer Studies Reveal Its Tetrahedral FeIISe ₄ Coordination Sphere. <i>European Journal of Inorganic Chemistry</i> , 2018 , 2018, 713-721	2.3	5
73	Interactions between H-bonded [Cu(EOH)] triangles; a combined magnetic susceptibility and EPR study. <i>Physical Chemistry Chemical Physics</i> , 2018 , 20, 17234-17244	3.6	13
72	Determination of the Distributions of the Spin-Hamiltonian Parameters in Spin Triangles: A Combined Magnetic Susceptometry and Electron Paramagnetic Resonance Spectroscopic Study of the Highly Symmetric [CrO(PhCOO)(py)](ClO ₄) \cdot 5py. <i>Inorganic Chemistry</i> , 2018 , 57, 13259-13269	5.1	10
71	First Demonstration of Magnetoelectric Coupling in a Polynuclear Molecular Nanomagnet: Single-Crystal EPR Studies of [Fe O(O CPh) (py)]ClO ₄ py under Static Electric Fields. <i>Chemistry - A European Journal</i> , 2018 , 24, 14896-14900	4.8	27
70	Dynamic versus Static Character of the Magnetic Jahn-Teller Effect: Magnetostructural Studies of [FeO(OCPh)(py)]ClO ₄ py. <i>Inorganic Chemistry</i> , 2017 , 56, 762-772	5.1	14
69	Towards ionic liquids with tailored magnetic properties: bmim salts of ferro- and antiferromagnetic Cu triangles. <i>Dalton Transactions</i> , 2017 , 46, 12263-12273	4.3	14

68	Site preferences in hetero-metallic [FeNi] clusters: a combined crystallographic, spectroscopic and theoretical analysis. <i>Dalton Transactions</i> , 2017 , 46, 12835-12844	4.3	3
67	Hexanuclear complexes from the use of a series of amino-alcohol ligands in Fe chemistry. <i>Polyhedron</i> , 2013 , 64, 218-230	2.7	7
66	Azide-bridged manganese(III) one-dimensional chain: synthesis, structure, and magnetic study. <i>Journal of Coordination Chemistry</i> , 2013 , 66, 9-17	1.6	10
65	Crystal structure and topological ferrimagnetic behavior of a new 2D metal-organic hybrid manganese complex $[Mn_3(N_3)_4(L)_2(H_2O)_2]n \cdot 6(C_2H_5OH) \cdot 4(H_2O)$ with the AF/AF/F alternating sequence (HL = N-((pyridine-2-yl)methylene)isonicotinohydrazide). <i>Polyhedron</i> , 2013 , 49, 61-66	2.7	10
64	Polymerization of a preformed Mn ₆ cluster to a one-dimensional chain. <i>Polyhedron</i> , 2013 , 52, 917-923	2.7	6
63	The first member of a second generation family of ligands derived from metal-ion assisted reactivity of di-2,6-(2-pyridylcarbonyl)pyridine: Synthesis and characterization of a MnII/III ₄ rhombus. <i>Inorganic Chemistry Communication</i> , 2012 , 15, 73-77	3.1	14
62	Pentanuclear complexes with unusual structural topologies from the initial use of two aliphatic amino-alcohol ligands in Fe chemistry. <i>Dalton Transactions</i> , 2012 , 41, 1544-52	4.3	12
61	Iron(III) chemistry with ferrocene-1,1'-dicarboxylic acid (fdch ₂): an Fe ₇ cluster with an oxidized fdc(-) ligand. <i>Inorganic Chemistry</i> , 2011 , 50, 5646-54	5.1	16
60	The first non-acetato members of the bis(anion)octacarboxylatotetrakis{di-2-pyridyl-methanediolate(2-)}enneametal(II) family of complexes: Synthesis, X-ray structures and magnetism of $[M_9(N_3)_2(O_2CCMe_3)_8\{(py)_2CO_2\}_4]$ (M = Co, Ni). <i>Polyhedron</i> , 2011 , 30, 3026-3033	2.7	13
59	Magnetic relaxation in basic iron(III) carboxylate $[Fe_3O(O_2CPh)_6(H_2O)_3]ClO_4 \cdot py$. <i>Dalton Transactions</i> , 2011 , 40, 6371-4	4.3	16
58	Structural variation from 1D chains to 3D networks: a systematic study of coordination number effect on the construction of coordination polymers using the terephthaloylbisglycinate ligand. <i>New Journal of Chemistry</i> , 2011 , 35, 1060	3.6	25
57	Expanding the 3d-4f heterometallic chemistry of the (py) ₂ CO and pyCOpyCOpy ligands: structural, magnetic and Mössbauer spectroscopic studies of two Fe(II)-Gd(III) complexes. <i>Dalton Transactions</i> , 2011 , 40, 8199-205	4.3	14
56	Coordination Polymers of Lanthanide(III): Toward Encapsulation of Well-Defined Assembly of Guest Molecules. <i>Crystal Growth and Design</i> , 2011 , 11, 547-554	3.5	24
55	Copper(II)-mediated oxime-tritrile coupling in non-aqueous solutions: Synthetic, structural and magnetic studies of the copper(II)-alicylaldehyde oxime reaction system. <i>Inorganica Chimica Acta</i> , 2011 , 370, 50-58	2.7	9
54	Isomorphous replacement of M(II) ions in M(II)-Gd(III) dimers (M(II) = Cu(II), Mn(II), Ni(II), Co(II), Zn(II)): magnetic studies of the products. <i>Dalton Transactions</i> , 2010 , 39, 5020-7	4.3	44
53	Mössbauer spectra of two extended series of basic iron(III) carboxylates $[Fe_3O(O_2CR)_6(H_2O)_6]A$ (A = ClO ₄ , NO ₃). <i>Hyperfine Interactions</i> , 2010 , 198, 229-241	0.8	11
52	Cluster based copper(II)-azide polymer: Synthesis, structure and magnetic study. <i>Polyhedron</i> , 2010 , 29, 3342-3348	2.7	24
51	Aromatic -oxide bridged copper(II) coordination polymers: Synthesis, characterization and magnetic properties. <i>Inorganica Chimica Acta</i> , 2010 , 363, 2279-2286	2.7	25

50	Selective adenine/cytosine cations in one-dimensional coordination polymers of manganese (II) and copper (II) 2,3-pyridinedicarboxylates. <i>Inorganic Chemistry Communication</i> , 2010 , 13, 1244-1248	3.1	12
49	Maleamate(II) and Maleate(II) Copper(II)2,2'-Bipyridine Complexes: Synthesis, Reactivity and Structural and Physical Studies. <i>European Journal of Inorganic Chemistry</i> , 2009 , 2009, 4554-4563	2.3	11
48	2-Pyridyl ketone oximes in iron(III) carboxylate chemistry: Synthesis, structural and physical studies of tetranuclear clusters containing the [Fe4(β-O)2]8+ Butterfly Core. <i>Polyhedron</i> , 2009 , 28, 3221-3226	2.7	9
47	A diferric complex from metal-assisted methanolysis of di-2,6-(2-pyridylcarbonyl)-pyridine: Structural, magnetic and spectroscopic (Mössbauer, EPR) studies. <i>Polyhedron</i> , 2009 , 28, 3251-3256	2.7	8
46	Initial use of 1,1'-oxalyldiimidazole for inorganic synthesis: Decomposition of the ligand as a means to the preparation of an imidazole- and oxalate(-2)-containing, 1D copper(II) complex. <i>Inorganic Chemistry Communication</i> , 2009 , 12, 402-405	3.1	7
45	Ferromagnetic Cu(II)4, Co(II)4, and Ni(II)6 azido complexes derived from metal-assisted methanolysis of di-2,6-(2-pyridylcarbonyl)pyridine. <i>Inorganic Chemistry</i> , 2009 , 48, 3167-76	5.1	76
44	Alcoholysis/hydrolysis of 1,1'-carbonyldiimidazole as a means of preparing unprecedented, imidazole-containing one-dimensional coordination polymers of copper(II). <i>Dalton Transactions</i> , 2009 , 3354-62	4.3	17
43	A metamagnetic 2D copper(ii)-azide complex with 1D ferromagnetism and a hysteretic spin-flop transition. <i>Dalton Transactions</i> , 2009 , 3215-21	4.3	56
42	Reversible core-interconversion of an iron(III) dihydroxo bridged complex. <i>Inorganic Chemistry</i> , 2008 , 47, 11314-23	5.1	13
41	Slow magnetic relaxation of a ferromagnetic Ni(II)5 cluster with an S = 5 ground state. <i>Inorganic Chemistry</i> , 2008 , 47, 10674-81	5.1	53
40	A family of enneanuclear iron(II) single-molecule magnets. <i>Chemistry - A European Journal</i> , 2008 , 14, 2514-86	4.3	102
39	A Tetragonal Core with Asymmetric Iron Environments Supported Solely by Bis(OH){(O)H} Bridging and Terminal Pyridine Amide (N, O) Coordination: A New Member of the Tetrairon(III) Family. <i>European Journal of Inorganic Chemistry</i> , 2008 , 2008, 2820-2829	2.3	11
38	Ferromagnetism in CuII4 and CoII4 Complexes Derived from Metal-Assisted Solvolysis of Di-2,6-(2-pyridylcarbonyl)pyridine: Syntheses, Structures, and Magnetic Properties. <i>European Journal of Inorganic Chemistry</i> , 2008 , 2008, 3796-3801	2.3	33
37	Aerobic FeIII/II Reduction in the Presence of 2,4,6-Tri(2-pyridyl)-1,3,5-triazine and Benzilic Acid: Synthesis and Characterization of a Heptacoordinate FeII Nitrate Complex. <i>European Journal of Inorganic Chemistry</i> , 2008 , 2008, 5632-5641	2.3	13
36	A systematic investigation of the CuCl2/H2mal/phen reaction system (H2mal=malonic acid): Solution and solid state studies of its products. <i>Polyhedron</i> , 2008 , 27, 2315-2326	2.7	11
35	Salicylaldoxime in manganese(III) carboxylate chemistry: Synthesis, structural characterization and physical studies of hexanuclear and polymeric complexes. <i>Polyhedron</i> , 2008 , 27, 3575-3586	2.7	24
34	Synthesis, spectroscopic, structural and electrochemical studies of carboxyl substituted 1,4-naphthoquinones. <i>Inorganica Chimica Acta</i> , 2008 , 361, 1681-1688	2.7	23
33	Two new coordination polymers containing the triangular [Mn3O(O2CR)6]0/+ units. <i>Inorganica Chimica Acta</i> , 2008 , 361, 4100-4106	2.7	15

32	An "S"-shaped pentanuclear CuII cluster derived from the metal-assisted hydrolysis of pyCOPyCOPy: structural, magnetic and spectroscopic studies. <i>Dalton Transactions</i> , 2007 , 3582-9	4.3	38
31	Synthesis and characterization of the tetranuclear iron(III) complex of a new asymmetric multidentate ligand. A structural model for purple acid phosphatases. <i>Dalton Transactions</i> , 2007 , 5132-9	4.3	30
30	Structure and properties of dinuclear manganese(III) complexes with pentaanionic pentadentate ligands including alkoxo, amido, and phenoxo donors. <i>Inorganic Chemistry</i> , 2007 , 46, 6902-10	5.1	13
29	A Diferrous Single-Molecule Magnet. <i>European Journal of Inorganic Chemistry</i> , 2007 , 2007, 2409-2415	2.3	18
28	Mixed-Valence Cobalt(II/III) Carboxylate Clusters: CoII4CoIII2 and CoII2CoIII2 Complexes from the Use of 2-(Hydroxymethyl)pyridine. <i>European Journal of Inorganic Chemistry</i> , 2007 , 2007, 5098-5104	2.3	45
27	The benzoate/nitrate/2,2':6',2''-terpyridine Blend in lanthanide(III) chemistry: Relevance to the separation of lanthanides and actinides by solvent extraction. <i>Inorganic Chemistry Communication</i> , 2007 , 10, 800-804	3.1	13
26	Ligand-induced electronic effects in the physical properties of transition metal clusters: Modulation of Mössbauer parameters and magnetic exchange couplings in complexes by the R substituent. <i>Inorganica Chimica Acta</i> , 2007 , 360, 39-47	2.7	11
25	A general synthetic route for the preparation of high-spin molecules: Replacement of bridging hydroxo ligands in molecular clusters by end-on azido ligands. <i>Polyhedron</i> , 2007 , 26, 2089-2094	2.7	24
24	Weak ferromagnetism in 1D coordination polymers: Syntheses, crystal structures, spectroscopic and magnetic properties of [Cu(L)(Memal)]n (Memal ²⁻ the dianion of methylmalonic acid, L=1,10-phenanthroline, 2,2'-bipyridine). <i>Polyhedron</i> , 2007 , 26, 2536-2542	2.7	21
23	J-strain and antisymmetric exchange in a polynuclear compound containing the {Fe3O}7+ core. <i>Comptes Rendus Chimie</i> , 2007 , 10, 116-124	2.7	12
22	Co(II) chemistry of 2,6-bis(2-pyridylcarbonyl)pyridine: an icosanuclear Co cluster exhibiting superparamagnetic relaxation. <i>Angewandte Chemie - International Edition</i> , 2006 , 45, 432-5	16.4	110
21	Syntheses, Structural, and Physical Studies of Basic CrIII and FeIII Benzilates and Benzoates: Evidence of Antisymmetric Exchange and Distributions of Isotropic and Antisymmetric Exchange Parameters. <i>European Journal of Inorganic Chemistry</i> , 2006 , 2006, 3710-3723	2.3	40
20	CoII Chemistry of 2,6-Bis(2-pyridylcarbonyl)pyridine: An Icosanuclear Co Cluster Exhibiting Superparamagnetic Relaxation. <i>Angewandte Chemie</i> , 2006 , 118, 446-449	3.6	4
19	Reactivity and structural and physical studies of tetranuclear iron(III) clusters containing the [Fe4(μ ₃ -O) ₂] ⁸⁺ "butterfly" core: an FeIII ₄ cluster with an S = 1 ground state. <i>Inorganic Chemistry</i> , 2006 , 45, 7372-81	5.1	34
18	Hexanuclear iron(III) salicylaldoximate complexes presenting the [Fe ₆ (μ ₃ -O) ₂ (μ ₂ -OR) ₂] ¹²⁺ core: syntheses, crystal structures, and spectroscopic and magnetic characterization. <i>Inorganic Chemistry</i> , 2006 , 45, 2317-26	5.1	47
17	An octanuclear complex containing the [Fe ₃ O] ⁷⁺ metal core: structural, magnetic, Mössbauer, and electron paramagnetic resonance studies. <i>Inorganic Chemistry</i> , 2006 , 45, 443-53	5.1	57
16	Formation of the core in copper(II) carboxylate chemistry via use of di-2-pyridyl ketone oxime [(py) ₂ CNOH]:[Cu ₃ (OH)(O ₂ CR) ₂ {(py) ₂ CNO} ₃] (R=Me, Ph). <i>Inorganic Chemistry Communication</i> , 2006 , 9, 814-818	3.1	63
15	Further investigations on the Fe(ClO ₄) ₃ /RCO ₂ -/phen reaction system: Syntheses, structural and physical characterization of complexes [Fe ₂ O(O ₂ CCCl ₃) ₂ (phen) ₂ (H ₂ O) ₂](ClO ₄) ₂ and [Fe ₂ O(O ₂ CCMe ₃) ₂ (phen) ₂ (H ₂ O) ₂](ClO ₄) ₂ . <i>Polyhedron</i> , 2006 , 25, 1391-1398	2.7	8

14	Cation-dependent nuclearity of the copper-azido moiety: synthesis, structure, and magnetic study. <i>Inorganic Chemistry</i> , 2005 , 44, 6379-85	5.1	67
13	Salicylaldoxime (H ₂ salox) in iron(III) carboxylate chemistry: Synthesis, X-ray crystal structure, spectroscopic characterization and magnetic behavior of trinuclear oxo-centered complexes. <i>Polyhedron</i> , 2005 , 24, 711-721	2.7	48
12	A trinuclear cluster containing the {Fe ₃ (μ -O)} ⁷⁺ core: Structural, magnetic and spectroscopic (IR, Mössbauer, EPR) studies. <i>Polyhedron</i> , 2005 , 24, 1540-1548	2.7	43
11	Hydrogen-Bonded Networks Featuring Yttrium(III) Complexes of N,N-Dimethylurea (DMU): Preparation and Characterization of [Y(DMU) ₆][YCl ₆] and [Y(NO ₃) ₃ (DMU) ₃]. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2005 , 60, 363-372	1	4
10	New poly-iron(II) complexes of N ₄ O dinucleating Schiff bases and pseudohalides: syntheses, structures, and magnetic and Mössbauer properties. <i>Inorganic Chemistry</i> , 2004 , 43, 1574-86	5.1	102
9	A nonanuclear iron(II) single-molecule magnet. <i>Angewandte Chemie - International Edition</i> , 2004 , 43, 2266-70	5.1	196
8	A Nonanuclear Iron(II) Single-Molecule Magnet. <i>Angewandte Chemie</i> , 2004 , 116, 2316-2320	3.6	33
7	A new example of a tetranuclear iron(III) cluster containing the [Fe ₄ O ₂] ⁸⁺ core: preparation, X-ray crystal structure, magnetochemistry and Mössbauer study of [Fe ₄ O ₂ (O ₂ CMe) ₆ (N ₃) ₂ (phen) ₂]. <i>Inorganica Chimica Acta</i> , 2004 , 357, 1345-1354	2.7	39
6	2,2'-Bipyrimidine (bpym) in iron(III) carboxylate chemistry: preparation and characterization of [Fe ₂ O(O ₂ CMe) ₂ (bpym) ₂ (H ₂ O) ₂](ClO ₄) ₂ , and the influence of weak interactions in the formation of metal clusters. <i>Polyhedron</i> , 2004 , 23, 1271-1277	2.7	20
5	Use of the di-2-pyridyl ketone/3,5-di-tert-butylcatechol blend in iron(III) chemistry: a cationic tetranuclear cluster and an anionic trinuclear complex. <i>Dalton Transactions</i> , 2003 , 3411-3418	4.3	54
4	Novel rectangular [Fe ₄ (μ ₄ -OHO)(μ -OH) ₂](7+) versus "butterfly" [Fe ₄ (μ ₃ -O ₂)](8+) core topology in the Fe(III)/RCO ₂ (-)/phen reaction systems (R = Me, Ph; phen = 1,10-phenanthroline): preparation and properties of [Fe ₄ (OHO)(OH) ₂ (O ₂ CMe) ₄ (phen) ₄](ClO ₄) ₃ , [Fe ₄ O ₂ (O ₂ CPh) ₇ (phen) ₂](ClO ₄) ₄ , and [Fe ₄ O ₂ (O ₂ CPh) ₈ (phen) ₂]. <i>Inorganic Chemistry</i> , 2002 , 41, 6474-87	5.1	81
3	Reactions of 2,2'-bipyridine (bpy) and 1,10-phenanthroline (phen) with yttrium(III) nitrate: preparation, X-ray crystal structures and spectroscopic characterization of the bis-bpy and bis-phen complexes. <i>Transition Metal Chemistry</i> , 2001 , 26, 276-281	2.1	65
2	Reaction between Yttrium Nitrate and 2,2':6',2"-Terpyridine (terpy) in MeCN: Preparation, Crystal Structures and Spectroscopic Characterization of [Y(NO ₃) ₃ (terpy)(H ₂ O)] and [Y(NO ₃) ₃ (terpy)(H ₂ O)] ₃ MeCN. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2001 , 56, 122-128	1	14
1	Tetranuclear iron(III) carboxylate clusters with 1,10-phenanthroline and 2,2'-bipyridine: a new [Fe ₄ (μ -OHO)(μ -OH) ₂] ⁷⁺ core. <i>Dalton Transactions RSC</i> , 2001 , 955-957		23