Margareta Nyman

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6605322/publications.pdf

Version: 2024-02-01

117453 114278 4,278 85 34 63 citations g-index h-index papers 86 86 86 5936 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Rapid determination of short-chain fatty acids in colonic contents and faeces of humans and rats by acidified water-extraction and direct-injection gas chromatography. Biomedical Chromatography, 2006, 20, 674-682.	0.8	397
2	Contribution of diet to the composition of the human gut microbiota. Microbial Ecology in Health and Disease, 2015, 26, 26164.	3.8	310
3	High-Fat Diet Reduces the Formation of Butyrate, but Increases Succinate, Inflammation, Liver Fat and Cholesterol in Rats, while Dietary Fibre Counteracts These Effects. PLoS ONE, 2013, 8, e80476.	1.1	249
4	Fermentation of dietary fibre components in the rat intestinal tract. British Journal of Nutrition, 1982, 47, 357-366.	1.2	206
5	Modulation of gut microbiota in rats fed highâ€fat diets by processing wholeâ€grain barley to barley malt. Molecular Nutrition and Food Research, 2015, 59, 2066-2076.	1.5	181
6	Fermentation of dietary fibre in the intestinal tract: comparison between man and rat. British Journal of Nutrition, 1986, 55, 487-496.	1.2	169
7	Survival of Lactobacillus plantarum DSM 9843 (299v), and effect on the short-chain fatty acid content of faeces after ingestion of a rose-hip drink with fermented oats. International Journal of Food Microbiology, 1998, 42, 29-38.	2.1	161
8	Binding of Cu2+, Zn2+, and Cd2+to Inositol Tri-, Tetra-, Penta-, and Hexaphosphates. Journal of Agricultural and Food Chemistry, 1998, 46, 3194-3200.	2.4	153
9	Fermentation and bulking capacity of indigestible carbohydrates: the case of inulin and oligofructose. British Journal of Nutrition, 2002, 87, S163-S168.	1.2	116
10	Variation in the content of glucosinolates, hydroxycinnamic acids, carotenoids, total antioxidant capacity and low-molecular-weight carbohydrates inBrassica vegetables. Journal of the Science of Food and Agriculture, 2006, 86, 528-538.	1.7	106
11	Distribution and characterisation of fructan in wheat milling fractions. Journal of Cereal Science, 2008, 48, 768-774.	1.8	105
12	Determination of short-chain fatty acids in serum by hollow fiber supported liquid membrane extraction coupled with gas chromatography. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2007, 846, 202-208.	1.2	95
13	Effects of two whole-grain barley varieties on caecal SCFA, gut microbiota and plasma inflammatory markers in rats consuming low- and high-fat diets. British Journal of Nutrition, 2015, 113, 1558-1570.	1.2	93
14	Changes in carbohydrate and glucosinolate composition in white cabbage (Brassica oleracea var.) Tj ETQq0 0 0 r	gBŢ.¦Over	lock ₂ 10 Tf 50
15	Probiotics lower plasma glucose in the high-fat fed C57BL/6J mouse. Beneficial Microbes, 2010, 1, 189-196.	1.0	92
16	Short-chain fatty acid formation in the hindgut of rats fed oligosaccharides varying in monomeric composition, degree of polymerisation and solubility. British Journal of Nutrition, 2005, 94, 705-713.	1.2	81
17	Lingonberries reduce atherosclerosis in <i>Apoe^{â€∤â€}</i> mice in association with altered gut microbiota composition and improved lipid profile. Molecular Nutrition and Food Research, 2016, 60, 1150-1160.	1.5	67
18	Minerals, phytate and dietary fibre in different fractions of oat-grain. Journal of Cereal Science, 1988, 7, 73-82.	1.8	62

#	Article	IF	CITATIONS
19	On the possibility of using high pressure treatment to modify physico-chemical properties of dietary fibre in white cabbage (Brassica oleracea var. capitata). Innovative Food Science and Emerging Technologies, 2004, 5, 171-177.	2.7	55
20	Carbohydrate composition and content of organic acids in fresh and stored apples. Journal of the Science of Food and Agriculture, 2000, 80, 1538-1544.	1.7	54
21	Dietary fibre fermentation in the rat intestinal tract: effect of adaptation period, protein and fibre levels, and particle size. British Journal of Nutrition, 1985, 54, 635-643.	1.2	53
22	Xylooligosaccharides Increase <i>Bifidobacteria</i> and <i>Lachnospiraceae</i> in Mice on a High-Fat Diet, with a Concomitant Increase in Short-Chain Fatty Acids, Especially Butyric Acid. Journal of Agricultural and Food Chemistry, 2021, 69, 3617-3625.	2.4	48
23	Addition of Various Carbohydrates to Beef Burgers Affects the Formation of Heterocyclic Amines during Frying. Journal of Agricultural and Food Chemistry, 2004, 52, 7561-7566.	2.4	47
24	Dietary supplementation with \hat{l}^2 -glucan enriched oat bran increases faecal concentration of carboxylic acids in healthy subjects. European Journal of Clinical Nutrition, 2008, 62, 978-984.	1.3	46
25	Designing future prebiotic fiber to target metabolic syndrome. Nutrition, 2014, 30, 497-502.	1.1	46
26	Binding of mineral elements by dietary fibre components in cerealsâ€"In vitro (III). Food Chemistry, 1991, 40, 169-183.	4.2	45
27	Binding of mineral elements by some dietary fibre components—In vitro (II). Food Chemistry, 1987, 26, 139-148.	4.2	43
28	Fermentation of Vegetable Fiber in the Intestinal Tract of Rats and Effects on Fecal Bulking and Bile Acid Excretion. Journal of Nutrition, 1990, 120, 459-466.	1.3	43
29	Serum Short-Chain Fatty Acids and Associations With Inflammation in Newly Diagnosed Patients With Multiple Sclerosis and Healthy Controls. Frontiers in Immunology, 2021, 12, 661493.	2.2	43
30	Rheological and Chemical Properties of Mucilage in Different Varieties from Linseed (<i>Linum) Tj ETQq0 0 0 rgBT</i>	/Overlock	10 Tf 50 30
31	Fasting serum concentration of short-chain fatty acids in subjects with microscopic colitis and celiac disease: no difference compared with controls, but between genders. Scandinavian Journal of Gastroenterology, 2013, 48, 696-701.	0.6	40
32	Formation of Short-Chain Fatty Acids, Excretion of Anthocyanins, and Microbial Diversity in Rats Fed Blackcurrants, Blackberries, and Raspberries. Journal of Nutrition and Metabolism, 2013, 2013, 1-12.	0.7	39
33	Determination of digestible energy values and fermentabilities of dietary fibre supplements: a European interlaboratory study in vivo. British Journal of Nutrition, 1995, 74, 289-302.	1.2	38
34	Physiological effects of cereal dietary fibre. Carbohydrate Polymers, 1993, 21, 183-187.	5.1	37
35	Influence of processing and cooking of carrots in mixed meals on satiety, glucose and hormonal response. International Journal of Food Sciences and Nutrition, 1995, 46, 3-12.	1.3	35
36	Blueberry Husks and Probiotics Attenuate Colorectal Inflammation and Oncogenesis, and Liver Injuries in Rats Exposed to Cycling DSS-Treatment. PLoS ONE, 2012, 7, e33510.	1.1	34

#	Article	IF	CITATIONS
37	Enzyme resistant starch fractions and dietary fibre. Scandinavian Journal of Gastroenterology, 1987, 22, 29-32.	0.6	33
38	Blueberry husks and multi-strain probiotics affect colonic fermentation in rats. British Journal of Nutrition, 2009, 101, 859-870.	1.2	31
39	Short-chain fatty acid formation at fermentation of indigestible carbohydrates. NĀĦngsforskning: Referattidskrift l NĀĦngsforskningsfrÃ¥gor, 2001, 45, 165-168.	0.0	30
40	Molar mass and rheological characterisation of an exopolysaccharide from Pediococcus damnosus 2.6. Carbohydrate Polymers, 2007, 68, 577-586.	5.1	30
41	Effects of Soluble and Insoluble Fractions from Bilberries, Black Currants, and Raspberries on Short-Chain Fatty Acid Formation, Anthocyanin Excretion, and Cholesterol in Rats. Journal of Agricultural and Food Chemistry, 2014, 62, 4359-4368.	2.4	30
42	Binding of mineral elements by some dietary fibre components—in vitro (I). Food Chemistry, 1987, 23, 295-303.	4.2	28
43	Wheat bran increases high-density-lipoprotein cholesterol in the rat. British Journal of Nutrition, 1981, 46, 385-393.	1.2	27
44	Satiety effects of spinach in mixed meals: Comparison with other vegetables. International Journal of Food Sciences and Nutrition, 1995, 46, 327-334.	1.3	25
45	Extraction of βâ€Glucan from Oat Bran in Laboratory Scale. Cereal Chemistry, 2009, 86, 601-608.	1.1	25
46	Lingonberries and their two separated fractions differently alter the gut microbiota, improve metabolic functions, reduce gut inflammatory properties, and improve brain function in ApoEâ^'/â^' mice fed high-fat diet. Nutritional Neuroscience, 2020, 23, 600-612.	1.5	25
47	Processing effects on the nutritional advancement of probiotics and prebiotics. Microbial Ecology in Health and Disease, 2004, 16 , 113 - 124 .	3.8	24
48	Determination of free and conjugated bile acids in serum of Apoe(\hat{a}^2/\hat{a}^2) mice fed different lingonberry fractions by UHPLC-MS. Scientific Reports, 2019, 9, 3800.	1.6	24
49	Bulk Laxatives: Their Dietary Fibre Composition, Degradation, and Faecal Bulking Capacity in the Rat. Scandinavian Journal of Gastroenterology, 1985, 20, 887-895.	0.6	22
50	Optimization of extraction methods for determination of the raffinose family oligosaccharides in leguminous vine peas (Pisum sativum L.) and effects of blanching. Journal of Food Composition and Analysis, 2007, 20, 13-18.	1.9	22
51	Fermentation of dietary fibre in the intestinal tract of rats â€" a comparison of flours with different extraction rates from six cereals. Journal of Cereal Science, 1985, 3, 207-219.	1.8	21
52	Blueberry husks, rye bran and multi-strain probiotics affect the severity of colitis induced by dextran sulphate sodium. Scandinavian Journal of Gastroenterology, 2009, 44, 1213-1225.	0.6	21
53	Barley malt increases hindgut and portal butyric acid, modulates gene expression of gut tight junction proteins and Toll-like receptors in rats fed high-fat diets, but high advanced glycation end-products partially attenuate the effects. Food and Function, 2015, 6, 3165-3176.	2.1	21
54	Monobutyrin Reduces Liver Cholesterol and Improves Intestinal Barrier Function in Rats Fed High-Fat Diets. Nutrients, 2019, 11, 308.	1.7	21

#	Article	IF	CITATIONS
55	Popping of whole-grain wheat: Effects on dietary fibre degradation in the rat intestinal tract. Journal of Cereal Science, 1987, 5, 67-72.	1.8	20
56	Content of low molecular weight carbohydrates in vining peas (Pisum sativum) related to harvest time, size and brine grade. Food Chemistry, 2006, 94, 513-519.	4.2	19
57	Monovalerin and trivalerin increase brain acetic acid, decrease liver succinic acid, and alter gut microbiota in rats fed high-fat diets. European Journal of Nutrition, 2019, 58, 1545-1560.	1.8	18
58	Degradation of water-soluble fibre polysaccharides in carrots after different types of processing. Food Chemistry, 1993, 47, 169-176.	4.2	17
59	Effects of harvest time and storage on dietary fibre components in various cultivars of white cabbage (Brassica oleracea varcapitata). Journal of the Science of Food and Agriculture, 2002, 82, 1405-1411.	1.7	17
60	Dietary Fiber in Bilberry Ameliorates Pre-Obesity Events in Rats by Regulating Lipid Depot, Cecal Short-Chain Fatty Acid Formation and Microbiota Composition. Nutrients, 2019, 11, 1350.	1.7	17
61	Dietary Oat Bran Reduces Systemic Inflammation in Mice Subjected to Pelvic Irradiation. Nutrients, 2020, 12, 2172.	1.7	16
62	Impact of dietary induced precocious gut maturation on cecal microbiota and its relation to the bloodâ€brain barrier during the postnatal period in rats. Neurogastroenterology and Motility, 2018, 30, e13285.	1.6	15
63	Highâ€Fat Diet Enriched with Bilberry Modifies Colonic Mucus Dynamics and Restores Marked Alterations of Gut Microbiome in Rats. Molecular Nutrition and Food Research, 2019, 63, e1900117.	1.5	14
64	Bifidobacterium lactis Bb-12 and Lactobacillus salivarius UCC500 Modify Carboxylic Acid Formation in the Hindgut of Rats Given Pectin, Inulin, and Lactitol. Journal of Nutrition, 2006, 136, 2175-2180.	1.3	13
65	Determination of bile acids by hollow fibre liquid-phase microextraction coupled with gas chromatography. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2014, 944, 69-74.	1.2	12
66	Monobutyrin and Monovalerin Affect Brain Short-Chain Fatty Acid Profiles and Tight-Junction Protein Expression in ApoE-Knockout Rats Fed High-Fat Diets. Nutrients, 2020, 12, 1202.	1.7	12
67	Carboxylic acids in the hindgut of rats fed highly soluble inulin andBifidobacterium lactis(Bb-12),Lactobacillus salivarius(UCC500) orLactobacillus rhamnosus(GG). Food Nutrition Research, 2007, 51, 13-21.	0.3	10
68	The Effect of Dietary Fiber from Wheat Processing Streams on the Formation of Carboxylic Acids and Microbiota in the Hindgut of Rats. Journal of Agricultural and Food Chemistry, 2011, 59, 3406-3413.	2.4	10
69	Oat Bran Increased Fecal Butyrate and Prevented Gastrointestinal Symptoms in Patients With Quiescent Ulcerative Colitis—Randomized Controlled Trial. Crohn's & Colitis 360, 2020, 2, .	0.5	9
70	Novel xylan degrading enzymes from polysaccharide utilizing loci of <i>Prevotella copri</i> DSM18205. Glycobiology, 2021, 31, 1330-1349.	1.3	9
71	Dietary Fibre in Type II Diabetes. Acta Medica Scandinavica, 1982, 210, 47-50.	0.0	8
72	Effects of Whole Brown Bean and Its Isolated Fiber Fraction on Plasma Lipid Profile, Atherosclerosis, Gut Microbiota, and Microbiota-Dependent Metabolites in Apoeâ ² /â ² Mice. Nutrients, 2022, 14, 937.	1.7	8

#	Article	IF	CITATIONS
73	Content of low molecular weight carbohydrates in vining peas (Pisum sativum) after blanching and freezing: effect of cultivar and cultivation conditions. Journal of the Science of Food and Agriculture, 2005, 85, 691-699.	1.7	7
74	Characterization of Indigestible Carbohydrates in Various Fractions from Wheat Processing. Cereal Chemistry, 2010, 87, 125-130.	1.1	6
75	Application of a dynamic gastrointestinal in vitro model combined with a rat model to predict the digestive fate of barley dietary fibre and evaluate potential impact on hindgut fermentation. Bioactive Carbohydrates and Dietary Fibre, 2017, 9, 7-13.	1.5	6
76	Intestinal degradation of dietary fibre in green beans-effects of microwave treatments. International Journal of Food Sciences and Nutrition, 1999, 50, 245-253.	1.3	5
77	Colorectal Oncogenesis and Inflammation in a Rat Model Based on Chronic Inflammation due to Cycling DSS Treatments. Gastroenterology Research and Practice, 2011, 2011, 1-15.	0.7	5
78	Prebiotic and synbiotic effects on rats fed malted barley with selected bacteria strains. Food and Nutrition Research, 2014, 58, 24848.	1.2	5
79	Vegetable, fruit and potato fibres. , 2013, , 193-207.		4
80	A Fiber-Rich Diet and Radiation-Induced Injury in the Murine Intestinal Mucosa. International Journal of Molecular Sciences, 2022, 23, 439.	1.8	4
81	Chemistry and fermentative breakdown of dietary fibre in bulk-laxatives compared to some fibre containing foods. Scandinavian Journal of Gastroenterology, 1987, 22, 52-54.	0.6	2
82	The satiety, glucose, and hormonal responses after mixed meals with vegetables. American Journal of Clinical Nutrition, 1994, 59, 793S.	2.2	2
83	Irradiation Induces Tuft Cell Hyperplasia and Myenteric Neuronal Loss in the Absence of Dietary Fiber in a Mouse Model of Pelvic Radiotherapy. Gastroenterology Insights, 2022, 13, 87-102.	0.7	2
84	Dietary Fiber and the Hippocampal Neurogenic Niche in a Model of Pelvic Radiotherapy. Neuroscience, 2021, 475, 137-147.	1.1	1
85	In memoriam of Nils-Georg Asp. Food and Nutrition Research, 2014, 58, 24317.	1.2	0