

Yuki Fujiwara

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6607240/publications.pdf>

Version: 2024-02-01

50
papers

1,199
citations

361045

20
h-index

414034

32
g-index

50
all docs

50
docs citations

50
times ranked

1829
citing authors

#	ARTICLE	IF	CITATIONS
1	The C-reactive Protein to Albumin Ratio Predicts Long-Term Outcomes in Patients with Pancreatic Cancer After Pancreatic Resection. <i>World Journal of Surgery</i> , 2016, 40, 2254-2260.	0.8	107
2	Pancreatic cancer treatment: better, but a long way to go. <i>Surgery Today</i> , 2020, 50, 1117-1125.	0.7	91
3	Negative Impact of Blood Transfusion on Recurrence and Prognosis of Hepatocellular Carcinoma After Hepatic Resection. <i>Journal of Gastrointestinal Surgery</i> , 2009, 13, 1636-1642.	0.9	86
4	Preoperative platelet to lymphocyte ratio predicts outcome of patients with pancreatic ductal adenocarcinoma after pancreatic resection. <i>Surgery</i> , 2015, 158, 360-365.	1.0	77
5	Blockade of the CD93 pathway normalizes tumor vasculature to facilitate drug delivery and immunotherapy. <i>Science Translational Medicine</i> , 2021, 13, .	5.8	54
6	Comprehensive assessment of the prognosis of pancreatic cancer: peripheral blood neutrophil-lymphocyte ratio and immunohistochemical analyses of the tumour site. <i>Scandinavian Journal of Gastroenterology</i> , 2016, 51, 610-617.	0.6	49
7	Preoperative Platelet-to-Albumin Ratio Predicts Prognosis of Patients with Pancreatic Ductal Adenocarcinoma After Pancreatic Resection. <i>Anticancer Research</i> , 2017, 37, 787-794.	0.5	46
8	Nafamostat mesilate can prevent adhesion, invasion and peritoneal dissemination of pancreatic cancer thorough nuclear factor kappa-B inhibition. <i>Journal of Hepato-Biliary-Pancreatic Sciences</i> , 2011, 18, 731-739.	1.4	40
9	Impact of the C-reactive protein to albumin ratio on long-term outcomes after hepatic resection for colorectal liver metastases. <i>American Journal of Surgery</i> , 2017, 214, 752-756.	0.9	36
10	Glasgow Prognostic Score Predicts Outcome After Surgical Resection of Gallbladder Cancer. <i>World Journal of Surgery</i> , 2015, 39, 753-758.	0.8	32
11	Perioperative change in peripheral blood monocyte count may predict prognosis in patients with colorectal liver metastasis after hepatic resection. <i>Journal of Surgical Oncology</i> , 2012, 106, 31-35.	0.8	29
12	Anti-tumor effect by inhibition of NF- κ B activation using nafamostat mesilate for pancreatic cancer in a mouse model. <i>Oncology Reports</i> , 2010, 24, 843-50.	1.2	29
13	Glasgow prognostic score is related to blood transfusion requirements and post-operative complications in hepatic resection for hepatocellular carcinoma. <i>Anticancer Research</i> , 2010, 30, 5129-36.	0.5	27
14	Combination chemotherapy of serine protease inhibitor nafamostat mesilate with oxaliplatin targeting NF- κ B activation for pancreatic cancer. <i>Cancer Letters</i> , 2013, 333, 89-95.	3.2	26
15	The Glasgow prognostic score is valuable for colorectal cancer with both synchronous and metachronous unresectable liver metastases. <i>Oncology Letters</i> , 2012, 4, 324-328.	0.8	25
16	Risk stratification using a novel liver functional reserve score of combination prothrombin time-international normalized ratio to albumin ratio and albumin in patients with hepatocellular carcinoma. <i>Surgery</i> , 2018, 164, 404-410.	1.0	24
17	Postoperative peak serum C-reactive protein predicts outcome of hepatic resection for hepatocellular carcinoma. <i>Anticancer Research</i> , 2013, 33, 705-9.	0.5	22
18	Combination chemotherapy of nafamostat mesylate with gemcitabine for gallbladder cancer targeting nuclear factor- κ B activation. <i>Journal of Surgical Research</i> , 2013, 184, 605-612.	0.8	21

#	ARTICLE	IF	CITATIONS
19	Perioperative serum albumin correlates with postoperative pancreatic fistula after pancreaticoduodenectomy. <i>Anticancer Research</i> , 2015, 35, 499-503.	0.5	21
20	Inhibition of Nuclear Factor- κ B Enhances the Antitumor Effect of Paclitaxel Against Gastric Cancer with Peritoneal Dissemination in Mice. <i>Digestive Diseases and Sciences</i> , 2013, 58, 123-131.	1.1	20
21	Dual inhibition of nuclear factor kappa-B and Mdm2 enhance the antitumor effect of radiation therapy for pancreatic cancer. <i>Cancer Letters</i> , 2016, 370, 177-184.	3.2	20
22	Pomalidomide Inhibits PD-L1 Induction to Promote Antitumor Immunity. <i>Cancer Research</i> , 2018, 78, 6655-6665.	0.4	20
23	Postoperative peripheral absolute blood lymphocyte-to-monocyte ratio predicts therapeutic outcome after pancreatic resection in patients with pancreatic adenocarcinoma. <i>Anticancer Research</i> , 2014, 34, 5163-8.	0.5	19
24	Negative impact of surgical site infection on long-term outcomes after hepatic resection for colorectal liver metastases. <i>Anticancer Research</i> , 2013, 33, 1697-703.	0.5	18
25	The Prognostic Significance of C-reactive Protein-To-Lymphocyte Ratio in Colorectal Liver Metastases. <i>Journal of Surgical Research</i> , 2021, 258, 414-421.	0.8	17
26	Neutrophil to Lymphocyte Ratio Predicts Therapeutic Outcome After Pancreaticoduodenectomy for Carcinoma of the Ampulla of Vater. <i>Anticancer Research</i> , 2016, 36, 403-8.	0.5	17
27	Inhibition of Nuclear Factor Kappa-B Enhances the Antitumor Effect of Combination Treatment with Tumor Necrosis Factor-Alpha Gene Therapy and Gemcitabine for Pancreatic Cancer in Mice. <i>Journal of the American College of Surgeons</i> , 2013, 216, 320-332e3.	0.2	16
28	Glasgow prognostic score predicts therapeutic outcome after hepatic resection for hepatocellular carcinoma. <i>Oncology Letters</i> , 2017, 14, 293-298.	0.8	16
29	C-Reactive Protein-based Prognostic Measures Are Superior at Predicting Survival Compared with Peripheral Blood Cell Count-based Ones in Patients After Curative Resection for Pancreatic Cancer. <i>Anticancer Research</i> , 2018, 38, 6491-6499.	0.5	15
30	GPR182 limits antitumor immunity via chemokine scavenging in mouse melanoma models. <i>Nature Communications</i> , 2022, 13, 97.	5.8	15
31	A novel postoperative inflammatory score predicts postoperative pancreatic fistula after pancreatic resection. <i>Anticancer Research</i> , 2013, 33, 5005-10.	0.5	14
32	The Comparison of Inflammation-Based Prognostic Scores in Patients With Extrahepatic Bile Duct Cancer After Pancreaticoduodenectomy. <i>Journal of Surgical Research</i> , 2019, 238, 102-112.	0.8	12
33	CD28H expression identifies resident memory CD8 ⁺ T cells with less cytotoxicity in human peripheral tissues and cancers. <i>Oncolmmunology</i> , 2019, 8, e1538440.	2.1	12
34	Glasgow prognostic score predicts therapeutic outcome after pancreaticoduodenectomy for carcinoma of the ampulla of vater. <i>Anticancer Research</i> , 2013, 33, 2715-21.	0.5	12
35	Combination treatment using adenovirus vector-mediated tumor necrosis factor-alpha gene transfer and a NF- κ B inhibitor for pancreatic cancer in mice. <i>Cancer Letters</i> , 2011, 306, 92-98.	3.2	11
36	Combination Paclitaxel and Inhibitor of Nuclear Factor κ B Activation Improves Therapeutic Outcome for Model Mice With Peritoneal Dissemination of Pancreatic Cancer. <i>Pancreas</i> , 2011, 40, 600-607.	0.5	11

#	ARTICLE	IF	CITATIONS
37	The GPR171 pathway suppresses T cell activation and limits antitumor immunity. <i>Nature Communications</i> , 2021, 12, 5857.	5.8	11
38	Negative Impact of Preoperative Endoscopic Biliary Drainage on Prognosis of Pancreatic Ductal Adenocarcinoma After Pancreaticoduodenectomy. <i>Anticancer Research</i> , 2015, 35, 5079-83.	0.5	11
39	Nuclear factor κ B activity correlates with the progression and prognosis of pancreatic cancer in a mouse model. <i>Surgery Today</i> , 2013, 43, 171-177.	0.7	10
40	Preoperative controlling nutritional status (CONUT) score predicts long-term outcomes in patients with non-B non-C hepatocellular carcinoma after curative hepatic resection. <i>Langenbeck's Archives of Surgery</i> , 2021, 406, 99-107.	0.8	10
41	Effectiveness of Anatomical Resection for Small Hepatocellular Carcinoma: a Propensity Score-Matched Analysis of a Multi-institutional Database. <i>Journal of Gastrointestinal Surgery</i> , 2021, 25, 2835-2841.	0.9	10
42	Perioperative change in white blood cell count predicts outcome of hepatic resection for hepatocellular carcinoma. <i>Journal of Hepato-Biliary-Pancreatic Sciences</i> , 2010, 17, 892-897.	1.4	8
43	Preoperative peripheral blood neutrophil count predicts long-term outcomes following hepatic resection for colorectal liver metastases. <i>Oncology Letters</i> , 2017, 13, 3688-3694.	0.8	7
44	Combination of Distance from Superior Mesenteric Artery and Serum CA19-9 as a Novel Prediction of Local Recurrence in Patients With Pancreatic Cancer Following Resection. <i>Anticancer Research</i> , 2019, 39, 1469-1478.	0.5	5
45	A Novel Prognostic Score Combining Preoperative Biliary Drainage and Inflammatory Status for Patients with Periampullary Cancers. <i>Journal of Gastrointestinal Surgery</i> , 2020, 24, 813-822.	0.9	5
46	Radical resection of T1 pancreatic adenocarcinoma with a pseudocyst of the tail due to acute obstructive pancreatitis: report of a case. <i>Surgical Case Reports</i> , 2016, 2, 144.	0.2	4
47	Hepatic abscess in the Spiegel lobe caused by foreign body penetration: report of a case report. <i>Surgical Case Reports</i> , 2017, 3, 24.	0.2	4
48	Glycogen synthase kinase-3 β activity plays a key role in the antitumor effect of nafamostat mesilate in pancreatic cancer cells. <i>Annals of Gastroenterological Surgery</i> , 2018, 2, 65-71.	1.2	3
49	Radical Resection of a Primarily Unresectable Pancreatic Cancer After Neoadjuvant Chemotherapy Using Gemcitabine, TS-1, and Nafamostat Mesilate; Report of a Case. <i>International Surgery</i> , 2015, 100, 287-291.	0.0	2
50	Hepatic Thermal Injury Promotes Colorectal Cancer Engraftment in C57/black 6 Mice. <i>American Journal of Physiology - Cell Physiology</i> , 2020, 320, C142-C151.	2.1	2