

# Ely Kosnicki

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6623870/publications.pdf>

Version: 2024-02-01

16  
papers

102  
citations

1307594

7  
h-index

1372567

10  
g-index

16  
all docs

16  
docs citations

16  
times ranked

119  
citing authors

#	ARTICLE	IF	CITATIONS
1	OUP accepted manuscript. Journal of Insect Science, 2022, 22, .	1.5	0
2	Determining Sexual Dimorphism of Living Aquatic Beetles, <i>Stygoparnus comalensis</i> (Coleoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 7 Journal of Insect Science, 2019, 19, .	1.5	1
3	Variation in the number of lateral telson spines of <i>Stygobromus flagellatus</i> (Benedict, 1896) (Amphipoda: Crangonyctidae), a subterranean species from Texas, USA. Journal of Crustacean Biology, 2019, 39, 186-188.	0.8	1
4	Hydrogeomorphic Reference Condition and Its Relationship with Macroinvertebrate Assemblages in Southeastern U.S. Sand Hills Streams. Journal of the American Water Resources Association, 2018, 54, 914-933.	2.4	0
5	Development of an Index of Biotic Integrity for the Sand Hills Ecoregion of the Southeastern United States. Transactions of the American Fisheries Society, 2017, 146, 112-127.	1.4	8
6	Factors Influencing Stream Fish Species Composition and Functional Properties at Multiple Spatial Scales in the Sand Hills of the Southeastern United States. Transactions of the American Fisheries Society, 2016, 145, 545-562.	1.4	8
7	A Stream Multimetric Macroinvertebrate Index (MMI) for the Sand Hills Ecoregion of the Southeastern Plains, USA. Environmental Management, 2016, 58, 741-751.	2.7	7
8	Empirical Estimation of Stream Discharge Using Channel Geometry in Low Gradient, Sand Bed Streams of the Southeastern Plains. Journal of the American Water Resources Association, 2015, 51, 1060-1071.	2.4	8
9	Defining the Reference Condition for Wadeable Streams in the Sand Hills Subdivision of the Southeastern Plains Ecoregion, USA. Environmental Management, 2014, 54, 494-504.	2.7	11
10	The influence of soil type, congeneric cues, and floodplain connectivity on the local distribution of the devil crayfish ( <i>Cambarus diogenes</i> Girard). Freshwater Science, 2013, 32, 1333-1344.	1.8	24
11	Seasonal predictability of benthic macroinvertebrate metrics and community structure with maturity-weighted abundances in a Missouri Ozark stream, USA. Ecological Indicators, 2011, 11, 704-714.	6.3	8
12	First Nearctic Records of <i>Oliveiriella</i> Wiedenbrug and Fittkau, with New Distributional Records for Two Other New World Species of Orthoclaadiinae (Diptera: Chironomidae). Entomological News, 2009, 120, 349-362.	0.2	3
13	Least-Desired Index for Assessing the Effectiveness of Grass Riparian Filter Strips in Improving Water Quality in an Agricultural Region. Environmental Entomology, 2007, 36, 713-724.	1.4	7
14	Least-Desired Index for Assessing the Effectiveness of Grass Riparian Filter Strips in Improving Water Quality in an Agricultural Region. Environmental Entomology, 2007, 36, 713-724.	1.4	6
15	Dehydration Tolerance of the Temporary Pond Mayfly <i>Siphonurus typicus</i> (Ephemeroptera: Tj ETQq1 1 0.784314 rgBT /Overlock 1.2 0	1.2	0
16	Life history aspects of the mayfly <i>Siphonurus typicus</i> (Ephemeroptera: Siphonuridae) with a new application for measuring nymphal development and growth. Hydrobiologia, 2003, 510, 131-146.	2.0	10