
J Herrstedt

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6627781/publications.pdf Version: 2024-02-01



I HEDDSTEDT

#	Article	IF	CITATIONS
1	2016 MASCC and ESMO guideline update for the prevention of chemotherapy- and radiotherapy-induced nausea and vomiting and of nausea and vomiting in advanced cancer patients. Annals of Oncology, 2016, 27, v119-v133.	0.6	454
2	Management of febrile neutropaenia: ESMO Clinical Practice Guidelines. Annals of Oncology, 2016, 27, v111-v118.	0.6	454
3	Bone health in cancer patients: ESMO Clinical Practice Guidelines. Annals of Oncology, 2014, 25, iii124-iii137.	0.6	450
4	Early recognition of malnutrition and cachexia in the cancer patient: a position paper of a European School of Oncology Task Force. Annals of Oncology, 2014, 25, 1492-1499.	0.6	240
5	Comorbidity in elderly cancer patients in relation to overall and cancer-specific mortality. British Journal of Cancer, 2012, 106, 1353-1360.	2.9	222
6	Management of oral and gastrointestinal mucosal injury: ESMO Clinical Practice Guidelines for diagnosis, treatment, and follow-up. Annals of Oncology, 2015, 26, v139-v151.	0.6	188
7	Ondansetron plus Metopimazine Compared with Ondansetron Alone in Patients Receiving Moderately Emetogenic Chemotherapy. New England Journal of Medicine, 1993, 328, 1076-1080.	13.9	84
8	Polypharmacy and drug use in elderly Danish cancer patients during 1996 to 2006. Journal of Geriatric Oncology, 2012, 3, 33-40.	0.5	58
9	Ondansetron Plus Metopimazine Compared With Ondansetron Plus Metopimazine Plus Prednisolone as Antiemetic Prophylaxis in Patients Receiving Multiple Cycles of Moderately Emetogenic Chemotherapy. Journal of Clinical Oncology, 2001, 19, 2091-2097.	0.8	42
10	Chemotherapy-induced nausea and vomiting: ESMO Clinical Recommendations for prophylaxis. Annals of Oncology, 2009, 20, iv156-iv158.	0.6	34
11	Randomized, double-blind comparison of ondansetron versus ondansetron plus metopimazine as antiemetic prophylaxis during platinum-based chemotherapy in patients with cancer Journal of Clinical Oncology, 1997, 15, 1690-1696.	0.8	30
12	Long-term safety in patients with recurrent ovarian cancer treated with niraparib versus placebo: Results from the phase III ENGOT-OV16/NOVA trial. Gynecologic Oncology, 2020, 159, 442-448.	0.6	28
13	Non-intercepted dose errors in prescribing anti-neoplastic treatment: a prospective, comparative cohort study. Annals of Oncology, 2015, 26, 981-986.	0.6	23
14	The impact of comprehensive geriatric assessment for optimal treatment of older patients with cancer: A randomized parallel-group clinical trial. Journal of Geriatric Oncology, 2020, 11, 488-495.	0.5	22
15	Interaction of the antiemetic metopimazine and anticancer agents with brain dopamine D2, 5-hydroxytryptamine3, histamine H1, muscarine cholinergic and ?1 receptors. Cancer Chemotherapy and Pharmacology, 1993, 33, 53-56.	1.1	21
16	MASCC/ESMO Antiemetic Guidelines: Introduction to the 2016 guideline update. Supportive Care in Cancer, 2017, 25, 267-269.	1.0	19
17	Randomized, double-blind trial comparing the antiemetic effect of tropisetron plus metopimazine with tropisetron plus placebo in patients receiving multiple cycles of multiple-day cisplatin-based chemotherapy. Supportive Care in Cancer, 2007, 15, 417-426.	1.0	18
18	Prognostic factors in young ovarian cancer patients: An analysis of four prospective phase III intergroup trials of the AGO Study Group, GINECO and NSGO. European Journal of Cancer, 2016, 66, 114-124.	1.3	16

J HERRSTEDT

#	Article	IF	CITATIONS
19	Organized breast cancer screening not only reduces mortality from breast cancer but also significantly decreases disability-adjusted life years: analysis of the Global Burden of Disease Study and screening programme availability in 130 countries. ESMO Open, 2021, 6, 100111.	2.0	13
20	The effect of food on serum concentrations of metopimazine British Journal of Clinical Pharmacology, 1990, 30, 237-243.	1.1	12
21	Increased myelosuppression during cytostatic treatment and pleural effusion in patients with small cell lung cancer. European Journal of Cancer, 1992, 28, 1070-1073.	1.3	12
22	Dose-finding study of oral metopimazine. Supportive Care in Cancer, 1997, 5, 38-43.	1.0	12
23	Amisulpride prevents nausea and vomiting associated with highly emetogenic chemotherapy: a randomised, double-blind, placebo-controlled, dose-ranging trial. Supportive Care in Cancer, 2019, 27, 2699-2705.	1.0	11
24	Early tumor regrowth is a contributor to impaired survival in patients with completely resected advanced ovarian cancer. An exploratory analysis of the Intergroup trial AGO-OVAR 12. Gynecologic Oncology, 2019, 152, 235-242.	0.6	10
25	Prevention of Chemotherapy-Induced Nausea and Vomiting in the Older Patient: Optimizing Outcomes. Drugs and Aging, 2022, 39, 1-21.	1.3	10
26	Development of Antiemetic Therapy in Cancer Patients. Acta OncolÃ ³ gica, 1995, 34, 637-640.	0.8	9
27	High-dose metoclopramide + lorazepam versus low-dose metoclopramide + lorazepam + dehydrobenzperidol in the treatment of cisplatin-induced nausea and vomiting. Annals of Oncology, 1991, 2, 223-227.	0.6	8
28	Bioavailability of the antiemetic metopimazine given as a microenema. British Journal of Clinical Pharmacology, 1996, 41, 613-615.	1.1	8
29	Impact of Age, Comorbidity, and FIGO Stage on Treatment Choice and Mortality in Older Danish Patients with Gynecological Cancer: A Retrospective Register-Based Cohort Study. Drugs - Real World Outcomes, 2018, 5, 225-235.	0.7	8
30	Integrated safety analysis of rolapitant with coadministered drugs from phase II/III trials: an assessment of CYP2D6 or BCRP inhibition by rolapitant. Annals of Oncology, 2017, 28, 1268-1273.	0.6	7
31	We still need common criteria for the assessment of nausea and vomiting. European Journal of Cancer, 1994, 30, 1217.	1.3	2
32	Addressing Chemotherapy-Induced Peripheral Neuropathy Using Multi-Frequency Vibrometry and Patient-Reported Outcomes. Journal of Clinical Medicine, 2022, 11, 1862.	1.0	2
33	Effect and Tolerability of Immunotherapy in Patients with NSCLC with or without Brain Metastasis. Cancers, 2022, 14, 1682.	1.7	2
34	The prevalence, distribution and impact of peripheral neuropathy among Danish patients with cancer – a population-based cross-sectional study. Acta Oncológica, 2022, 61, 363-370.	0.8	1
35	A randomized, phase III study (AGO-OVAR-9, GINECO-TCG, NSGO-OC-0102): Gemcitabine-paclitaxel-carboplatin (TCG) versus paclitaxel-carboplatin (TC) as first-line treatment of ovarian cancer (OC): Survival of FIGO stage I-IIA patients. Journal of Clinical Oncology, 2009, 27, LBA5510-LBA5510.	0.8	0