
Dante Zarlenga

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6628032/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Horizontal gene transfer provides insights into the deep evolutionary history and biology of Trichinella. Food and Waterborne Parasitology, 2022, 27, e00155.	2.7	Ο
2	Ostertagia ostertagi Mediates Early Host Immune Responses via Macrophage and Toll-Like Receptor Pathways. Infection and Immunity, 2021, 89, .	2.2	2
3	Repeated, drug-truncated infections with Ostertagia ostertagi elicit strong humoral and cell-mediated immune responses and confer partial protection in cattle. Veterinary Parasitology, 2021, 296, 109510.	1.8	2
4	A simple molecular method to identify and quantify genera of gastrointestinal nematodes of cattle. Parasitology Research, 2021, 120, 3979-3986.	1.6	1
5	Immune reactivity and host modulatory roles of two novel Haemonchus contortus cathepsin B-like proteases. Parasites and Vectors, 2021, 14, 580.	2.5	2
6	A new paraprobiotic-based treatment for control of Haemonchus contortus in sheep. International Journal for Parasitology: Drugs and Drug Resistance, 2020, 14, 230-236.	3.4	16
7	Trichinella species and genotypes. Research in Veterinary Science, 2020, 133, 289-296.	1.9	48
8	Preface: International Commission on Trichinellosis recommendations for the detection and control of Trichinella. Food and Waterborne Parasitology, 2019, 17, e00063.	2.7	0
9	International Commission on Trichinellosis: Recommendations for genotyping Trichinella muscle stage larvae. Food and Waterborne Parasitology, 2019, 15, e00033.	2.7	14
10	Characterization of IL-10-producing neutrophils in cattle infected with Ostertagia ostertagi. Scientific Reports, 2019, 9, 20292.	3.3	12
11	A tale of three kingdoms: members of the Phylum Nematoda independently acquired the detoxifying enzyme cyanase through horizontal gene transfer from plants and bacteria. Parasitology, 2019, 146, 445-452.	1.5	9
12	Trichinella spiralis : Adaptation and parasitism. Veterinary Parasitology, 2016, 231, 8-21.	1.8	14
13	Abomasal mucosal immune responses of cattle with limited or continuous exposure to pasture-borne gastrointestinal nematode parasite infection. Veterinary Parasitology, 2016, 229, 118-125.	1.8	8
14	Hybridization is limited between two lineages of freeze-resistant Trichinella during coinfection in a mouse model. Infection, Genetics and Evolution, 2016, 38, 146-151.	2.3	6
15	Phage display for identifying peptides that bind the spike protein of transmissible gastroenteritis virus and possess diagnostic potential. Virus Genes, 2015, 51, 51-56.	1.6	10
16	Ostertagia ostertagi macrophage migration inhibitory factor is present in all developmental stages and may cross-regulate host functions through interaction with the host receptor. International Journal for Parasitology, 2014, 44, 355-367.	3.1	20
17	TsDAF-21/Hsp90 is expressed in all examined stages of Trichinella spiralis. Veterinary Parasitology, 2013, 194, 171-174.	1.8	7
18	Exploring metazoan evolution through dynamic and holistic changes in protein families and domains. BMC Evolutionary Biology, 2012, 12, 138.	3.2	9

DANTE ZARLENGA

#	Article	IF	CITATIONS
19	A calcium-activated nucleotidase secreted from <i>Ostertagia ostertagi</i> 4th-stage larvae is a member of the novel salivary apyrases present in blood-feeding arthropods. Parasitology, 2011, 138, 333-343.	1.5	14
20	Bacterial expression of antigenic sites A and D in the spike protein of transmissible gastroenteritis virus and evaluation of their inhibitory effects on viral infection. Virus Genes, 2011, 43, 335-341.	1.6	8
21	Integrating genomics and phylogenetics in understanding the history of Trichinella species. Veterinary Parasitology, 2009, 159, 210-213.	1.8	16
22	Human dispersal of Trichinella spiralis in domesticated pigs. Infection, Genetics and Evolution, 2008, 8, 799-805.	2.3	64
23	Post-Miocene expansion, colonization, and host switching drove speciation among extant nematodes of the archaic genus Trichinella. Proceedings of the National Academy of Sciences of the United States of America, 2006, 103, 7354-7359.	7.1	142
24	Cytokine responses in immunized and non-immunized calves after Ostertagia ostertagi infection. Parasite Immunology, 2005, 27, 325-331.	1.5	35
25	Recent advances on the taxonomy, systematics and epidemiology of Trichinella. International Journal for Parasitology, 2005, 35, 1191-1204.	3.1	137
26	Inhibition of bovine T lymphocyte responses by extracts of the stomach worm Ostertagia ostertagi. Veterinary Parasitology, 2004, 120, 199-214.	1.8	30
27	Molecular identification of natural hybrids between Trichinella nativa and Trichinella T6 provides evidence of gene flow and ongoing genetic divergence. International Journal for Parasitology, 2003, 33, 209-216.	3.1	44
28	A single, multiplex PCR for differentiating all species ofTrichinella. Parasite, 2001, 8, S24-S26.	2.0	26
29	A multiplex PCR assay for differentiating economically important gastrointestinal nematodes of cattle. Veterinary Parasitology, 2001, 97, 201-211.	1.8	81
30	Trichinella pseudospiralis populations of the Palearctic region and their relationship with populations of the Nearctic and Australian regions. International Journal for Parasitology, 2001, 31, 297-305.	3.1	53
31	Molecular and biochemical methods for parasite differentiation within the genus Trichinella. Veterinary Parasitology, 2000, 93, 279-292.	1.8	21
32	A multiplex PCR for unequivocal differentiation of all encapsulated and non-encapsulated genotypes of Trichinella. International Journal for Parasitology, 1999, 29, 1859-1867.	3.1	269
33	Identification and semi-quantitation of Ostertagia ostertagi eggs by enzymatic amplification of ITS-1 sequences. Veterinary Parasitology, 1998, 77, 245-257.	1.8	35
34	Characterization of protective immune responses in local lymphoid tissues after drug-attenuated infections with Ostertagia ostertagi in calves. Veterinary Parasitology, 1998, 80, 53-64.	1.8	27
35	Comparisons of two polymorphic species of Ostertagia and phylogenetic relationships within the Ostertagiinae (Nematoda: Trichostrongyloidea) inferred from ribosomal DNA repeat and mitochondrial DNA sequences. Journal of Parasitology, 1998, 84, 806-12.	0.7	18
36	Cytokine profile induced by a primary infection with Ostertagia ostertagi in cattle. Veterinary Immunology and Immunopathology, 1997, 58, 63-75.	1.2	51

#	Article	IF	CITATIONS
37	Isolation and phenotypic characterization of abomasal mucosal lymphocytes in the course of a primary Ostertagia ostertagi infection in calves. Veterinary Immunology and Immunopathology, 1997, 57, 87-98.	1.2	27
38	Variations in microsatellite sequences provide evidence for population differences and multiple ribosomal gene repeats within Trichinella pseudospiralis. Journal of Parasitology, 1996, 82, 534-8.	0.7	9
39	Enzymatic amplification and molecular cloning of cDNA encoding the small and large subunits of bovine interleukin 12. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 1995, 1270, 215-217.	3.8	30
40	Method for constructing internal standards for use in competitive PCR. BioTechniques, 1995, 19, 324-6.	1.8	16
41	A Taenia crassiceps cDNA sequence encoding a putative immunodiagnostic antigen for bovine cysticercosis. Molecular and Biochemical Parasitology, 1994, 67, 215-223.	1.1	34
42	Cloning and Characterization of Ribosomal RNA Genes from Three Species of Haemonchus (Nematoda:) Tj ETQq0 Parasitology, 1994, 78, 28-36.	0 0 rgBT 1.2	/Overlock 10 67
43	Cloning and sequence analysis of the small subunit ribosomal RNA gene from Nematodirus battus. Journal of Parasitology, 1994, 80, 342-4.	0.7	0
44	The identification and characterization of a break within the large subunit ribosomal RNA of Trichinella spiralis: comparison of gap sequences within the genus. Molecular and Biochemical Parasitology, 1992, 51, 281-289.	1.1	48
45	Characterization and detection of a newly described Asian taeniid using cloned ribosomal DNA fragments and sequence amplification by the polymerase chain reaction. Experimental Parasitology, 1991, 72, 174-183.	1.2	72
46	A repetitive DNA probe specific for a North American sylvatic genotype of Trichinella. Molecular and Biochemical Parasitology, 1991, 48, 131-137.	1.1	40

47	The differentiation of a newly described Asian taeniid from Taenia saginata using enzymatically amplified non-transcribed ribosomal DNA repeat sequences. Southeast Asian Journal of Tropical Medicine and Public Health, 1991, 22 Suppl, 251-5.	1.0	4
48	DNA analysis in the diagnosis of infection and in the speciation of nematodes parasites. OIE Revue Scientifique Et Technique, 1990, 9, 533-554.	1.2	17