

Akihide Ohkuchi

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6635711/akihide-ohkuchi-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

161
papers

2,843
citations

27
h-index

48
g-index

164
ext. papers

3,268
ext. citations

3.1
avg, IF

4.55
L-index

#	Paper	IF	Citations
161	Human villous trophoblasts express and secrete placenta-specific microRNAs into maternal circulation via exosomes. <i>Biology of Reproduction</i> , 2009 , 81, 717-29	3.9	373
160	Guidelines for obstetrical practice in Japan: Japan Society of Obstetrics and Gynecology (JSOG) and Japan Association of Obstetricians and Gynecologists (JAOG) 2014 edition. <i>Journal of Obstetrics and Gynaecology Research</i> , 2014 , 40, 1469-99	1.9	220
159	Hydroxysteroid (17- β)dehydrogenase 1 is dysregulated by miR-210 and miR-518c that are aberrantly expressed in preeclamptic placentas: a novel marker for predicting preeclampsia. <i>Hypertension</i> , 2012 , 59, 265-73	8.5	135
158	Important surgical measures and techniques at cesarean hysterectomy for placenta previa accreta. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2013 , 92, 372-7	3.8	107
157	Guidelines for obstetrical practice in Japan: Japan Society of Obstetrics and Gynecology (JSOG) and Japan Association of Obstetricians and Gynecologists (JAOG) 2011 edition. <i>Journal of Obstetrics and Gynaecology Research</i> , 2011 , 37, 1174-97	1.9	96
156	Uterine compression sutures for postpartum hemorrhage: an overview. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2013 , 92, 378-85	3.8	81
155	Alterations in placental growth factor levels before and after the onset of preeclampsia are more pronounced in women with early onset severe preeclampsia. <i>Hypertension Research</i> , 2007 , 30, 151-9	4.7	73
154	Evaluation of a new and automated electrochemiluminescence immunoassay for plasma sFlt-1 and PlGF levels in women with preeclampsia. <i>Hypertension Research</i> , 2010 , 33, 422-7	4.7	71
153	Effect of recombinant placental growth factor 2 on hypertension induced by full-length mouse soluble fms-like tyrosine kinase 1 adenoviral vector in pregnant mice. <i>Hypertension</i> , 2009 , 54, 1129-35	8.5	71
152	Human exosomal placenta-associated miR-517a-3p modulates the expression of PRKG1 mRNA in Jurkat cells. <i>Biology of Reproduction</i> , 2014 , 91, 129	3.9	48
151	Effect of maternal age on blood loss during parturition: a retrospective multivariate analysis of 10,053 cases. <i>Journal of Perinatal Medicine</i> , 2003 , 31, 209-15	2.7	48
150	Nanosilica-induced placental inflammation and pregnancy complications: Different roles of the inflammasome components NLRP3 and ASC. <i>Nanotoxicology</i> , 2015 , 9, 554-67	5.3	46
149	Palmitic acid induces interleukin-1 β secretion via NLRP3 inflammasomes and inflammatory responses through ROS production in human placental cells. <i>Journal of Reproductive Immunology</i> , 2016 , 116, 104-12	4.2	46
148	NLRP3 Deficiency Improves Angiotensin II-Induced Hypertension But Not Fetal Growth Restriction During Pregnancy. <i>Endocrinology</i> , 2015 , 156, 4281-92	4.8	44
147	Normal and high-normal blood pressures, but not body mass index, are risk factors for the subsequent occurrence of both preeclampsia and gestational hypertension: a retrospective cohort study. <i>Hypertension Research</i> , 2006 , 29, 161-7	4.7	42
146	Anterior placentation as a risk factor for massive hemorrhage during cesarean section in patients with placenta previa. <i>Journal of Obstetrics and Gynaecology Research</i> , 2014 , 40, 1243-8	1.9	40
145	Onset threshold of the plasma levels of soluble fms-like tyrosine kinase 1/placental growth factor ratio for predicting the imminent onset of preeclampsia within 4 weeks after blood sampling at 19-31 weeks of gestation. <i>Hypertension Research</i> , 2013 , 36, 1073-80	4.7	39

144	Establishing reference values for both total soluble Fms-like tyrosine kinase 1 and free placental growth factor in pregnant women. <i>Hypertension Research</i> , 2005 , 28, 727-32	4.7	38
143	Vaginal lactobacilli and preterm birth. <i>Journal of Perinatal Medicine</i> , 2002 , 30, 458-66	2.7	38
142	Alteration of serum soluble endoglin levels after the onset of preeclampsia is more pronounced in women with early-onset. <i>Hypertension Research</i> , 2008 , 31, 1541-8	4.7	37
141	Threshold of soluble fms-like tyrosine kinase 1/placental growth factor ratio for the imminent onset of preeclampsia. <i>Hypertension</i> , 2011 , 58, 859-66	8.5	36
140	Opening the bladder for cesarean hysterectomy for placenta previa percreta with bladder invasion. <i>Journal of Obstetrics and Gynaecology Research</i> , 2009 , 35, 359-63		33
139	Endogenous and exogenous miR-520c-3p modulates CD44-mediated extravillous trophoblast invasion. <i>Placenta</i> , 2017 , 50, 25-31	3.4	30
138	Antithrombin III activity and platelet count are more likely to decrease in twin pregnancies than in singleton pregnancies. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2002 , 81, 840-5	3.8	30
137	Predicting the risk of pre-eclampsia and a small-for-gestational-age infant by quantitative assessment of the diastolic notch in uterine artery flow velocity waveforms in unselected women. <i>Ultrasound in Obstetrics and Gynecology</i> , 2000 , 16, 171-8	5.8	28
136	AGEs and HMGB1 Increase Inflammatory Cytokine Production from Human Placental Cells, Resulting in an Enhancement of Monocyte Migration. <i>American Journal of Reproductive Immunology</i> , 2016 , 75, 557-68	3.8	28
135	Interferon-tau attenuates uptake of nanoparticles and secretion of interleukin-1 β in macrophages. <i>PLoS ONE</i> , 2014 , 9, e113974	3.7	27
134	Gestational hypertension as a subclinical preeclampsia in view of serum levels of angiogenesis-related factors. <i>Hypertension Research</i> , 2011 , 34, 212-7	4.7	26
133	Serum sFlt1:PlGF ratio, PlGF, and soluble endoglin levels in gestational proteinuria. <i>Hypertension in Pregnancy</i> , 2009 , 28, 95-108	2	24
132	Prediction and prevention of hypertensive disorders of pregnancy. <i>Hypertension Research</i> , 2017 , 40, 5-14.	4.7	23
131	Relationship between blood pressure level in early pregnancy and subsequent changes in blood pressure during pregnancy. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2002 , 81, 918-25	3.8	22
130	Expansion of the fraction of Th1 cells in women with preeclampsia: inverse correlation between the percentage of Th1 cells and the plasma level of PAI-2. <i>American Journal of Reproductive Immunology</i> , 2001 , 46, 252-9	3.8	22
129	Value of adding second-trimester uterine artery Doppler to patient characteristics in identification of nulliparous women at increased risk for pre-eclampsia: an individual patient data meta-analysis. <i>Ultrasound in Obstetrics and Gynecology</i> , 2013 , 42, 257-67	5.8	21
128	A novel 'uterine sandwich' for haemorrhage at caesarean section for placenta praevia. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , 2014 , 54, 283-6	1.7	20
127	Plasma level of hydroxysteroid (17- β)dehydrogenase 1 in the second trimester is an independent risk factor for predicting preeclampsia after adjusting for the effects of mean blood pressure, bilateral notching and plasma level of soluble fms-like tyrosine kinase 1/placental growth factor ratio. <i>Hypertension Research</i> , 2012 , 35, 1152-8	4.7	20

126	Establishing a reference value for the frequency of fetal movements using modified 'count to 10' method. <i>Journal of Obstetrics and Gynaecology Research</i> , 2008 , 34, 318-23		19
125	Matsubara-Takahashi cervix-holding technique for massive postpartum hemorrhage in patients with placenta previa with or without placenta accreta spectrum disorders. <i>International Journal of Gynecology and Obstetrics</i> , 2018 , 140, 357-364	4	19
124	Calculating probability of requiring allogeneic blood transfusion using three preoperative risk factors on cesarean section for placenta previa. <i>Archives of Gynecology and Obstetrics</i> , 2015 , 291, 281-5	2.5	18
123	Advanced glycation end products regulate interleukin-1 β production in human placenta. <i>Journal of Reproduction and Development</i> , 2017 , 63, 401-408	2.1	18
122	Establishing reference values for mean notch depth index, pulsatility index and resistance index in the uterine artery at 16-23 weeks' gestation. <i>Journal of Obstetrics and Gynaecology Research</i> , 2012 , 38, 1275-85	1.9	18
121	A novel method for evaluating uterine cervical consistency using vaginal ultrasound gray-level histogram. <i>Journal of Perinatal Medicine</i> , 2010 , 38, 491-4	2.7	18
120	Fetal heart rate pattern reflecting the severity of placental abruption. <i>Archives of Gynecology and Obstetrics</i> , 2008 , 277, 249-53	2.5	18
119	The risk of birth defects in dichorionic twins conceived by assisted reproductive technology. <i>Twin Research and Human Genetics</i> , 2004 , 7, 223-7		18
118	Extravillous trophoblast cell invasion is promoted by the CD44-hyaluronic acid interaction. <i>Placenta</i> , 2014 , 35, 163-70	3.4	17
117	Causes of stillbirth: an analysis of 77 cases. <i>Journal of Obstetrics and Gynaecology Research</i> , 1999 , 25, 419-24		17
116	Moderate Hypoxia Down-Regulates Interleukin-6 Secretion and TLR4 Expression in Human Sw.71 Placental Cells. <i>Cellular Physiology and Biochemistry</i> , 2015 , 36, 2149-60	3.9	16
115	Spontaneous resolution of post-delivery or post-abortion uterine artery pseudoaneurysm: A report of three cases. <i>Journal of Obstetrics and Gynaecology Research</i> , 2016 , 42, 730-733	1.9	16
114	A novel three-step approach for predicting the imminent onset of preeclampsia within 4 weeks after blood sampling at 19-31 weeks of gestation. <i>Hypertension Research</i> , 2014 , 37, 519-25	4.7	16
113	Blood loss in low-lying placenta: placental edge to cervical internal os distance of less vs. more than 2 cm. <i>Journal of Perinatal Medicine</i> , 2008 , 36, 507-12	2.7	16
112	Maternal peripheral blood natural killer cells incorporate placenta-associated microRNAs during pregnancy. <i>International Journal of Molecular Medicine</i> , 2015 , 35, 1511-24	4.4	15
111	Uterine rupture during pregnancy soon after a laparoscopic adenomyomectomy. <i>Reproductive Medicine and Biology</i> , 2007 , 6, 175-177	4.1	15
110	Isolated gestational proteinuria preceding the diagnosis of preeclampsia - an observational study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2016 , 95, 1048-54	3.8	15
109	Hemostatic effect of intrauterine balloon for postpartum hemorrhage with special reference to concomitant use of "holding the cervix" procedure (Matsubara). <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2017 , 210, 281-285	2.4	13

108	Oxygen concentration modulates cellular senescence and autophagy in human trophoblast cells. <i>American Journal of Reproductive Immunology</i> , 2018 , 79, e12826	3.8	13
107	Fetus with long QT syndrome manifested by tachyarrhythmia: a case report. <i>Prenatal Diagnosis</i> , 1999 , 19, 990-2	3.2	13
106	Vaginal Ureaplasma species increase chorioamnionitis in very preterm infants with preterm premature rupture of the membranes at European Journal of Clinical Microbiology and Infectious Diseases, 2018 , 37, 2371-2380	5.3	13
105	Galectin-1 as a novel risk factor for both gestational hypertension and preeclampsia, specially its expression at a low level in the second trimester and a high level after onset. <i>Hypertension Research</i> , 2018 , 41, 45-52	4.7	12
104	Ambulatory BP monitoring and clinic BP in predicting small-for-gestational-age infants during pregnancy. <i>Journal of Human Hypertension</i> , 2016 , 30, 62-7	2.6	12
103	The ovarian artery: an unusual feeding artery of uterine artery pseudoaneurysm necessitating repetitive transarterial embolisation. <i>Journal of Obstetrics and Gynaecology</i> , 2015 , 35, 656-7	1.3	11
102	Adenomyomectomy, curettage, and then uterine artery pseudoaneurysm occupying the entire uterine cavity. <i>Journal of Obstetrics and Gynaecology Research</i> , 2013 , 39, 1103-6	1.9	11
101	Granulocyte-colony stimulating factor for the treatment of ritodrine-induced neutropenia. <i>Journal of Obstetrics and Gynaecology Research</i> , 2008 , 34, 286-90		11
100	Prolonged activated partial thromboplastin time in thromboprophylaxis with unfractionated heparin in patients undergoing cesarean section. <i>Journal of Obstetrics and Gynaecology Research</i> , 2010 , 36, 58-63	1.9	10
99	Fetus with heart failure due to congenital atrioventricular block treated by maternally administered ritodrine. <i>Archives of Gynecology and Obstetrics</i> , 2008 , 278, 85-8	2.5	10
98	Increase in systolic blood pressure of > or = 30 mm Hg and/or diastolic blood pressure of > or = 15 mm Hg during pregnancy: is it pathologic?. <i>Hypertension in Pregnancy</i> , 2003 , 22, 275-85	2	10
97	Preeclamptic patient-derived circulating cell-free DNA activates the production of inflammatory cytokines via toll-like receptor 9 signalling in the human placenta. <i>Journal of Hypertension</i> , 2019 , 37, 2452-2460 ¹⁰	1.9	10
96	Olive Leaf Extract (OleaVita) Suppresses Inflammatory Cytokine Production and NLRP3 Inflammasomes in Human Placenta. <i>Nutrients</i> , 2019 , 11,	6.7	9
95	Palmitic acid activates NLRP3 inflammasome and induces placental inflammation during pregnancy in mice. <i>Journal of Reproduction and Development</i> , 2020 , 66, 241-248	2.1	9
94	Markedly higher sFlt-1/PlGF ratio in a woman with acute fatty liver of pregnancy compared with HELLP syndrome. <i>Journal of Obstetrics and Gynaecology Research</i> , 2019 , 45, 96-103	1.9	9
93	Biophysical interaction between blood pressure and uterine artery Doppler for the occurrence of early-onset preeclampsia: A prospective cohort study. <i>Pregnancy Hypertension</i> , 2013 , 3, 270-7	2.6	9
92	Hofbauer cell activation and its increased glucose-6-phosphate dehydrogenase activity in second trimester-spontaneous abortion: an ultrastructural dual staining enzyme-cytochemical study. <i>American Journal of Reproductive Immunology</i> , 2003 , 49, 202-9	3.8	9
91	Matsubara-Yano suture: a simple uterine compression suture for postpartum hemorrhage during cesarean section. <i>Archives of Gynecology and Obstetrics</i> , 2019 , 299, 113-121	2.5	9

90	Urinary protein-to-creatinine ratio in pregnant women after dipstick testing: prospective observational study. <i>BMC Pregnancy and Childbirth</i> , 2015 , 15, 331	3.2	8
89	Mutual Balance between Vasohibin-1 and Soluble VEGFR-1 in Endothelial Cells. <i>Pharmaceuticals</i> , 2011 , 4, 782-793	5.2	8
88	First-trimester ultrasonographic investigation of cardiovascular anatomy in thoracoabdominally conjoined twins. <i>Journal of Perinatal Medicine</i> , 2001 , 29, 77-80	2.7	8
87	Factors Contributing to Massive Blood Loss on Peripartum Hysterectomy for Abnormally Invasive Placenta: Who Bleeds More?. <i>Obstetrics and Gynecology International</i> , 2016 , 2016, 5349063	2	8
86	Temporary hypertension and white coat hypertension in the first trimester as risk factors for preeclampsia. <i>Hypertension Research</i> , 2019 , 42, 2002-2012	4.7	7
85	Which type of placenta previa requires blood transfusion more frequently? A new concept of indiscernible edge total previa. <i>Journal of Obstetrics and Gynaecology Research</i> , 2016 , 42, 1502-1508	1.9	7
84	Additive effects of mean blood pressure and bilateral notching in the second trimester on subsequent angiogenesis-related factors. <i>Hypertension Research</i> , 2014 , 37, 76-81	4.7	7
83	Effects of campaign for postpartum vaccination on seronegative rate against rubella among Japanese women. <i>BMC Infectious Diseases</i> , 2014 , 14, 152	4	7
82	Initial dose of vancomycin based on body weight and creatinine clearance to minimize inadequate trough levels in Japanese adults. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2012 , 31, 2537-43	5.3	7
81	Acupuncture for premenstrual dysphoric disorder. <i>Archives of Gynecology and Obstetrics</i> , 2009 , 280, 877-883	3.5	7
80	Irregular peak-to-peak intervals between ascending aortic flows during fetal ventricular tachycardia in long QT syndrome. <i>Ultrasound in Obstetrics and Gynecology</i> , 2009 , 33, 118-20	5.8	7
79	Hydrocephalus after intraventricular hemorrhage in eclamptic woman with HELLP syndrome. <i>Hypertension in Pregnancy</i> , 2006 , 25, 255-7	2	7
78	Clinical usefulness of serum levels of soluble fms-like tyrosine kinase 1/placental growth factor ratio to rule out preeclampsia in women with new-onset lupus nephritis during pregnancy. <i>CEN Case Reports</i> , 2019 , 8, 95-100	1	7
77	Conservative management of retained products of conception in the normal placental position: A retrospective observational study. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2019 , 240, 87-92	2.4	6
76	Increased serum levels of sFlt-1/PlGF ratio in preeclamptic women with onset at . <i>Pregnancy Hypertension</i> , 2018 , 12, 96-103	2.6	6
75	Congenital mesoblastic nephroma: Its diverse clinical features - A literature review with a case report. <i>Journal of Obstetrics and Gynaecology</i> , 2016 , 36, 340-4	1.3	6
74	The Frequency of Peripheral Blood CD4FoxP3 Regulatory T Cells in Women With Pre-eclampsia and Those With High-risk Factors for Pre-eclampsia. <i>Hypertension in Pregnancy</i> , 2015 , 34, 443-455	2	6
73	Maternal manifestation of Ballantyne's syndrome occurring concomitantly with the development of fetal congenital mesoblastic nephroma. <i>Journal of Obstetrics and Gynaecology Research</i> , 2014 , 40, 1114-7	1.9	6

72	Validation and development of models using clinical, biochemical and ultrasound markers for predicting pre-eclampsia: an individual participant data meta-analysis. <i>Health Technology Assessment</i> , 2020 , 24, 1-252	4.4	6
71	Aggregation of Human Trophoblast Cells into Three-Dimensional Culture System Enhances Anti-Inflammatory Characteristics through Cytoskeleton Regulation. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	6
70	Drainage failure of Bakri balloon: no drainage does not indicate no bleeding. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2015 , 94, 336	3.8	5
69	Postpartum hemorrhage: is angiographically detectable "sac" mandatory for diagnosis of ruptured pseudoaneurysm?. <i>Archives of Gynecology and Obstetrics</i> , 2016 , 293, 1361-2	2.5	5
68	A trio of risk factors for the onset of preeclampsia in the second and early third trimesters. <i>Pregnancy Hypertension</i> , 2014 , 4, 224-30	2.6	5
67	Ehlers-Danlos type IV in pregnancy with a history of myocardial infarction. <i>Journal of Obstetrics and Gynaecology Research</i> , 2009 , 35, 797-800		5
66	Liver dysfunction in late pregnancy: cytomegalovirus-induced hepatitis or the HELLP syndrome?. <i>Journal of Obstetrics and Gynaecology Research</i> , 2001 , 27, 319-23		5
65	Influence of chorioamnionitis on survival and morbidity in singletons live-born at . <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2000 , 79, 861-865	3.8	5
64	External validation of prognostic models predicting pre-eclampsia: individual participant data meta-analysis. <i>BMC Medicine</i> , 2020 , 18, 302	11.4	4
63	Effect of eicosapentaenoic acid agent on aggravated hypertriglyceridemia during pregnancy. <i>Journal of Obstetrics and Gynaecology Research</i> , 2013 , 39, 1541-4	1.9	4
62	C-reactive protein levels at pre-/post-indicated cervical cerclage predict very preterm birth. <i>Journal of Perinatal Medicine</i> , 2011 , 39, 151-5	2.7	4
61	Laterally-positioned placenta in placenta previa. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2020 , 33, 2642-2648	2	4
60	Serum soluble LIGHT in the early third trimester as a novel biomarker for predicting late-onset preeclampsia. <i>Pregnancy Hypertension</i> , 2018 , 14, 174-176	2.6	4
59	LncRNA -Derived Accelerates the Invasion of Extravillous Trophoblast Cells by Inhibiting and Subsequently Activating Matrix Metalloproteinases. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	4
58	Supporting rural remote physicians to conduct a study and write a paper: experience of Clinical Research Support Team (CRST)-Jichi. <i>Rural and Remote Health</i> , 2014 , 14, 2883	1.3	4
57	Synergic interaction between ritodrine and magnesium sulfate on the occurrence of critical neonatal hyperkalemia: A Japanese nationwide retrospective cohort study. <i>Scientific Reports</i> , 2020 , 10, 7804	4.9	3
56	Magnetic resonance imaging of degeneration of uterine adenomyosis during pregnancy and post-partum period. <i>Journal of Obstetrics and Gynaecology Research</i> , 2018 , 44, 1169-1173	1.9	3
55	Urinary protein-to-creatinine ratio indicative of significant proteinuria in normotensive pregnant women. <i>Journal of Obstetrics and Gynaecology Research</i> , 2016 , 42, 784-8	1.9	3

54	Changes in Central Hemodynamics in Women With Hypertensive Pregnancy Between Before and After Delivery. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 329-36	2.3	3
53	Immune status among Japanese during nationwide rubella outbreak in Japan 2012-2013. <i>Journal of Infection</i> , 2014 , 68, 300-2	18.9	3
52	Need for systematic classification of various uterine compression sutures. <i>Journal of Obstetrics and Gynaecology Research</i> , 2015 , 41, 1676	1.9	3
51	Serious adverse drug reaction in a woman with hyperemesis gravidarum after first exposure to vitamin B complex containing vitamins B1, B6 and B12. <i>Journal of Obstetrics and Gynaecology Research</i> , 2009 , 35, 790-3		3
50	Mirror syndrome associated with hydropic acardius in triplet pregnancy. <i>Fetal Diagnosis and Therapy</i> , 2008 , 24, 429-33	2.4	3
49	Acute promyelocytic leukemia: an unusual cause showing prolonged disseminated intravascular coagulation after placental abruption. <i>Archives of Gynecology and Obstetrics</i> , 2008 , 277, 267-70	2.5	3
48	Intrauterine death of one twin, with rescue of the other, in twin-twin transfusion syndrome. <i>Ultrasound in Obstetrics and Gynecology</i> , 2002 , 19, 293-6	5.8	3
47	Woman with postpartum ventricular tachycardia and hypomagnesemia. <i>Journal of Obstetrics and Gynaecology Research</i> , 2003 , 29, 92-5		3
46	Neonatal nucleated red blood cell counts in twins. <i>Journal of Perinatal Medicine</i> , 2001 , 29, 144-50	2.7	3
45	Extravillous trophoblast invasion accelerated by WNT3A, 5A, and 10B via CD44. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2021 , 34, 3377-3385	2	3
44	Independent risk factors for a small placenta and a small-for-gestational-age infant at 35-41 weeks of gestation: An association with circulating angiogenesis-related factor levels at 19-31 weeks of gestation. <i>Journal of Obstetrics and Gynaecology Research</i> , 2017 , 43, 1285-1292	1.9	2
43	Intrauterine ritodrine exposure and childhood asthma: Is ritodrine a real culprit?. <i>Pediatric Allergy and Immunology</i> , 2018 , 29, 332-333	4.2	2
42	Thrombus just beneath a retrievable inferior vena cava filter in a pregnant woman with deep vein thrombosis: its removal requiring catheter thrombus fragmentation with fibrinolysis. <i>Journal of Obstetrics and Gynaecology Research</i> , 2014 , 40, 590-4	1.9	2
41	Respiratory Arrest in an Obese Pregnant Woman with Hyperemesis Gravidarum. <i>Case Reports in Obstetrics and Gynecology</i> , 2015 , 2015, 278391	0.8	2
40	Prolapse of the small intestine from the uterine perforation at dilatation and curettage. <i>Case Reports in Obstetrics and Gynecology</i> , 2014 , 2014, 164356	0.8	2
39	Serum leptin concentration during in vitro fertilization treatment with controlled ovarian hyperstimulation: Serum leptin concentration decreases in the early luteal phase in pregnancy. <i>Reproductive Medicine and Biology</i> , 2003 , 2, 177-182	4.1	2
38	Perioperative changes in plasma antithrombin activity and platelet counts in patients undergoing gynecologic surgery. <i>Seminars in Thrombosis and Hemostasis</i> , 2002 , 28, 519-24	5.3	2
37	Video image: Matsubara's Nelaton and Fishing methods for easier Bakri balloon insertion and avoiding its prolapse during cesarean section. <i>Hypertension Research in Pregnancy</i> , 2018 , 6, 73-75	0.7	2

36	Timing of surgery for placenta previa with suspected abnormally invasive placentation: a test of team competency?. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2017 , 96, 1029	3.8	1
35	Association of antenatal antithrombin activity with perinatal liver dysfunction: A prospective multicenter study. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017 , 32, 1378-1386	4	1
34	Interstitial Pregnancy in the Third Trimester with Severe Preeclampsia: A Case Report and Literature Review. <i>Case Reports in Obstetrics and Gynecology</i> , 2020 , 2020, 9408501	0.8	1
33	Re: Peripartum hysterectomy: an economic analysis of direct healthcare costs using routinely collected data: Some concerns from obstetricians' viewpoint. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2018 , 125, 905-906	3.7	1
32	Racial difference in postpartum hemorrhage outcome: pathophysiologic, clinical, and social significance?. <i>American Journal of Obstetrics and Gynecology</i> , 2018 , 219, 515-516	6.4	1
31	Absent Ductus Venosus Associated with Partial Liver Defect. <i>Case Reports in Obstetrics and Gynecology</i> , 2018 , 2018, 6591025	0.8	1
30	Increased antiangiogenic factors in severe proteinuria without hypertension in pregnancy: is kidney biopsy necessary?. <i>CEN Case Reports</i> , 2014 , 3, 86-89	1	1
29	Is it time to classify various uterine compression suture techniques?. <i>Archives of Gynecology and Obstetrics</i> , 2013 , 288, 1195-6	2.5	1
28	Response to Dr Sabour: 'Prediction and prevention of hypertensive disorders of pregnancy: a methodological mistake'. <i>Hypertension Research</i> , 2017 , 40, 791	4.7	1
27	Maternal Perception of Decreased Fetal Movement in One Twin: A Clue Leading to the Early Detection of Absent Variability due to Acute Twin-to-Twin Transfusion Syndrome. <i>Case Reports in Obstetrics and Gynecology</i> , 2013 , 2013, 345808	0.8	1
26	Statistical model predicting a short duration to birth in women with preterm labor at 22-35 weeks' gestation: the importance of large vaginal Gram-positive rods. <i>Journal of Perinatal Medicine</i> , 2009 , 37, 244-50	2.7	1
25	The Risk of Birth Defects in Dichorionic Twins Conceived by Assisted Reproductive Technology		1
24	Decreased circulating levels of plasmacytoid dendritic cells in women with early-onset preeclampsia. <i>Journal of Reproductive Immunology</i> , 2020 , 141, 103170	4.2	1
23	Expectant management of retained products of conception following abortion: A retrospective cohort study. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2021 , 260, 1-5	2.4	1
22	Surgical management of abnormally invasive placenta: is decreased blood loss due to participation of gynecologic oncologists?. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2016 , 95, 119	3.8	1
21	Perinatal outcome of pregnancy after adenomyomectomy: summary of 10 cases with a brief literature review. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2020 , 33, 4145-4149	2	1
20	Economic evaluation of the sFlt-1/PlGF ratio for the short-term prediction of preeclampsia in a Japanese cohort of the PROGNOSIS Asia study. <i>Hypertension Research</i> , 2021 , 44, 822-829	4.7	1
19	Posterior Reversible Encephalopathy Syndrome with Stroke in Puerperal Woman with High Titer of Anti-Phospholipid IgM Antibody. <i>Case Reports in Obstetrics and Gynecology</i> , 2018 , 2018, 7438676	0.8	1

18	Changes in standing body sway of pregnant women after long-term bed rest. <i>Journal of Obstetrics and Gynaecology</i> , 2016 , 36, 479-82	1.3	○
17	Supporting less experienced physicians to write a paper: A proposal to introduce such a system on an academic society basis. <i>Journal of Obstetrics and Gynaecology Research</i> , 2016 , 42, 1891-1892	1.9	○
16	Misquoting Page's classification for placental abruption in Japan?. <i>Journal of Obstetrics and Gynaecology Research</i> , 2011 , 37, 481	1.9	○
15	Relation between viability of vaginal polymorphonuclear leukocytes and presence of histologic chorioamnionitis. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2000 , 79, 818-823	3.8	○
14	Difficulty of predicting early-onset super-imposed preeclampsia in pregnant women with hemodialysis due to diabetic nephropathy by serum levels of sFlt-1, PlGF, and sEng. <i>CEN Case Reports</i> , 2020 , 9, 101-105	1	○
13	Effects of prophylactic vaginal progesterone administration on mild cervical shortening (TROPICAL study): a multicenter, double-blind, randomized trial. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2021 , 1-7	2	○
12	Elevated S100A9 in preeclampsia induces soluble endoglin and IL-1 β secretion and hypertension via the NLRP3 inflammasome. <i>Journal of Hypertension</i> , 2022 , 40, 84-93	1.9	○
11	A randomized phase 3 trial evaluating antithrombin gamma treatment in Japanese patients with early-onset severe preeclampsia (KOUNO-TORI study): Study protocol. <i>Contemporary Clinical Trials</i> , 2021 , 107, 106490	2.3	○
10	Bleeding-related outcomes of low-risk total placenta previa are equivalent to those of partial/marginal placenta previa. <i>Taiwanese Journal of Obstetrics and Gynecology</i> , 2022 , 61, 447-452	1.6	○
9	MicroRNA. <i>Comprehensive Gynecology and Obstetrics</i> , 2018 , 209-224	0	
8	Insertion of the Bakri balloon: The earlier, the better?. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , 2016 , 56, 117	1.7	
7	Is anterior placentation per se a risk for cesarean section-related morbidity in placenta previa?. <i>Journal of Perinatology</i> , 2014 , 34, 649	3.1	
6	Pregnancy-related thromboembolism and contributing risk factors: From 10 years of experience at a Japanese tertiary obstetrics institute. <i>Hypertension Research in Pregnancy</i> , 2017 , 5, 59-64	0.7	
5	Effectiveness of bromocriptine treatment in a patient with peripartum cardiomyopathy. <i>Journal of Cardiology Cases</i> , 2010 , 2, e28-e31	0.6	
4	Hemostatic effect of combined procedures for placenta previa: cervix-holding, intrauterine balloon, and uterine compression suture. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2021 , 1-7	2	
3	The role of NLRP3 inflammasome on preeclampsia. <i>Reproductive Immunology and Biology</i> , 2019 , 34, 1-8	0	○
2	Low placental growth factor levels and high soluble endoglin levels at 26-31 weeks of gestation precede light placenta with and without relatively heavy infant, respectively: A retrospective cohort study. <i>Journal of Obstetrics and Gynaecology Research</i> , 2021 , 47, 1040-1051	1.9	
1	Prognosis and Prognostic Factors of Patients with Emergent Cerclage: A Japanese Single-Center Study.. <i>Obstetrics and Gynecology International</i> , 2021 , 2021, 4351783	2	

