

Simon M Fredholm

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6640381/publications.pdf>

Version: 2024-02-01

22
papers

814
citations

566801

15
h-index

676716

22
g-index

22
all docs

22
docs citations

22
times ranked

1147
citing authors

#	ARTICLE	IF	CITATIONS
1	Antibiotics inhibit tumor and disease activity in cutaneous T-cell lymphoma. <i>Blood</i> , 2019, 134, 1072-1083.	0.6	94
2	Staphylococcal enterotoxin A (SEA) stimulates STAT3 activation and IL-17 expression in cutaneous T-cell lymphoma. <i>Blood</i> , 2016, 127, 1287-1296.	0.6	86
3	Single-cell heterogeneity in SÄ©zary syndrome. <i>Blood Advances</i> , 2018, 2, 2115-2126.	2.5	78
4	Jak3, STAT3, and STAT5 inhibit expression of miR-22, a novel tumor suppressor microRNA, in cutaneous T-Cell lymphoma. <i>Oncotarget</i> , 2015, 6, 20555-20569.	0.8	78
5	Butyrate and propionate inhibit antigen-specific CD8+ T cell activation by suppressing IL-12 production by antigen-presenting cells. <i>Scientific Reports</i> , 2017, 7, 14516.	1.6	77
6	Analysis of STAT4 expression in cutaneous T-cell lymphoma (CTCL) patients and patient-derived cell lines. <i>Cell Cycle</i> , 2014, 13, 2975-2982.	1.3	62
7	MicroRNA expression in early mycosis fungoides is distinctly different from atopic dermatitis and advanced cutaneous T-cell lymphoma. <i>Anticancer Research</i> , 2014, 34, 7207-17.	0.5	55
8	STAT5 induces miR-21 expression in cutaneous T cell lymphoma. <i>Oncotarget</i> , 2016, 7, 45730-45744.	0.8	45
9	SATB1 in Malignant T Cells. <i>Journal of Investigative Dermatology</i> , 2018, 138, 1805-1815.	0.3	38
10	STAT3/5-Dependent IL9 Overexpression Contributes to Neoplastic Cell Survival in Mycosis Fungoides. <i>Clinical Cancer Research</i> , 2016, 22, 3328-3339.	3.2	36
11	Staphylococcal alpha-toxin tilts the balance between malignant and non-malignant CD4⁺ T cells in cutaneous T-cell lymphoma. <i>Oncolmmunology</i> , 2019, 8, e1641387.	2.1	32
12	IL-15 and IL-17F are differentially regulated and expressed in mycosis fungoides (MF). <i>Cell Cycle</i> , 2014, 13, 1306-1312.	1.3	27
13	<i>Staphylococcus aureus</i> alpha-toxin inhibits CD8 ⁺ T cell-mediated killing of cancer cells in cutaneous T-cell lymphoma. <i>Oncolmmunology</i> , 2020, 9, 1751561.	2.1	24
14	A novel BLK-induced tumor model. <i>Tumor Biology</i> , 2017, 39, 101042831771419.	0.8	19
15	Human P2Y11 Expression Level Affects Human P2X7 Receptor-Mediated Cell Death. <i>Frontiers in Immunology</i> , 2018, 9, 1159.	2.2	17
16	STAT3 activation and infiltration of eosinophil granulocytes in mycosis fungoides. <i>Anticancer Research</i> , 2014, 34, 5277-86.	0.5	15
17	The Thioredoxin-Interacting Protein TXNIP Is a Putative Tumour Suppressor in Cutaneous T-Cell Lymphoma. <i>Dermatology</i> , 2021, 237, 283-290.	0.9	8
18	MicroRNA-93 Targets p21 and Promotes Proliferation in Mycosis Fungoides T Cells. <i>Dermatology</i> , 2021, 237, 277-282.	0.9	8

#	ARTICLE	IF	CITATIONS
19	Low SATB1 Expression Promotes IL-5 and IL-9 Expression in SÅ©zary Syndrome. Journal of Investigative Dermatology, 2020, 140, 713-716.	0.3	5
20	The Expression of IL-21 Is Promoted by MEKK4 in Malignant T Cells and Associated with Increased Progression Risk in Cutaneous T-Cell Lymphoma. Journal of Investigative Dermatology, 2016, 136, 866-869.	0.3	4
21	Expression and function of Kv1.3 channel in malignant T cells in SÅ©zary syndrome. Oncotarget, 2019, 10, 4894-4906.	0.8	3
22	Proinflammatory biomarkers are associated with prediabetes in patients with schizophrenia. CNS Spectrums, 2022, 27, 347-354.	0.7	3