

Catherine Curie

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6646642/publications.pdf>

Version: 2024-02-01

41
papers

8,594
citations

147801

31
h-index

265206

42
g-index

42
all docs

42
docs citations

42
times ranked

5490
citing authors

#	ARTICLE	IF	CITATIONS
1	<scp>NRAMP6</scp> and <scp>NRAMP1</scp> cooperatively regulate root growth and manganese translocation under manganese deficiency in Arabidopsis. <i>Plant Journal</i> , 2022, 110, 1564-1577.	5.7	22
2	Manganese triggers phosphorylation-mediated endocytosis of the Arabidopsis metal transporter NRAMP1. <i>Plant Journal</i> , 2021, 106, 1328-1337.	5.7	29
3	AtDTX25, a member of the multidrug and toxic compound extrusion family, is a vacuolar ascorbate transporter that controls intracellular iron cycling in Arabidopsis. <i>New Phytologist</i> , 2021, 231, 1956-1967.	7.3	18
4	<i>Paspalum urvillei</i> and <i>Setaria parviflora</i> , two grasses naturally adapted to extreme iron-rich environments. <i>Plant Physiology and Biochemistry</i> , 2020, 151, 144-156.	5.8	23
5	Split green fluorescent protein as a tool to study infection with a plant pathogen, Cauliflower mosaic virus. <i>PLoS ONE</i> , 2019, 14, e0213087.	2.5	10
6	New routes for plant iron mining. <i>New Phytologist</i> , 2017, 214, 521-525.	7.3	76
7	Phosphatidylinositol 3-phosphate-binding protein AtPH1 controls the localization of the metal transporter NRAMP1 in <i>Arabidopsis</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E3354-E3363.	7.1	54
8	Intracellular Distribution of Manganese by the <i>Trans</i> -Golgi Network Transporter NRAMP2 Is Critical for Photosynthesis and Cellular Redox Homeostasis. <i>Plant Cell</i> , 2017, 29, 3068-3084.	6.6	87
9	The high-affinity metal Transporters NRAMP1 and IRT1 Team up to Take up Iron under Sufficient Metal Provision. <i>Scientific Reports</i> , 2016, 6, 37222.	3.3	131
10	Inventory of metal complexes circulating in plant fluids: a reliable method based on HPLC coupled with dual elemental and high-resolution molecular mass spectrometric detection. <i>New Phytologist</i> , 2016, 211, 1129-1141.	7.3	87
11	Ascorbate Efflux as a New Strategy for Iron Reduction and Transport in Plants. <i>Journal of Biological Chemistry</i> , 2014, 289, 2515-2525.	3.4	153
12	Over-expression of the Bacterial Phytase US417 in Arabidopsis Reduces the Concentration of Phytic Acid and Reveals Its Involvement in the Regulation of Sulfate and Phosphate Homeostasis and Signaling. <i>Plant and Cell Physiology</i> , 2014, 55, 1912-1924.	3.1	23
13	New insights into Fe localization in plant tissues. <i>Frontiers in Plant Science</i> , 2013, 4, 350.	3.6	99
14	The <i>Arabidopsis</i> YELLOW STRIPE LIKE4 and 6 Transporters Control Iron Release from the Chloroplast. <i>Plant Cell</i> , 2013, 25, 1040-1055.	6.6	114
15	Proteasome-mediated turnover of the transcriptional activator FIT is required for plant iron-deficiency responses. <i>Plant Journal</i> , 2011, 66, 1044-1052.	5.7	112
16	Ubiquitination of transporters at the forefront of plant nutrition. <i>Plant Signaling and Behavior</i> , 2011, 6, 1597-1599.	2.4	14
17	Plant Cell Nucleolus as a Hot Spot for Iron. <i>Journal of Biological Chemistry</i> , 2011, 286, 27863-27866.	3.4	81
18	The FRD3 Citrate Effluxer Promotes Iron Nutrition between Symplastically Disconnected Tissues throughout <i>Arabidopsis</i> Development. <i>Plant Cell</i> , 2011, 23, 2725-2737.	6.6	147

#	ARTICLE	IF	CITATIONS
19	High-Affinity Manganese Uptake by the Metal Transporter NRAMP1 Is Essential for <i>Arabidopsis</i> Growth in Low Manganese Conditions. <i>Plant Cell</i> , 2010, 22, 904-917.	6.6	449
20	Straightforward histochemical staining of Fe by the adaptation of an old-school technique. <i>Plant Signaling and Behavior</i> , 2010, 5, 56-57.	2.4	21
21	Increased sensitivity to iron deficiency in <i>Arabidopsis thaliana</i> overaccumulating nicotianamine. <i>Journal of Experimental Botany</i> , 2009, 60, 1249-1259.	4.8	66
22	Identification of the Endodermal Vacuole as the Iron Storage Compartment in the <i>Arabidopsis</i> Embryo. <i>Plant Physiology</i> , 2009, 151, 1329-1338.	4.8	203
23	<i>Arabidopsis</i> IRT2 cooperates with the high-affinity iron uptake system to maintain iron homeostasis in root epidermal cells. <i>Planta</i> , 2009, 229, 1171-1179.	3.2	161
24	Metal movement within the plant: contribution of nicotianamine and yellow stripe 1-like transporters. <i>Annals of Botany</i> , 2009, 103, 1-11.	2.9	703
25	The NRAMP6 metal transporter contributes to cadmium toxicity. <i>Biochemical Journal</i> , 2009, 422, 217-228.	3.7	235
26	Cytokinins negatively regulate the root iron uptake machinery in <i>Arabidopsis</i> through a growth-dependent pathway. <i>Plant Journal</i> , 2008, 55, 289-300.	5.7	188
27	Iron Acquisition from Fe-Pyoverdine by <i>Arabidopsis thaliana</i> . <i>Molecular Plant-Microbe Interactions</i> , 2007, 20, 441-447.	2.6	225
28	Iron utilization and metabolism in plants. <i>Current Opinion in Plant Biology</i> , 2007, 10, 276-282.	7.1	374
29	Mobilization of vacuolar iron by AtNRAMP3 and AtNRAMP4 is essential for seed germination on low iron. <i>EMBO Journal</i> , 2005, 24, 4041-4051.	7.8	562
30	A loss-of-function mutation in AtYSL1 reveals its role in iron and nicotianamine seed loading. <i>Plant Journal</i> , 2005, 44, 769-782.	5.7	238
31	A Putative Function for the <i>Arabidopsis</i> Fe-Phytosiderophore Transporter Homolog AtYSL2 in Fe and Zn Homeostasis. <i>Plant and Cell Physiology</i> , 2005, 46, 762-774.	3.1	163
32	Dual Regulation of the <i>Arabidopsis</i> High-Affinity Root Iron Uptake System by Local and Long-Distance Signals. <i>Plant Physiology</i> , 2003, 132, 796-804.	4.8	262
33	IRT1, an <i>Arabidopsis</i> Transporter Essential for Iron Uptake from the Soil and for Plant Growth. <i>Plant Cell</i> , 2002, 14, 1223-1233.	6.6	1,464
34	<i>Arabidopsis</i> IRT2 gene encodes a root-periphery iron transporter. <i>Plant Journal</i> , 2001, 26, 181-189.	5.7	272
35	Maize yellow stripe1 encodes a membrane protein directly involved in Fe(III) uptake. <i>Nature</i> , 2001, 409, 346-349.	27.8	905
36	Involvement of NRAMP1 from <i>Arabidopsis thaliana</i> in iron transport. <i>Biochemical Journal</i> , 2000, 347, 749-755.	3.7	474

#	ARTICLE	IF	CITATIONS
37	Molecular biology of male gametogenesis. <i>Euphytica</i> , 1994, 79, 245-250.	1.2	6
38	Modular organization and developmental activity of an <i>Arabidopsis thaliana</i> EF-1 α gene promoter. <i>Molecular Genetics and Genomics</i> , 1993, 238, 428-436.	2.4	85
39	The activation process of <i>Arabidopsis thaliana</i> A1 gene encoding the translation elongation factor EF-1 α is conserved among angiosperms. <i>Plant Molecular Biology</i> , 1992, 18, 1083-1089.	3.9	15
40	Cis and trans-acting elements involved in the activation of <i>Arabidopsis thaliana</i> A1 gene encoding the translation elongation factor EF-1 α . <i>Nucleic Acids Research</i> , 1991, 19, 1305-1310.	14.5	72
41	The gene family encoding the <i>Arabidopsis thaliana</i> translation elongation factor EF-1 α : Molecular cloning, characterization and expression. <i>Molecular Genetics and Genomics</i> , 1989, 219, 106-112.	2.4	161