Wilfredo L GonzÃ;les

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6647351/publications.pdf

Version: 2024-02-01

22 papers 503 citations

759233 12 h-index 21 g-index

22 all docs 22 docs citations

times ranked

22

714 citing authors

#	Article	IF	CITATIONS
1	Temperature regime influences accessions and effectiveness of germination promoters in the highâ€Andean crop maca. Agronomy Journal, 2021, 113, 2557-2566.	1.8	1
2	Influence of seed color on germination performance and storage organ color in the high-Andean crop Lepidium meyenii (Maca). Archives of Agronomy and Soil Science, 2020, , 1-11.	2.6	2
3	Sand-swimming behaviour reduces ectoparasitism in an iguanian lizard. Die Naturwissenschaften, 2019, 106, 53.	1.6	1
4	Seed weight predicts seedling emergence, and extremely acid soil and low availability of Phosphorus are associated with poor plant performances in Lepidium meyenii Walpers (maca). Scientia Horticulturae, 2019, 253, 341-348.	3.6	2
5	Genetic variation in the reduction of attractive floral traits of an annual tarweed in response to drought and apical damage. Journal of Plant Ecology, 2016, 9, 629-635.	2.3	5
6	Testing non-additive effects of nectar-robbing ants and hummingbird pollination on the reproductive success of a parasitic plant. Plant Ecology, 2013, 214, 633-640.	1.6	14
7	Local Host Adaptation and Use of a Novel Host in the Seed Beetle Megacerus eulophus. PLoS ONE, 2013, 8, e53892.	2.5	4
8	Community Impacts of Prosopis juliflora Invasion: Biogeographic and Congeneric Comparisons. PLoS ONE, 2012, 7, e44966.	2.5	99
9	Foliar damage modifies floral attractiveness to pollinators in Alstroemeria exerens. Evolutionary Ecology, 2009, 23, 545-555.	1.2	20
10	Induction of glandular and non-glandular trichomes by damage in leaves of Madia sativa under contrasting water regimes. Acta Oecologica, 2008, 33, 128-132.	1.1	71
11	Water availability limits tolerance of apical damage in the Chilean tarweed Madia sativa. Acta Oecologica, 2008, 34, 104-110.	1.1	30
12	Host-associated variation in sexual size dimorphism and fitness effects of adult feeding in a bruchid beetle. Entomologia Experimentalis Et Applicata, 2007, 122, 233-237.	1.4	13
13	Outcrossing increases infection success in the holoparasitic mistletoe Tristerix aphyllus (Loranthaceae). Evolutionary Ecology, 2007, 21, 173-183.	1.2	13
14	Cost and benefits of attractive floral traits in the annual species Madia sativa (Asteraceae). Evolutionary Ecology, 2007, 21, 247-257.	1.2	29
15	Phenotypic plasticity in the holoparasitic mistletoe Tristerix aphyllus (Loranthaceae): consequences of trait variation for successful establishment. Evolutionary Ecology, 2007, 21, 431-444.	1.2	12
16	Natural selection on ecophysiological traits of a fern species in a temperate rainforest. Evolutionary Ecology, 2007, 21, 651-662.	1.2	49
17	A new species of Uroleucon (Hemiptera: Aphididae) on Madia (Asteraceae) in Chile. Neotropical Entomology, 2005, 34, 221-225.	1.2	4
18	Evaluation of induced responses, insect population growth, and host-plant fitness may change the outcome of tests of the preference-performance hypothesis: a case study. Entomologia Experimentalis Et Applicata, 2003, 109, 211-216.	1.4	9

#	Article	IF	CITATIONS
19	Host plant and natural enemy impact on cereal aphid competition in a seasonal environment. Oikos, 2002, 96, 481-491.	2.7	25
20	Historia natural cuantitativa de una relación parásito-hospedero: el sistema Tristerix-cactáceas en Chile semiárido. Revista Chilena De Historia Natural, 2002, 75, 127.	1.2	29
21	Plant quality vs. risk of parasitism: within-plant distribution and performance of the corn leaf aphid, Rhopalosiphum maidis. Agricultural and Forest Entomology, 2001, 3, 29-33.	1.3	15
22	Feeding by the aphid Sipha flava produces a reddish spot on leaves of Sorghum halepense: an induced defense?. Journal of Chemical Ecology, 2001, 27, 273-283.	1.8	56