

Orio Yamamura

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6648128/publications.pdf>

Version: 2024-02-01

32
papers

719
citations

840119

11
h-index

552369

26
g-index

32
all docs

32
docs citations

32
times ranked

865
citing authors

#	ARTICLE	IF	CITATIONS
1	Abundance Estimates of Steller Sea Lions (<i>Eumetopias jubatus</i>) Off the Western Coast of Hokkaido, Japan. <i>Mammal Study</i> , 2021, 46, .	0.2	2
2	Aging steller sea lions by growth layer groups in teeth. <i>Wildlife Society Bulletin</i> , 2019, 43, 238-243.	1.6	1
3	Transported zooplankton from the Okhotsk Sea facilitate feeding and growth of juvenile walleye pollock on a continental shelf along the Oyashio Current, western subarctic Pacific. <i>Marine Biology</i> , 2018, 165, 1.	0.7	3
4	Diets of Steller sea lions off the coast of Hokkaido, Japan: An interdecadal and geographic comparison. <i>Marine Ecology</i> , 2017, 38, e12477.	0.4	6
5	Beyond dichotomy in the protection and management of marine mammals in Japan. <i>Therya</i> , 2015, 6, 283-296.	0.2	7
6	Comparison of factors affecting recruitment variability of walleye pollock <i>Theragra chalcogramma</i> in the Pacific Ocean and the Sea of Japan off northern Japan. <i>Fisheries Science</i> , 2014, 80, 117-126.	0.7	13
7	Abiotic and biotic factors affecting recruitment variability of walleye pollock (<i>Theragra chalcogramma</i>) off the Pacific coast of Hokkaido, Japan. <i>Fisheries Oceanography</i> , 2013, 22, 193-206.	0.9	17
8	Interannual variation in diets of walleye pollock in the Doto area, in relation to climate variation. <i>Marine Ecology - Progress Series</i> , 2013, 491, 221-234.	0.9	8
9	Food habits of threadfin hake <i>Laemonema longipes</i> along the Pacific coast of northern Japan. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2012, 92, 613-621.	0.4	2
10	Winter energy allocation and deficit of juvenile walleye pollock <i>Theragra chalcogramma</i> in the Doto area, northern Japan. <i>Environmental Biology of Fishes</i> , 2012, 94, 389-402.	0.4	21
11	Temporal variation in chum salmon, <i>Oncorhynchus keta</i> , diets in the central Bering Sea in summer and early autumn. <i>Environmental Biology of Fishes</i> , 2012, 93, 319-331.	0.4	4
12	Characterization of an Alternative Chromatin Remodeling to Parasperm in a Cottid Fish, <i>Hemilepidotus gilberti</i> . <i>Zoological Science</i> , 2011, 28, 438-443.	0.3	0
13	The relationship between pink salmon biomass and the body condition of short-tailed shearwaters in the Bering Sea: can fish compete with seabirds?. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2011, 278, 2584-2590.	1.2	25
14	Size-dependent thermal preferences in a pelagic fish. <i>Oikos</i> , 2010, 119, 1265-1272.	1.2	85
15	Biomass and mortality of chum salmon in the pelagic Bering Sea. <i>Marine Ecology - Progress Series</i> , 2010, 403, 219-230.	0.9	4
16	The distribution of Steller sea lions (<i>Eumetopias jubatus</i>) in the Sea of Japan off Hokkaido, Japan: A preliminary report. <i>Marine Mammal Science</i> , 2009, 25, 949-954.	0.9	6
17	Resightings of branded Steller sea lions at wintering haul-out sites in Hokkaido, Japan 2003-2006. <i>Marine Mammal Science</i> , 2009, 26, 698.	0.9	6
18	Winter lipid depletion of juvenile walleye pollock <i>Theragra chalcogramma</i> in the Doto area, northern Japan. <i>Journal of Fish Biology</i> , 2009, 75, 186-202.	0.7	12

#	ARTICLE	IF	CITATIONS
19	Dietary shift and feeding intensity of <i>Stenobrachius leucopsarus</i> in the Bering Sea. <i>Journal of Oceanography</i> , 2008, 64, 185-194.	0.7	9
20	Feeding habits of mesopelagic fish <i>Lampanyctus jordani</i> (Family: Myctophidae) over the continental slope off Tohoku area, northern Japan. <i>Fisheries Science</i> , 2008, 74, 69-76.	0.7	4
21	II-2. Comparison of sampling gears for micronekton in PICES-MIE. <i>Nippon Suisan Gakkaishi</i> , 2007, 73, 931-932.	0.0	3
22	Rate of energy depletion and overwintering mortality of juvenile walleye pollock in cold water. <i>Journal of Fish Biology</i> , 2007, 71, 1714-1734.	0.7	11
23	NEMURO's a lower trophic level model for the North Pacific marine ecosystem. <i>Ecological Modelling</i> , 2007, 202, 12-25.	1.2	293
24	Optimum temperature for growth of juvenile walleye pollock <i>Theragra chalcogramma</i> . <i>Journal of Experimental Marine Biology and Ecology</i> , 2007, 347, 69-76.	0.7	19
25	Contributions of the VENFISH program: meso-zooplankton, Pacific saury (<i>Cololabis saira</i>) and walleye pollock (<i>Theragra chalcogramma</i>) in the northwestern Pacific. <i>Fisheries Oceanography</i> , 2004, 13, 1-9.	0.9	18
26	Trophodynamic modeling of walleye pollock (<i>Theragra chalcogramma</i>) in the Doto area, northern Japan: model description and baseline simulations. <i>Fisheries Oceanography</i> , 2004, 13, 138-154.	0.9	22
27	Assimilation efficiency of Rhinoceros Auklet (<i>Cerorhinca monocerata</i>) chicks fed Japanese anchovy (<i>Engraulis japonicus</i>) and Japanese sand lance (<i>Ammodytes personatus</i>). <i>Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology</i> , 2004, 139, 97-101.	0.8	11
28	Demersal fish fauna in the Doto nearshore waters, northern Japan: Species composition and seasonal variation. <i>Fisheries Science</i> , 2003, 69, 445-455.	0.7	8
29	Elemental analysis of otoliths of walleye pollock: A trial to discriminate the local stocks around Hokkaido, Japan. <i>Fisheries Science</i> , 2003, 69, 1209-1217.	0.7	1
30	Diet of the mesopelagic fish <i>Notoscopelus japonicus</i> (Family: Myctophidae) associated with the continental slope off the Pacific coast of Honshu, Japan. <i>Fisheries Science</i> , 2002, 68, 1034-1040.	0.7	10
31	Diets of walleye pollock <i>Theragra chalcogramma</i> in the Doto area, northern Japan: ontogenetic and seasonal variations. <i>Marine Ecology - Progress Series</i> , 2002, 238, 187-198.	0.9	73
32	Feeding Habits of the Mesopelagic Fish <i>Gonostoma gracile</i> in the Northwestern North Pacific. <i>Journal of Oceanography</i> , 2001, 57, 509-517.	0.7	15