


List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6648962/publications.pdf

Version: 2024-02-01



YONG SU

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Clinical target volume design of postoperative intensity-modulated radiotherapy for major salivary gland tumours according to surgical principles: an innovative method. Journal of Cancer Research and Clinical Oncology, 2022, 148, 921-930. | 2.5 | 0 |
| 2 | Individualized clinical target volume delineation and efficacy analysis in unilateral nasopharyngeal carcinoma treated with intensity-modulated radiotherapy (IMRT): 10-year summary. Journal of Cancer Research and Clinical Oncology, 2022, 148, 1931-1942. | 2.5 | 6 |
| 3 | Failure patterns and prognostic factors for cervical nodeâ€negative nasopharyngeal carcinoma in the intensityâ€modulated radiotherapy era. Asia-Pacific Journal of Clinical Oncology, 2021, 17, 330-337. | 1.1 | 1 |
| 4 | Efficacy of concurrent chemoradiotherapy in subgroups of stage III nasopharyngeal carcinoma: an analysis based on 10-year follow-up. Radiation Oncology, 2021, 16, 215. | 2.7 | 7 |
| 5 | Late-course accelerated hyperfractionated intensity-modulated radiotherapy for nasopharyngeal adenoid cystic carcinoma: A case report. Medical Dosimetry, 2020, 45, 46-51. | 0.9 | 1 |
| 6 | Is Surgery an Inevitable Treatment for Advanced Salivary Lymphoepithelial Carcinoma? Three Case Reports. Ear, Nose and Throat Journal, 2020, 100, 014556132092317. | 0.8 | 8 |
| 7 | Can neoadjuvant chemotherapy improve survival in stage T3-4N1 nasopharyngeal carcinoma? A propensity matched analysis. Radiation Oncology, 2020, 15, 160. | 2.7 | 4 |
| 8 | Target delineation and dose prescription of adaptive replanning intensityâ€nodulated radiotherapy for nasopharyngeal carcinoma. Cancer Communications, 2019, 39, 1-4. | 9.2 | 8 |
| 9 | Reduction of Target Volume and the Corresponding Dose for the Tumor Regression Field after Induction Chemotherapy in Locoregionally Advanced Nasopharyngeal Carcinoma. Cancer Research and Treatment, 2019, 51, 685-695. | 3.0 | 27 |
| 10 | Development and validation of quality of life scale of nasopharyngeal carcinoma patients: the QOL-NPC (version 2). Health and Quality of Life Outcomes, 2016, 14, 76. | 2.4 | 12 |
| 11 | Using CT or MRI to assess locoregional spread to determine the radiotherapy target of hypopharyngeal carcinoma. Asia-Pacific Journal of Clinical Oncology, 2014, 10, e21-7. | 1.1 | 2 |
| 12 | Prognostic value of tumor volume for patients with nasopharyngeal carcinoma treated with concurrent chemotherapy and intensity-modulated radiotherapy. Journal of Cancer Research and Clinical Oncology, 2014, 140, 69-76. | 2.5 | 46 |
| 13 | Prognostic significance of tumor volume in patients with nasopharyngeal carcinoma undergoing intensityâ€modulated radiation therapy. Head and Neck, 2013, 35, 689-694. | 2.0 | 42 |
| 14 | Analysis of risk factors for retropharyngeal lymph node metastasis in carcinoma of the hypopharynx. Head and Neck, 2013, 35, 1274-1277. | 2.0 | 20 |
| 15 | Analysis of cervical and retropharyngeal lymph node metastases in the patients with hypopharyngeal carcinoma with computed tomography and magnetic resonance imaging. Chinese Journal of Cancer, 2010, 29, 189-193. | 4.9 | 8 |