## Michele Carraro

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6673633/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	FP115KIDNEY BIOPSY IN THE ELDERLY: FRAILTY INDEX IS A GOOD TOOL TO PREDICT CLINICAL OUTCOMES. Nephrology Dialysis Transplantation, 2018, 33, i15-i16.	0.7	0
2	Hemicentin 1 influences podocyte dynamic changes in glomerular diseases. American Journal of Physiology - Renal Physiology, 2018, 314, F1154-F1165.	2.7	11
3	The renal phenotype of allopurinol-treated HPRT-deficient mouse. PLoS ONE, 2017, 12, e0173512.	2.5	8
4	Soluble CD40 ligand directly alters glomerular permeability and may act as a circulating permeability factor in FSGS. PLoS ONE, 2017, 12, e0188045.	2.5	25
5	A nanoporous surface is essential for glomerular podocyte differentiation in three-dimensional culture. International Journal of Nanomedicine, 2016, Volume 11, 4957-4973.	6.7	11
6	Rapid and cost-effective xenograft hepatocellular carcinoma model in Zebrafish for drug testing. International Journal of Pharmaceutics, 2016, 515, 583-591.	5.2	21
7	Podocyte Developmental Defects Caused by Adriamycin in Zebrafish Embryos and Larvae: A Novel Model of Glomerular Damage. PLoS ONE, 2014, 9, e98131.	2.5	24
8	Small Molecule Membrane Transporters in the Mammalian Podocyte: A Pathogenic and Therapeutic Target. International Journal of Molecular Sciences, 2014, 15, 21366-21380.	4.1	5
9	Warfarin-related nephropathy: possible role for the warfarin pharmacogenetic profile. CKJ: Clinical Kidney Journal, 2014, 7, 605-608.	2.9	14
10	ROLE OF COMPONENTS OF FRAILTY ON QUALITY OF LIFE IN DIALYSIS PATIENTS: A CROSSâ€6ECTIONAL STUDY. Journal of Renal Care, 2013, 39, 96-102.	1.2	11
11	Podocyte Expression of Membrane Transporters Involved in Puromycin Aminonucleoside-Mediated Injury. PLoS ONE, 2013, 8, e66159.	2.5	7
12	Podocyte Glutamatergic Signaling Contributes to the Function of the Glomerular Filtration Barrier. Journal of the American Society of Nephrology: JASN, 2009, 20, 1929-1940.	6.1	77
13	Proteomics of Plasma and Urine in Primary Nephrotic Syndrome in Children. , 2008, 160, 17-28.		12
14	HIV-1 Tat reduces nephrin in human podocytes: a potential mechanism for enhanced glomerular permeability in HIV-associated nephropathy. Aids, 2007, 21, 423-432.	2.2	39
15	Transitions of serum albumin in patients with glomerulosclerosis â€~in vivo' characterization by electrophoretic titration curves. Electrophoresis, 2006, 27, 2960-2969.	2.4	12
16	Direct effect of plasma permeability factors from patients with idiopatic FSGS on nephrin and podocin expression in human podocytes. International Journal of Molecular Medicine, 2005, 16, 49.	4.0	7
17	Glomerular Permeability Defect in Hypertension Is Dependent on Renin Angiotensin System Activation. American Journal of Hypertension, 2005, 18, 844-850.	2.0	9
18	Direct effect of plasma permeability factors from patients with idiopatic FSGS on nephrin and podocin expression in human podocytes. International Journal of Molecular Medicine, 2005, 16, 49-58.	4.0	18

MICHELE CARRARO

#	Article	IF	CITATIONS
19	The effect of proteinase inhibitors on glomerular albumin permeability induced in vitro by serum from patients with idiopathic focal segmental glomerulosclerosis. Nephrology Dialysis Transplantation, 2004, 19, 1969-1975.	0.7	15
20	Recurrent focal glomerulosclerosis in the era of genetics of podocyte proteins: theory and therapy. Nephrology Dialysis Transplantation, 2004, 19, 1036-1040.	0.7	28
21	Glomerular albumin permeability as anin vitromodel for characterizing the mechanism of focal glomerulosclerosis and predicting post-transplant recurrence. Pediatric Transplantation, 2004, 8, 339-343.	1.0	15
22	Apolipoprotein E in idiopathic nephrotic syndrome and focal segmental glomerulosclerosis. Kidney International, 2003, 63, 686-695.	5.2	23
23	Nephrotic urine prevents increased rat glomerular albumin permeability induced by serum from the same patient with idiopathic nephrotic syndrome. Nephrology Dialysis Transplantation, 2003, 18, 689-693.	0.7	14
24	Serum Glomerular Permeability Activity in Patients with Podocin Mutations (NPHS2) and Steroid-ResistantNephrotic Syndrome. Journal of the American Society of Nephrology: JASN, 2002, 13, 1946-1952.	6.1	77
25	Characterization of plasma factors that alter the permeability to albumin within isolated glomeruli. Proteomics, 2002, 2, 197-205.	2.2	43
26	PROTEINURIA IN FOCAL SEGMENTAL GLOMERULOSCLEROSIS: ROLE OF CIRCULATING FACTORS AND THERAPEUTIC APPROACH. Renal Failure, 2001, 23, 533-541.	2.1	15
27	Apolipoproteins Prevent Glomerular Albumin Permeability Induced In Vitro by Serum from Patients with Focal Segmental Glomerulosclerosis. Journal of the American Society of Nephrology: JASN, 2001, 12, 143-150.	6.1	57
28	Prevalence, Genetics, and Clinical Features of Patients Carrying Podocin Mutations in Steroid-Resistant Nonfamilial Focal Segmental Glomerulosclerosis. Journal of the American Society of Nephrology: JASN, 2001, 12, 2742-2746.	6.1	155
29	Prediction and treatment of recurrent focal segmental glomerulosclerosis after renal transplantation in children. American Journal of Kidney Diseases, 1999, 34, 1048-1055.	1.9	188
30	Relationship between Whole-Body Protein Turnover and Serum Creatinine in Chronically Uremic Patients. Mineral and Electrolyte Metabolism, 1998, 24, 267-272.	1.1	10
31	Nifedipine Reduces Postexercise Proteinuria in Young Volunteers. Kidney and Blood Pressure Research, 1995, 18, 306-310.	2.0	0
32	Involvement of the Renin Angiotensin System in the Pathogenesis of Postexercise Proteinuria. Scandinavian Journal of Urology and Nephrology, 1993, 27, 301-304.	1.4	14