

Ingvild Vik

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6676995/publications.pdf>

Version: 2024-02-01

10
papers

155
citations

1683934

5
h-index

1372474

10
g-index

10
all docs

10
docs citations

10
times ranked

186
citing authors

#	ARTICLE	IF	CITATIONS
1	Ibuprofen versus pivmecillinam for uncomplicated urinary tract infection in women – A double-blind, randomized non-inferiority trial. <i>PLoS Medicine</i> , 2018, 15, e1002569.	3.9	88
2	Ibuprofen versus mecillinam for uncomplicated cystitis - a randomized controlled trial study protocol. <i>BMC Infectious Diseases</i> , 2014, 14, 693.	1.3	16
3	Reducing antibiotic use in uncomplicated urinary tract infections in adult women: a systematic review and individual participant data meta-analysis. <i>Clinical Microbiology and Infection</i> , 2022, 28, 1558-1566.	2.8	11
4	Bacteriology in uncomplicated urinary tract infections in Norwegian general practice from 2001 – 2015. <i>BJGP Open</i> , 2018, 1, bjgpopen17X101145.	0.9	9
5	Pivmecillinam for Uncomplicated Lower Urinary Tract Infections Caused by <i>Staphylococcus saprophyticus</i> – Cumulative Observational Data from Four Recent Clinical Studies. <i>Antibiotics</i> , 2019, 8, 57.	1.5	7
6	Treatment of uncomplicated UTI in males: a systematic review of the literature. <i>BJGP Open</i> , 2021, 5, bjgpopen20X101140.	0.9	7
7	Strategies to reduce antibiotic use in women with uncomplicated urinary tract infection in primary care: protocol of a systematic review and meta-analysis including individual patient data. <i>BMJ Open</i> , 2020, 10, e035883.	0.8	6
8	Antibiotic switch after treatment with UTI antibiotics in male patients. <i>Infectious Diseases</i> , 2020, 52, 405-412.	1.4	5
9	Predictors of Symptom Duration and Bacteriuria in Uncomplicated Urinary Tract Infection. <i>Scandinavian Journal of Primary Health Care</i> , 2018, 36, 446-454.	0.6	4
10	Predicting the use of antibiotics after initial symptomatic treatment of an uncomplicated urinary tract infection: analyses performed after a randomised controlled trial. <i>BMJ Open</i> , 2020, 10, e035074.	0.8	2